



Swansea University
Prifysgol Abertawe



Cronfa - Swansea University Open Access Repository

This is an author produced version of a paper published in :
Physical Review D

Cronfa URL for this paper:
<http://cronfa.swan.ac.uk/Record/cronfa22986>

Paper:

Alston, S., Dunbar, D. & Perkins, W. (2015). n-point amplitudes with a single negative-helicity graviton. *Physical Review D*, 92(6)

<http://dx.doi.org/10.1103/PhysRevD.92.065024>

This article is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Authors are personally responsible for adhering to publisher restrictions or conditions. When uploading content they are required to comply with their publisher agreement and the SHERPA RoMEO database to judge whether or not it is copyright safe to add this version of the paper to this repository.

<http://www.swansea.ac.uk/iss/researchsupport/cronfa-support/>

September 1, 2015

n -point amplitudes with a single negative-helicity graviton

Sam D. Alston, David C. Dunbar and Warren B. Perkins

*College of Science,
Swansea University,
Swansea, SA2 8PP, UK
September 1, 2015*

Abstract

We construct an expression for the n -point one-loop graviton scattering amplitude with a single negative helicity external leg using an augmented recursion technique. We analyse the soft-limits of these amplitudes and demonstrate that they have soft behaviour beyond the conjectured universal behaviour.

PACS numbers: 04.65.+e

I. INTRODUCTION

Exploring the singularities of scattering amplitudes has been a key theme of research in recent years [1] with the aim being to compute amplitudes entirely from a knowledge of their singular behaviour. This reprise of earlier work [2] has been invigorated both by developments in the understanding of underlying symmetries [3] and technical progress.

One important technique has been BCFW [4] recursion which applies complex analysis to amplitudes. Using Cauchy's theorem, if a complex function is analytic except at simple poles z_i (all non-zero) and $f(z) \rightarrow 0$ as $|z| \rightarrow \infty$ then by considering the integral

$$\oint_C f(z) \frac{dz}{z} \quad (1.1)$$

where the contour C is the circle at infinity, we obtain

$$f(0) = - \sum_i \frac{\text{Residue}(f, z_i)}{z_i} \quad (1.2)$$

The recursion procedure for amplitudes utilises this by complexifying two of the external momenta¹ by shifting a pair of spinors

$$\bar{\lambda}_a \rightarrow \bar{\lambda}_{\hat{a}} = \bar{\lambda}_a - z \bar{\lambda}_b \quad , \quad \lambda_b \rightarrow \lambda_{\hat{b}} = \lambda_b + z \lambda_a. \quad (1.3)$$

This shifts the momenta p_a and p_b to complex values $p_a(z)$ and $p_b(z)$ which are both still null and preserves overall momentum conservation. The resultant amplitude $A(z)$ plays the role of $f(z)$ above.

As tree amplitudes are rational functions of λ_i and $\bar{\lambda}_i$, provided the complexified amplitude $A(z)$ satisfies the conditions above, $A(0)$ may be determined in terms of residues. These residues arise from simple poles corresponding to factorisations of the amplitude [5]. The target amplitude is thus readily expressed in terms of lower point amplitudes evaluated with specific complex momenta,

$$A_n^{\text{tree}}(0) = \sum_{i,\lambda} A_{r_i+1}^{\text{tree},\lambda}(z_i) \frac{i}{K^2} A_{n-r_i+1}^{\text{tree},-\lambda}(z_i), \quad (1.4)$$

where the summation over i is only over factorisations where the a and b legs are on opposite sides of the pole. This is the on-shell recursive expression of [4].

One-loop amplitudes in a massless theory can be expressed as [6]

$$A_n^{\text{1-loop}} = \sum_{i \in \mathcal{C}} a_i I_4^i + \sum_{j \in \mathcal{D}} b_j I_3^j + \sum_{k \in \mathcal{E}} c_k I_2^k + R_n + O(\epsilon), \quad (1.5)$$

where the I_r^i are r -point scalar integral functions and the a_i etc. are rational coefficients. R_n is a purely rational term. In terms of complex momentum this means the amplitude has both poles and discontinuities. Eq.(1.5) may simplify in specific theories, e.g. in maximal

¹ As usual, a null momentum is represented as a pair of two component spinors $p^\mu = \sigma_{\alpha\dot{\alpha}}^\mu \lambda^\alpha \bar{\lambda}^{\dot{\alpha}}$. For real momenta $\lambda = \pm \bar{\lambda}^*$ but for complex momenta λ and $\bar{\lambda}$ are independent [3]. We are using a spinor helicity formalism with the usual spinor products $\langle a b \rangle = \epsilon_{\alpha\beta} \lambda_a^\alpha \lambda_b^\beta$ and $[a b] = -\epsilon_{\dot{\alpha}\dot{\beta}} \bar{\lambda}_a^{\dot{\alpha}} \bar{\lambda}_b^{\dot{\beta}}$.

supersymmetric Yang-Mills and supergravity where only the first "box functions" appear [7, 8] and in specific helicity amplitudes such as the "all-plus" and "single-minus" amplitudes of both Yang-Mills and gravity where only the rational terms R_n appear.

The appearance of discontinuities is in fact enormously helpful in computing amplitudes. These discontinuities allow the direct computation of the coefficients of the scalar integral functions using unitarity methods. Dividing the amplitude into integral functions with rational coefficients has been very fruitful: a range of specialised techniques have been devised to determine the rational coefficients based on unitarity techniques rather than Feynman diagrams [6, 7, 9–11]. Progress has been made both via the two-particle cuts [6, 7, 12] and using generalisations of unitarity [10] where, for example, triple [13–16] and quadruple cuts [11] are utilised to identify the triangle and box coefficients respectively.

Returning to the amplitude, there is a further problem related to complex factorisation. In general, beyond tree level, amplitudes with complex momenta may have multi-pole singularities and consequently may have poles of higher order. Mathematically, this is not a barrier to using complex analysis, since, if we have a function whose expansion about z_i is

$$f(z) = \frac{a_{-2}}{(z - z_i)^2} + \frac{a_{-1}}{(z - z_i)} + \text{finite} \quad (1.6)$$

then

$$\text{Residue}\left(\frac{f(z)}{z}, z_i\right) = -\frac{a_{-2}}{z_i^2} + \frac{a_{-1}}{z_i} \quad (1.7)$$

However for one-loop amplitudes only the leading singularities have been determined in general and there are no general theorems for the sub-leading terms.

In one-loop amplitudes poles in momenta arise in two ways: firstly from explicit poles in Feynman diagrams such as in fig. 1, and secondly from loop momentum integrals. Specifically a $(P^2)^{-1}$ pole can arise from the loop momentum integral of diagrams of the form shown in fig. 2.




FIG. 1: Diagrams with an explicit pole.




FIG. 2: Diagrams with a pole from loop integration.

When the pole is a two-particle factorisation $P^2 = (k_a + k_b)^2$, these two sources can overlap as in fig. 3 and produce double poles.




FIG. 3: Double poles may arise from these diagrams.

Figure 3 illustrates the challenge in determining the sub-leading contribution. Although the poles are physical and gauge independent, the diagram effectively contains an off-shell current whose sub-leading term in s_{ab} is gauge and scheme dependent. The leading term can be shown to be expressible in terms of an "effective vertex" times an on-shell tree amplitude [17, 18]. The vertex vanishes unless both legs a and b have the same helicity in which case the double pole is

$$V(a^+, b^+, P^+) \times \frac{1}{s_{ab}} \times A^{\text{tree}}(P^-, \dots) \quad (1.8)$$

The tree amplitude in this expression is the lower point amplitude where legs a and b have been replaced by a single leg of negative helicity. In Yang-Mills theory the effective vertex takes the form

$$V^{YM}(a^+, b^+, P^+) = -\frac{i}{48\pi^2} \frac{[ab][bP][Pa]}{s_{ab}} \propto \frac{[ab]}{\langle ab \rangle} \quad (1.9)$$

Note that this is only singular for complex momentum where we can have $\langle ab \rangle = 0$ without necessarily having $[ab] = 0$. For graviton scattering amplitudes the equivalent effective vertex is

$$V^{\text{grav}}(a^+, b^+, P^+) = \frac{i}{360\pi^2} \frac{([ab][bP][Pa])^2}{s_{ab}} \quad (1.10)$$

In this article we develop techniques to determine the sub-leading pole in the amplitude. Double poles generally are present in most non-supersymmetric amplitudes but the starting point to study these is in the simpler purely rational amplitudes. At one-loop level these are the all-plus amplitude $M_n(+, +, +, \dots, +)$ and the single-minus amplitude $M_n(-, +, +, \dots, +)$. These two amplitudes are vanishing at tree level. Of these the all-plus amplitude does not have double poles since the leading pole will multiply a vanishing tree amplitude. The single-minus amplitude however does have double poles in s_{bc} with b, c positive helicity legs. The leading term has a MHV tree amplitude as a factor. For Yang-Mills the single-minus all- n forms are known [19] and in ref. [17, 20] a general form for the sub-leading was presented for this specific amplitude. The equivalent gravity amplitude has explicit forms for four [21], five and six-points [22] and but as yet no conjectured all- n sub-leading behaviour.

In this article we will isolate the sub-leading term which is not present in the explicit on-shell factorisation and give expressions for this extra term. This allows a recursive expression for the all- n amplitude to be given. The form of this is not particularly compact but it is explicit.

As an application we examine the soft-limits of the amplitude. Gravity amplitudes obey

soft-factorisation theorems at tree level [23–25]. Motivated by the presence of BMS symmetry [26], these are conjectured to also apply at loop level [25]. We confirm the result, obtained for $n \leq 6$ [27, 28], that one-loop amplitudes do not obey the soft-factorisation theorems.

II. SINGLE-MINUS AMPLITUDE COMPUTATION

We now turn to the explicit computation of the n -point single-minus amplitude in gravity $M_n(a^-, b^+, \dots, n^+)^2$.

We utilise a shift on the negative helicity leg, a , and one of the positive helicity legs b ,

$$\bar{\lambda}_a \rightarrow \bar{\lambda}_{\hat{a}} = \bar{\lambda}_a - z\bar{\lambda}_b, \quad , \quad \lambda_b \rightarrow \lambda_{\hat{b}} = \lambda_b + z\lambda_a \quad (2.1)$$

We will assume, to be justified later, that under this shift $M(z) \rightarrow 0$ as $z \rightarrow \infty$ so that $M(0)$ may be determined by its residues. This shift appears to have good asymptotic behaviour, $M \sim z^{-2}$ for $n \leq 7$, but excites double poles.

Under the shift (2.1) the factorisation channels of $M^{1\text{-loop}}(\hat{a}^-, \hat{b}^+, c^+, \dots, n^+)$ can be arranged into terms which arise from factorisations into products of on-shell tree and loop amplitudes and an extra contribution which must be computed separately.

The on-shell factorisations fall into three sets of diagrams which we label A, B and C together with the extra term labeled Δ_n . The full amplitude is the sum of the contributions of each type:

$$\begin{aligned} M^{1\text{-loop}}(a^-, b^+, c^+, \dots, n^+) = & A_{\text{multi}}(a^-, b^+, c^+, \dots, n^+) + B_{\text{3ptMHV}}(a^-, b^+, c^+, \dots, n^+) \\ & + C_{\text{3ptMHV}}(a^-, b^+, c^+, \dots, n^+) + \Delta_n(a^-, b^+, c^+, \dots, n^+). \end{aligned} \quad (2.2)$$

The first three terms may be written down straightforwardly in terms of shifted lower point amplitudes. Firstly,

$$A_{\text{multi}}(a^-, b^+, c^+, \dots, n^+) = \sum_{\text{Part}} M_{n+2-r}^{\text{tree}}(\hat{a}^-, \hat{P}^-, \{R\}) \frac{1}{P_{a, \{R\}}^2} M_r^{1\text{-loop}}(-\hat{P}^+, \hat{b}^+, \{\bar{R}\}) \Big|_{\hat{P}^2=0} \quad (2.3)$$

where the sum is over all distinct partitions of $\{c, \dots, n\}$ into two sets $\{R\}$ and $\{\bar{R}\}$ which each contain at least two members. The on-shell amplitudes are evaluated at the value of z such that $\hat{P}^2 = 0$. When the set $\{R\}$ contains a single element we have the second type of term,

$$B_{\text{3ptMHV}}(a^-, b^+, c^+, \dots, n^+) = \sum_{\gamma \in \{c, \dots, n\}} M_3^{\text{tree}}(\hat{a}^-, \hat{P}^-, \gamma^+) \frac{1}{s_{a\gamma}} M_{n-1}^{1\text{-loop}}(-\hat{P}^+, \hat{b}^+, c^+, \dots, \not{\gamma}, \dots, n^+) \Big|_{[\hat{a}\gamma]=0} \quad (2.4)$$

where $\not{\gamma}$ denotes that leg γ is omitted from the argument list. When the set $\{\bar{R}\}$ contains a

² We use the normalisation for the full physical amplitudes $\mathcal{M}^{\text{tree}} = i(\kappa/2)^{n-2} M^{\text{tree}}$, $\mathcal{M}^{1\text{-loop}} = i(2\pi)^{-2} (\kappa/2)^n M^{1\text{-loop}}$.

single element there are also the factorisations,

$$C_{3\text{pt}\overline{\text{MHV}}}(a^-, b^+, c^+, \dots, n^+) = \sum_{\gamma \in \{c, \dots, n\}} M_3^{\text{tree}}(-\hat{P}^-, \hat{b}^+, \gamma^+) \frac{1}{s_{b\gamma}} M_{n-1}^{1\text{-loop}}(\hat{a}^-, \hat{P}^+, c^+, \dots, \cancel{\gamma}, \dots, n^+) \Big|_{\langle \hat{b}\gamma \rangle = 0} \quad (2.5)$$

These factorisations are illustrated in figs 4-6.




FIG. 4: Factorisations of type A. $\{R\} \cup \{\bar{R}\} = \{c, \dots, n\}$ and both contain at least two elements.




FIG. 5: Factorisation of type B.




FIG. 6: Factorisation of type C.

Note that these factorisations require a knowledge of both the all-plus and single-minus lower point amplitudes. Only the $n-1$ point single-minus amplitude appears.

The remaining contribution can be determined [22, 29] by considering diagrams of the form of fig 7. We use an axial gauge formalism [30, 31] in which helicity labels can be used for internal lines and off-shell internal legs in the vertices are nullified using a reference spinor: given a reference null momentum η , any off-shell leg with momentum K can be *nullified* using

$$K^b = K - \frac{K^2}{[\eta|K|\eta]}\eta \quad (2.6)$$

which gives spinors

$$\lambda_K = \alpha K|\eta], \quad \bar{\lambda}_K = \alpha^{-1} \frac{K|\eta\rangle}{[\eta|K|\eta]}. \quad (2.7)$$

Such a formalism has not been completely specified for gravity, however we only need a few simple properties of the three point vertex, specifically

$$V_3^{grav}(\alpha^+, \beta^+, \gamma^-) = (V_3^{Y-M}(\alpha^+, \beta^+, \gamma^-))^2 = \left(\frac{[\alpha \beta] \langle \gamma \eta \rangle^2}{\langle \alpha \eta \rangle \langle \beta \eta \rangle} \right)^2. \quad (2.8)$$

We choose $\lambda_\eta = \lambda_a$ and $\bar{\lambda}_\eta = \bar{\lambda}_b$ to simplify the computation.




FIG. 7: Non-factorising contribution arising along with the double poles.

In fig. 7

$$B = \ell + \hat{b} \quad C = -\ell + \gamma \quad (2.9)$$

so that

$$B + C = \hat{b} + \gamma \quad (2.10)$$

and so although B and C are loop momentum dependent $B + C$ is not and $s_{BC} = s_{\hat{b}\gamma}$.

The diagram shown in fig. 7 gives a contribution

$$\int d^D \ell \left(\frac{[b|\ell|a][\gamma|\ell|a] \langle Ca \rangle^2}{\langle ba \rangle \langle \gamma a \rangle \langle Ba \rangle^2} \right)^2 \frac{1}{\ell^2 B^2 C^2} \tau(B^-, C^+, g^+ \dots n^+, \hat{a}^-, c^+, \dots, f^+) \quad (2.11)$$

which involves the current $\tau(B^-, C^+, g^+ \dots n^+, \hat{a}^-, c^+, \dots, f^+)$ with two off-shell legs B and C . Fortunately, as we are only interested in the residues generated by these diagrams we do not need τ exactly, only its leading and sub-leading behaviour at the pole. The loop integration introduces a factor of $s_{\hat{b}\gamma}$ for each s_{BC} , B^2 or C^2 appearing in τ , thus we can consider the expansion of τ in these quantities. To see how this arises, we perform a standard Feynman parametrisation of the loop propagators:

$$\frac{1}{\ell^2 B^2 C^2} \rightarrow 2 \int_0^1 dx_1 dx_2 dx_3 \delta(x_1 + x_2 + x_3 - 1) \frac{1}{[(\ell + x_2 \hat{b} - x_3 \gamma)^2 + x_2 x_3 s_{\hat{b}\gamma}]^3} \quad (2.12)$$

and make the change of variables:

$$\begin{aligned} \ell &= p - x_2 \hat{b} + x_3 \gamma \\ B &= p + (1 - x_2) \hat{b} + x_3 \gamma \\ C &= -p - x_2 \hat{b} - (1 - x_3) \gamma \end{aligned} \quad (2.13)$$

leading to

$$\frac{1}{\ell^2 B^2 C^2} = 2 \int_0^1 dx_1 dx_2 dx_3 \delta(x_1 + x_2 + x_3 - 1) \frac{1}{[p^2 + x_2 x_3 s_{\hat{b}\gamma}]^3} \quad (2.14)$$

In the rest of the integrand the change of variables replaces s_{BC} , B^2 and C^2 with bilinear combinations of the loop momentum, \hat{b} and γ . In dimensional regularization the standard tensor integral is,

$$\int \frac{d^D \ell}{(2\pi)^{2w}} \frac{\ell^{a_1} \dots \ell^{a_n}}{(\ell^2 + 2Q \cdot \ell + M^2)^A} = \frac{(-1)^n \Gamma(A - D/2)}{(4\pi)^w \Gamma(A)} \left[\frac{Q^{a_1} \dots Q^{a_n}}{(Q^2 - M^2)^{A-D/2}} + \mathcal{O}\left(\frac{1}{(Q^2 - M^2)^{A-D/2-1}}\right) \right]. \quad (2.15)$$

In this case $Q = 0$ and $M^2 = s_{\hat{b}\gamma}$, thus the leading term has a factor of $s_{\hat{b}\gamma}^{-1}$ and any product of two loop momentum factors in the integrand will generate a factor of $s_{\hat{b}\gamma}$. With this in mind, each factor of s_{BC} , B^2 or C^2 in the numerator of τ will ultimately introduce a factor of $s_{\hat{b}\gamma}$ which reduces the order of any pole.

III. GRAVITY CURRENTS

In order to determine (2.11) we would ideally use the full gravity MHV current with two massive legs,

$$\tau_{\text{grav}}^{\text{MHV}}(a^-, B^-, C^+, d^+, \dots, n^+) \quad (3.1)$$

where B and C are non-null, however this is not available. Fortunately we only need the leading and sub-leading terms in the current around $B^2 = C^2 = s_{BC} = 0$. In this region we may use an approximation to the current which arises from the Kawai Lewellen and Tye (KLT) relations for on-shell amplitudes [32] which we continue (sufficiently) off-shell. While the KLT relations only hold exactly on-shell, it turns out that we can use them to compute the pole we require from YM currents. The justification for this is explicitly given in the appendix and by the fact it generates a consistent amplitude.

The explicit form for the KLT relations used is [33]

$$M_n(B, C, \dots, n, \hat{a}) = i(1)^{n+1} \left[A_n^{\text{YM}}(B, C, \dots, n, \hat{a}) \times \sum_{\mathcal{P}_1, \mathcal{P}_2} f(i_1, \dots, i_k; B) \bar{f}(j_1, \dots, j_{k'}) A_n^{\text{YM}}(i_1, \dots, i_k, B, n, j_1, \dots, j_{k'}, \hat{a}) \right] + \mathcal{P}_{\text{KLT}}(C, d, \dots, n-1), \quad (3.2)$$

where \mathcal{P}_{KLT} represents a permutation over the $n-3$ legs $\{C, \dots, n-1\}$. These $n-3$ legs are split into two subsets; for even n , $\{C, \dots, n/2\}$ and $\{n/2+1, \dots, n-1\}$. The \mathcal{P}_1 are the permutations $\{i_1, \dots, i_k\}$ of $\{C, \dots, n/2\}$, \mathcal{P}_2 are the permutations $\{j_1, \dots, j_{k'}\}$ of $\{n/2+1, \dots, n-1\}$, $k = n/2 - 1$ and $k' = n/2 - 2$. For n odd, $n/2$ is replaced by $(n+1)/2$. The KLT expression is extremely useful, however the number of terms grows rapidly with n . For n even the total number of terms is

$$(n-3)! \times \left(\frac{n}{2} - 1\right)! \times \left(\frac{n}{2} - 2\right)! \quad (3.3)$$

The functions f and \bar{f} are given by

$$\begin{aligned} f(i_1, \dots, i_k; B) &= s_{Bi_k} \prod_{m=1}^{k-1} \left(s_{Bi_m} + \sum_{p=m+1}^k g(i_m, i_p) \right) \\ \bar{f}(j_1, \dots, j_{k'}) &= s_{j_1 n} \prod_{m=2}^{k'} \left(s_{j_m n} + \sum_{p=1}^{m-1} g(j_p, j_m) \right) \end{aligned} \quad (3.4)$$

where

$$g(i, j) = \begin{cases} s_{ij} & \text{if } i > j \\ 0 & \text{otherwise} \end{cases} \quad (3.5)$$

The inequality above is interpreted in terms of the ordering of the legs in the original gravity amplitude, i.e. $i > j$ means leg i is to the right of leg j in the argument list of M_n in (3.2). This definition is applied to the term of \mathcal{P}_{KLT} shown explicitly in (3.2) and the permutation \mathcal{P}_{KLT} is applied to the resulting function.

The particular form of the KLT relation has been chosen to help us isolate the double pole term and allow us to compute the subleading term from this. In appendix A it is shown explicitly that a quantity τ_n^g that captures the leading and sub-leading behaviour of τ_{grav}^{MHV} can be obtained from the KLT relations (3.2) by replacing the Yang-Mills amplitudes by Yang-Mills currents. When legs B and C are adjacent the Yang-Mills current is

$$\begin{aligned} \tau_n^{\text{YM}}(\hat{a}^-, b^+, \dots, f^+, B^-, C^+, g^+, \dots, n^+) &= \frac{\langle Ba \rangle^2}{\langle Ca \rangle^2} \frac{1}{\langle ab \rangle \dots \langle ef \rangle} \frac{1}{\langle gh \rangle \dots \langle na \rangle} \\ &\times \left(\frac{\langle aB \rangle \langle aC \rangle \langle ag \rangle \langle fa \rangle}{\langle fB \rangle \langle Cg \rangle \langle gf \rangle} + \frac{\langle aB \rangle \langle af \rangle^2}{\langle Bf \rangle \langle gf \rangle} \frac{[\eta|C|a]}{[\eta|B+C|f]} + \frac{\langle aC \rangle \langle ag \rangle^2}{\langle Cg \rangle \langle gf \rangle} \frac{[\eta|B|a]}{[\eta|B+C|g]} \right. \\ &\left. - \frac{\langle a|BC|a \rangle}{s_{BC}} \frac{[\eta|B+C|a]^2}{[\eta|B+C|f][\eta|B+C|g]} \right) + \mathcal{O}(B^2) + \mathcal{O}(C^2) \end{aligned} \quad (3.6)$$

For the special case $\tau_n^{\text{YM}}(\hat{a}^-, B^-, C^+, g^+, \dots, n^+)$ we have

$$\begin{aligned} \tau_n^{\text{YM}}(\hat{a}^-, B^-, C^+, g^+, \dots, n^+) &= \frac{\langle Ba \rangle^2}{\langle Ca \rangle^2} \frac{1}{\langle gh \rangle \dots \langle na \rangle} \\ &\times \left(\frac{\langle aC \rangle \langle ga \rangle}{\langle Cg \rangle} \frac{[\eta|B|a]}{[\eta|B+C|g]} - \frac{\langle a|BC|a \rangle}{s_{BC}} \frac{[\eta|B+C|a]}{[\eta|B+C|g]} \right) + \mathcal{O}(B^2) + \mathcal{O}(C^2) \end{aligned} \quad (3.7)$$

When B and C are separated the current reduces to the Parke-Taylor form.

As discussed above, one factor of $s_{b\gamma}^{-1}$ arises from the integration, so double poles in the amplitude arise when single powers of s_{BC}^{-1} occur in τ_n^g . While there appear to be double poles in τ_n^g when \mathcal{P}_{KLT} leaves C adjacent to B in the first Yang-Mills factor and $i_k = C$ in the second, there is a factor of $s_{BC} = s_{b\gamma}$ in f which lowers the order of the pole in these cases. Furthermore, in these terms with legs B and C adjacent in both Yang-Mills amplitudes the loop momentum dependent factors of f have a restricted form. Consider first the term in \mathcal{P}_{KLT} explicitly shown in (3.2) when $i_k = C$. In this case f has no loop momentum dependence. Other than B (which is not involved in the permutations), C is the left most leg in the argument list of τ_n^g , therefore when $i_k = C$, the final term in the sum: $\sum_{p=m+1}^k g(i_m, i_p)$

is always $g(i_m, C) = s_{i_m C}$. All of the factors in the product part of f therefore contain:

$$s_{B i_m} + s_{C i_m} = B^2 + C^2 + [i_m | B + C | i_m] = B^2 + C^2 + [i_m | \hat{b} + \gamma | i_m]. \quad (3.8)$$

As \mathcal{P}_{KLT} doesn't involve B , any of these permutations that leave C alone leave both B and C untouched and the arguments above are unaffected. Those elements of \mathcal{P}_{KLT} that move C around separate B and C in the first Yang-Mills factor and hence are not in this class.

An s_{BC} factor also arises when legs B and C are adjacent in just one of the Yang-Mills factors, however the loop momentum dependence of the $f\bar{f}$ factors in these is not always simple.

Crucially the only pole in τ_n^g occurs in the terms with B and C adjacent in both Yang-Mills factors, hence τ_n^g only needs to be determined beyond leading order when B and C are adjacent in both Yang-Mills factors.

The KLT based expression for τ_n^g can now be used to evaluate the non-factorising contributions to the amplitude: Δ_n . Using the explicit form of the KLT relations (3.2), the non-factorising diagram depicted in fig. 7 gives,

$$\begin{aligned} \Delta_n^{\text{pre-res}} = & i(-1)^{n+1} \int d^D \ell \left(\frac{[b|\ell|a][\gamma|\ell|a] \langle Ca \rangle^2}{\langle ba \rangle \langle \gamma a \rangle \langle Ba \rangle^2} \right)^2 \frac{1}{\ell^2 B^2 C^2} \\ & \times \left[\tau_n^{\text{YM}}(B, C, \dots, n, a) \sum_{\mathcal{P}_1, \mathcal{P}_2} f(i_1, \dots, i_k; B) \bar{f}(j_1, \dots, j_{k'}; C) \tau_n^{\text{YM}}(i_1, \dots, i_k, B, n, j_1, \dots, j_{k'}, a) \right] \\ & + \mathcal{P}_{KLT}(C, d, \dots, n-1) \end{aligned} \quad (3.9)$$

where $\Delta_n^{\text{pre-res}}$ denotes that the residue has yet to be extracted. Within the summation there are four distinct cases that are treated separately. These are distinguished by legs B and C being adjacent or separated in the two Yang-Mills currents. We denote the contributions to the final amplitude from terms of each type by Δ_{SS} , Δ_{AS} , Δ_{SA} and Δ_{AA} .

For terms where legs B and C are separated in both Yang-Mills currents, the Yang-Mills currents are finite at $\langle \hat{b}\gamma \rangle = 0$ and they are simply the Parke-Taylor amplitudes. The factor

$$\left(\frac{\langle Ca \rangle^2}{\langle Ba \rangle^2} \right)^2 \tau_n^{\text{YM}}(B, \dots, C, \dots, n, a) \tau_n^{\text{YM}}(i_1, \dots, i_k, B, n, j_1, \dots, j_{k'}, a) \quad (3.10)$$

then carries no net spinor weight in C or B and does not contain either $\bar{\lambda}_B$ or $\bar{\lambda}_C$. Using

$$\frac{\langle CX \rangle}{\langle CY \rangle} = \frac{\langle CX \rangle \langle \gamma a \rangle}{\langle CY \rangle \langle \gamma a \rangle} = \frac{\langle Ca \rangle \langle \gamma X \rangle + \langle C\gamma \rangle \langle Xa \rangle}{\langle Ca \rangle \langle \gamma Y \rangle + \langle C\gamma \rangle \langle Ya \rangle} = \frac{\langle \gamma X \rangle}{\langle \gamma Y \rangle} + \mathcal{O}(\langle C\gamma \rangle) \quad (3.11)$$

λ_C can be replaced by λ_γ in this factor. λ_B could similarly be replaced by $\lambda_{\hat{b}}$. As there is only a single pole in these contributions, the residue simply involves this factor evaluated at $\langle \hat{b}\gamma \rangle = 0$. Since this factor contains no net spinor weight in B , in the residue λ_B is ultimately replaced by λ_γ .

As discussed above, the only contributions to the residue come from scalar integrals, therefore within the f and \bar{f} factors terms involving B and C are substituted as:

$$s_{CX} \rightarrow (1 - x_3) s_{\gamma X} + x_2 s_{\hat{b}X} \quad s_{BX} \rightarrow x_3 s_{\gamma X} + (1 - x_2) s_{\hat{b}X} \quad (3.12)$$

The residue again involves these quantities evaluated on the pole where

$$s_{CX} \rightarrow (1 - x_3)s_{\gamma X} + x_2[X|b|a] \frac{\langle \gamma X \rangle}{\langle \gamma a \rangle} \quad s_{BX} \rightarrow x_3s_{\gamma X} + (1 - x_2)[X|b|a] \frac{\langle \gamma X \rangle}{\langle \gamma a \rangle} \quad (3.13)$$

After making these substitutions, the $f\bar{f}$ factor in each term in this class can be expanded:

$$x_2x_3[f\bar{f}]_i := \sum_r \sum_v H_{i:rv}^{SS} x_2^r x_3^v \quad (3.14)$$

where i labels the different permutations in this class. The contribution of term i to Δ_{SS} is then

$$\begin{aligned} \Delta_{SS:i} = i(-1)^{n+1} \frac{[b\gamma]^4}{s_{b\gamma}} \sum_r \sum_v H_{i:rv}^{SS} \frac{r!v!}{(r+v+2)!} \\ \times \left[A_n^{\text{YM}}(\gamma^-, \dots, \gamma^+, \dots, n, a) A_n^{\text{YM}}(i_1, \dots, \gamma^+, \dots, i_k, \gamma^-, n, j_1, \dots, j_{k'}, a) \right] \end{aligned} \quad (3.15)$$

where the γ 's in the argument lists of the Yang-Mills amplitudes appear where B and C appeared originally.

As discussed above, when legs B and C are adjacent in either of the Yang-Mills currents in (3.9), $f\bar{f}$ always contains a factor of s_{BC} . If legs B and C are adjacent in just one of the currents, there is a single pole overall, only the singular part of the current with B and C adjacent contributes and the Parke-Taylor form can be used for the other. Again we are dealing with a simple pole, so in the residue all shifted quantities are evaluated on the pole.

If legs B and C are adjacent in only the first Yang-Mills current, $\tau_n^{\text{YM}}(\hat{a}, B, C, g, \dots, n)$, the Parke-Taylor amplitude is used for the second current with the same substitutions as in the separated-separated case. The Feynman parameter integrations can also be performed in a similar fashion: after making the substitutions (3.13), $f\bar{f}$ is again expanded for each term in the class:

$$x_2x_3(x_3 + x_2 - 1)[f\bar{f}]_i := \sum_r \sum_v H_{i:rv}^{AS} x_2^r x_3^v \quad (3.16)$$

To help track the leg orderings, the string of $\langle xy \rangle$ factors in the denominator of (3.6) is denoted

$$F_n^{\text{YM}}(a^-, b^+, \dots, f^+, \mathcal{B}^-, \mathcal{C}^+, g^+, \dots, n^+) = \frac{1}{\langle ab \rangle \dots \langle ef \rangle} \frac{1}{\langle gh \rangle \dots \langle na \rangle} \quad (3.17)$$

where \mathcal{B}^- and \mathcal{C}^+ denote the positions of these legs in the original current and the slashes highlight that the function does not depend on the values of B and C . g denotes the leg following C in the first Yang-Mills current. The contribution of each term in this class to Δ_{AS} is

$$\begin{aligned} \Delta_{AS:i} = i(-1)^{n+1} \frac{[b\gamma]^4 \langle a|b\gamma|a \rangle}{s_{b\gamma}} \sum_r \sum_v H_{i:rv}^{AS} \frac{r!v!}{(r+v+2)!} \\ \times \left[\frac{\langle \gamma a \rangle}{\langle \gamma g \rangle} F_n^{\text{YM}}(a, \mathcal{B}, \mathcal{C}, g, \dots, n) A_n^{\text{YM}}(i_1, \dots, \gamma^+, \dots, i_k, \gamma^-, n, j_1, \dots, j_{k'}, a) \right] \end{aligned} \quad (3.18)$$

When legs B and C are adjacent in only the second Yang-Mills current, $\tau_n^{\text{YM}}(i_1, \dots, g, C, B, n, j_1, \dots, j_{k'}, \hat{a})$, the analogous definitions are

$$x_2 x_3 (x_3 + x_2 - 1) [f \bar{f}]_i := \sum_r \sum_v H_{rv}^{i:SA} x_2^r x_3^v \quad (3.19)$$

and leg g is the leg appearing before C in the second Yang-Mills current. The contribution of each term in this class to Δ_{SA} is

$$\begin{aligned} \Delta_{SA:i} = & i(-1)^{n+1} \frac{[b\gamma]^4 \langle a|b\gamma|a \rangle}{s_{b\gamma}} \sum_r \sum_v H_{rv}^{i:SA} \frac{r!v!}{(r+v+2)!} \\ & \times \left[A_n^{\text{YM}}(a, \gamma^-, \dots, \gamma^+, \dots, n) \frac{\langle \gamma a \rangle^2}{\langle \gamma g \rangle \langle \gamma n \rangle} F_n^{\text{YM}}(i_1, \dots, g, \mathcal{C}, \mathcal{B}, n, j_1, \dots, j_{k'}, a) \right] \end{aligned} \quad (3.20)$$

Finally there are contributions from terms where legs B and C are adjacent in both Yang-Mills currents. The form (3.6) is used for both currents. As the $f\bar{f}$ piece of these terms always have a factor of $s_{b\gamma}$ there is no pole in the terms involving the product of the regular parts of the two currents, a single pole in the terms where the singular part of one current multiplies the regular piece of the other (we denote these contributions $\Delta_{A_r A_s}$ and $\Delta_{A_s A_r}$) and a double pole in the product of the two singular parts (we denote these contributions $\Delta_{A_s A_s}$).

Note that, in this particular case, the loop momentum dependence of $f\bar{f}$ has the very restricted form given in (3.8). The pieces involving B^2 and C^2 cancel the corresponding propagator giving massless bubbles which are discarded in the usual dimensional regularisation prescription. The surviving piece of $f\bar{f}$ has no loop momentum dependence and thus no x dependence. Although not loop momentum dependent, we must expand $f\bar{f}$ about the pole. This corresponds to substituting

$$s_{CX} \longrightarrow s_{\gamma X} \quad s_{BX} \longrightarrow [X|b|a] \frac{\langle \gamma X \rangle}{\langle \gamma a \rangle} + \Delta z [b|X|a] \quad (3.21)$$

Then

$$\frac{[f\bar{f}]}{s_{BC}} = [f\bar{f}]_0 + \Delta z [f\bar{f}]_1 + O(\Delta z^2) \quad (3.22)$$

Schematically expanding the Yang-Mills currents and $f\bar{f}$ factors in terms of $s_{b\gamma}$, the adjacent-adjacent contribution has the form

$$\left(\frac{\tau_{-1}}{s_{b\gamma}} + \tau_0 \right) s_{b\gamma} ([f\bar{f}]_0 + s_{b\gamma} [f\bar{f}]_1) \left(\frac{\tau_{-1}}{s_{b\gamma}} + \tau_0 \right) \quad (3.23)$$

which gives the pole term

$$\frac{1}{s_{b\gamma}} \tau_{-1} \times [f\bar{f}]_0 \times \tau_{-1} \quad (3.24)$$

along with finite pieces

$$\tau_{-1} \times [f\bar{f}]_1 \times \tau_{-1} + \tau_{-1} \times [f\bar{f}]_0 \times \tau_0 + \tau_0 \times [f\bar{f}]_0 \times \tau_{-1} \quad (3.25)$$

Extracting a common factor of

$$\begin{aligned}
C_0 &= -\frac{[b\gamma]^4 \langle a|b\gamma|a \rangle}{360 s_{b\gamma}} \times F_n[a, \mathcal{B}, \mathcal{C}, g, \dots, n] \times F_n[i_1, \dots, f, \mathcal{C}, \mathcal{B}, n, j_1, \dots, j_k, a] \\
&= -\frac{[b\gamma]^4 \langle a|b\gamma|a \rangle \langle a\gamma \rangle \langle \gamma g \rangle \langle f\gamma \rangle \langle \gamma n \rangle}{360 \langle a\gamma \rangle^8 s_{b\gamma}} \\
&\quad \times A_{n-1}[a^-, \gamma^-, g, \dots, n] \times A_{n-1}[i_1, \dots, f, \gamma^-, n, j_1, \dots, j_k, a^-]
\end{aligned} \tag{3.26}$$

the adjacent-adjacent contribution is

$$\Delta_{AA:i} = C_0 \times \left(\frac{J_{r,r}}{s_{b\gamma}} [f\bar{f}]_0 + J_{r,r}^1 [f\bar{f}]_1 + (J_{r,s} + J_{s,r}) [f\bar{f}]_0 \right) \tag{3.27}$$

where

$$J_{r,s} = \frac{\langle \gamma a \rangle^4 \langle a g \rangle}{\langle \gamma g \rangle^2 \langle \gamma f \rangle \langle \gamma n \rangle} \tag{3.28}$$

$$J_{s,r} = \left(\frac{\langle \gamma a \rangle^3}{\langle \gamma g \rangle \langle f n \rangle} \right) \times \left(3 \frac{\langle a f \rangle \langle n a \rangle}{\langle n \gamma \rangle \langle \gamma f \rangle} - 2 \frac{\langle a n \rangle^2}{\langle n \gamma \rangle^2} - \frac{\langle a f \rangle^2}{\langle f \gamma \rangle^2} \right) \tag{3.29}$$

$$J_{r,r} = \frac{\langle \gamma a \rangle^3 \langle a|b\gamma|a \rangle}{\langle \gamma g \rangle \langle \gamma n \rangle \langle \gamma f \rangle} \tag{3.30}$$

$$J_{r,r}^1 = \frac{\langle \gamma a \rangle^2 \langle a|b\gamma|a \rangle}{[b\gamma] \langle \gamma g \rangle \langle \gamma n \rangle \langle \gamma f \rangle} \tag{3.31}$$

The full non-factorising contribution is then

$$\Delta_n(a^-, b^+, c^+, \dots, n^+) = \sum_i \Delta_{SS:i} + \sum_i \Delta_{AS:i} + \sum_i \Delta_{SA:i} + \sum_i \Delta_{AA:i} \tag{3.32}$$

where each sum is over all terms in the relevant class in (3.9).

We have used these expressions to generate the single-minus amplitudes for $n \leq 8$: our expressions have the correct symmetries and collinear limits as detailed in Appendix C.

IV. SOFT LIMITS

Graviton scattering amplitudes are singular as a leg becomes soft. Weinberg [23] many years ago presented the leading soft limit. If we parameterise the momentum of the n -th leg as $k_n^\mu = t \times k_s^\mu$ then in the limit $t \rightarrow 0$ the singularity in the n -point amplitude is

$$M_n \rightarrow \frac{1}{t} \times S^{(0)} \times M_{n-1} + O(t^0) \tag{4.1}$$

where M_{n-1} is the $n-1$ point amplitude. The soft-factor $S^{(0)}$ is universal and Weinberg showed that (4.1) does not receive corrections in loop amplitudes.

Recently it has also been proposed [24, 25] that the sub-leading and sub-sub leading terms are also universal. This can be best exposed, when a positive helicity leg becomes soft, by

setting

$$\lambda_n = t \times \lambda_s, \quad \bar{\lambda}_n = \bar{\lambda}_s, \quad (4.2)$$

In the $t \rightarrow 0$ limit the amplitude now has t^{-3} singularities. Effectively the choice (4.2) introduces a t^{-2} factor into the polarisation tensor $\epsilon_{\mu\nu}^+$ and a t^{+2} factor into $\epsilon_{\mu\nu}^-$. This is because the polarisation tensors for gravitons are a product of gluonic polarisations vectors, $\epsilon_{\mu\nu} = \epsilon_\mu \epsilon'_\nu$, and the gluonic polarisation vectors for a leg with momentum k using reference momentum q are

$$\epsilon_\mu^+(k; q) = \frac{\langle q | \gamma^\mu | k \rangle}{\sqrt{2} \langle q k \rangle} \quad \epsilon_\mu^-(k; q) = \frac{[q | \gamma^\mu | k \rangle}{\sqrt{2} [k q]} \quad (4.3)$$

Thus the choice (4.2) engineers a t^{-1} singularity in $\epsilon_\mu^+(k; q)$.

At tree level the soft behaviour is [25]

$$M_n^{\text{tree}} = \left(\frac{1}{t^3} S^{(0)} + \frac{1}{t^2} S^{(1)} + \frac{1}{t} S^{(2)} \right) M_{n-1}^{\text{tree}} + O(t^0) \quad (4.4)$$

Where, for a positive helicity-leg becoming soft [27, 34]

$$S^{(0)} = - \sum_{i=1}^{n-1} \frac{[s i] \langle i \alpha \rangle \langle i \beta \rangle}{\langle s i \rangle \langle s \alpha \rangle \langle s \beta \rangle} \quad (4.5)$$

$$S^{(1)} = - \frac{1}{2} \sum_{i=1}^{n-1} \frac{[s i]}{\langle s i \rangle} \left(\frac{\langle i \alpha \rangle}{\langle s \alpha \rangle} + \frac{\langle i \beta \rangle}{\langle s \beta \rangle} \right) \bar{\lambda}_s^{\dot{a}} \frac{\partial}{\partial \bar{\lambda}_i^{\dot{a}}} \quad (4.6)$$

$$S^{(2)} = \frac{1}{2} \sum_{i=1}^{n-1} \frac{[i s]}{\langle i s \rangle} \bar{\lambda}_s^{\dot{a}} \bar{\lambda}_s^{\dot{b}} \frac{\partial}{\partial \bar{\lambda}_i^{\dot{a}}} \frac{\partial}{\partial \bar{\lambda}_i^{\dot{b}}} \quad (4.7)$$

There are a few points to note regarding these expressions. The expressions are independent of the spinors α and β which arise from the choice of reference spinors for the soft leg as in (4.3). Also, we cannot simply act with these operators on on-shell expressions. For an on-shell expression, the spinors satisfy $\sum_i \lambda_i^a \bar{\lambda}_i^{\dot{a}} = 0$ which implies that not all the variables should be treated as independent. A solution to this is to choose two of the λ 's to eliminate and solve via³

$$\bar{\lambda}_1 = - \sum_{i=3}^n \frac{\langle 2 i \rangle}{\langle 2 1 \rangle} \bar{\lambda}_i \quad \bar{\lambda}_2 = - \sum_{i=3}^n \frac{\langle 1 i \rangle}{\langle 1 2 \rangle} \bar{\lambda}_i \quad (4.8)$$

The universal soft theorem has been proven to hold for tree level [27, 35–38] amplitudes and a bold generalisation is conjectured to remain true for loop amplitudes [25],

$$M_n^{1\text{-loop}} = \left(\frac{1}{t^3} S^{(0)} + \frac{1}{t^2} S^{(1)} + \frac{1}{t} S^{(2)} \right) M_{n-1}^{1\text{-loop}} + O(t^0) \quad (4.9)$$

Unfortunately this does not in fact hold. A possible extension at one-loop would be

$$M_n^{1\text{-loop}} = \left(\frac{1}{t^3} S^{(0)} + \frac{1}{t^2} S^{(1)} + \frac{1}{t} S^{(2)} \right) M_{n-1}^{1\text{-loop}} + \left(\frac{1}{t^2} S_{1\text{-loop}}^{(1)} + \frac{1}{t} S_{1\text{-loop}}^{(2)} \right) M_{n-1}^{\text{tree}} + O(t^0) \quad (4.10)$$

³ With this prescription the momenta k_1 and k_2 are complex valued.

where the operators $S_{1\text{-loop}}^{(i)}$ are loop soft factors acting upon the tree. Since the leading soft term has no loop corrections the loop corrections must start at t^{-2} . In this section we confirm and extend the evidence [27, 28] for loop amplitudes having loop corrections to the sub-sub leading terms beyond (4.10).

The two classes of helicity amplitude which are zero at tree level and are consequently rational at one-loop are the all-plus amplitudes, $M_n^{1\text{-loop}}(1^+, 2^+, \dots, n^+)$, and the single-minus amplitudes, $M_n^{1\text{-loop}}(1^-, 2^+, \dots, n^+)$. For both of these classes the tree amplitudes vanish, so (4.10) will reduce to (4.9). The two classes have different stories [27, 28]: the all-plus amplitudes have the expected behaviour (4.9) whereas the single-minus amplitudes have anomalous sub-sub-leading terms. One difference between the classes is the appearance of double-poles in the single-minus amplitudes. Previously the single-minus amplitudes were only known for $n \leq 6$. Having computed a recursive representation for the single-minus amplitudes we have checked whether they satisfy the soft-theorem (4.10). For $n \leq 8$ we find

- (4.9) is satisfied for a negative helicity leg going soft
- The leading and sub-leading terms in (4.9) are correct for a positive helicity leg going soft
- The sub-sub-leading terms in the amplitudes *do not* match (4.9).

These results are in agreement with those of [27, 28] where the soft theorems were examined in the five and six point cases.

V. CONCLUSIONS

In this article we have used augmented recursive techniques to obtain an explicit form for the non-factorising piece of the single-minus graviton scattering amplitude. To do so we have had to identify the difficult sub-leading poles in the recursion process. These are endemic in non-supersymmetric one-loop computations.

Explicit amplitudes have repeatedly been crucial to test and challenge various hypothesis in interacting field theory. We have generated hard to reach amplitudes which we hope will be useful. For example, we have used our expressions to confirm the presence of loop anomalies to the conjectured universal soft-limit of gravity amplitudes. The explicit forms of the five, six, seven and eight-point amplitudes are available at <http://pyweb.swan.ac.uk/~dunbar/graviton.html>

VI. ACKNOWLEDGEMENTS

This work was supported by STFC grant ST/L000369/1.

Appendix A: Using the KLT relations

The non-factorising contributions to the amplitude depend on MHV currents with two massive legs, e.g.

$$\tau_{\text{grav}}^{\text{MHV}}(a^-, B^-, C^+, d^+, \dots, n^+) \quad (\text{A1})$$

where B and C are non-null. Specifically the amplitude is sensitive to both the leading and sub-leading terms in this current around $B^2 = C^2 = s_{BC} = 0$.

Using a reference spinor, η , a *nullified* version of any massive momentum can be defined

via

$$P^b = P - \frac{P^2}{[\eta|P|\eta]}\eta \rightarrow P(P^b, P^2) = P^b + \frac{P^2}{[\eta|P^b|\eta]}\eta \quad (\text{A2})$$

so that

$$\tau_n = \tau_n(a, B^b, C^b, d, \dots, n, B^2, C^2). \quad (\text{A3})$$

τ_n can be expanded initially around $B^2 = C^2 = 0$,

$$\tau_n = F_1 + B^2 F_2 + C^2 F_3 + \dots \quad \text{where} \quad F_i = F_i(a, B^b, C^b, d, \dots, n). \quad (\text{A4})$$

At tree level the current is not expected to have double poles, but there are single s_{BC} poles. The F_i can therefore be expanded about $s_{BC} = 0$ as

$$F_i = \frac{G_i}{s_{BC}} + \sum_{j=0}^{\infty} H_i^j s_{BC}^j, \quad (\text{A5})$$

where the G_i and H_i^j are evaluated on $B^2 = C^2 = s_{BC} = 0$. The expansion of τ_n is then

$$\tau_n = \frac{G_1 + B^2 G_2 + C^2 G_3}{s_{BC}} + \sum_{j=0}^{\infty} H_1^j s_{BC}^j + \mathcal{O}(B^2) + \mathcal{O}(C^2). \quad (\text{A6})$$

Now consider the quantity τ_n^g which is *good enough* for the current purposes. The expansion of the difference between the two quantities is,

$$\tau_n - \tau_n^g = \frac{\delta G_1 + B^2 \delta G_2 + C^2 \delta G_3}{s_{BC}} + \sum_{j=0}^{\infty} \delta H_1^j s_{BC}^j + \mathcal{O}(B^2) + \mathcal{O}(C^2). \quad (\text{A7})$$

Two constraints must be imposed on τ_n^g :

$$\begin{aligned} \text{C.1 :} & \quad \lim_{s_{BC} \rightarrow 0} \left(s_{BC} \tau_n - s_{BC} \tau_n^g \right) = 0 \\ \text{C.2 :} & \quad \lim_{\substack{B^2 \rightarrow 0, C^2 \rightarrow 0 \\ s_{BC} \neq 0}} \left(\tau_n - \tau_n^g \right) = 0 \end{aligned} \quad (\text{A8})$$

C.1 requires

$$\delta G_1 + B^2 \delta G_2 + C^2 \delta G_3 = 0. \quad (\text{A9})$$

As the G_i are evaluated on $B^2 = C^2 = 0$, this can only hold if $\delta G_1 = \delta G_2 = \delta G_3 = 0$. While C.2 requires

$$\frac{\delta G_1}{s_{BC}} + \sum_{j=0}^{\infty} \delta H_1^j s_{BC}^j = 0. \quad (\text{A10})$$

As the H_1^j are evaluated at $s_{BC} = 0$ this can only hold for any s_{BC} if $\delta H_1^j = 0$. Combining the constraints from C.1 and C.2,

$$\tau_n = \tau_n^g + \mathcal{O}(B^2) + \mathcal{O}(C^2). \quad (\text{A11})$$

Hence conditions C.1 and C.2 ensure that τ_n^g correctly encompasses the leading and sub-leading behaviour of τ_n which determine the non-factorising contributions to the amplitude.

τ_n^g can be obtained by a generalisation of (3.2) to currents:

$$\tau_n^g(B, C, \dots, n, \hat{a}) = i(1)^{n+1} \left[\tau_n^{\text{YM}}(B, C, \dots, n, \hat{a}) \times \sum_{\mathcal{P}_1, \mathcal{P}_2} f(i_1, \dots, i_k; B) \bar{f}(j_1, \dots, j_{k'}) \tau_n^{\text{YM}}(i_1, \dots, i_k, B, n, j_1, \dots, j_{k'}, \hat{a}) \right] + \mathcal{P}_{KLT}(C, d, \dots, n-1), \quad (\text{A12})$$

where the Yang-Mills currents are given by

$$\tau_n^{\text{YM}}(\hat{a}^-, b^+, \dots, f^+, B^-, C^+, g^+, \dots, n^+) = \frac{\langle Ba \rangle^2}{\langle Ca \rangle^2} \frac{1}{\langle ab \rangle \dots \langle ef \rangle} \frac{1}{\langle gh \rangle \dots \langle na \rangle} \left(\frac{\langle aB \rangle \langle aC \rangle \langle ag \rangle \langle fa \rangle}{\langle fB \rangle \langle Cg \rangle \langle gf \rangle} + \frac{\langle aB \rangle \langle af \rangle^2}{\langle Bf \rangle \langle gf \rangle} \frac{[\eta|C|a]}{[\eta|B+C|f]} + \frac{\langle aC \rangle \langle ag \rangle^2}{\langle Cg \rangle \langle gf \rangle} \frac{[\eta|B|a]}{[\eta|B+C|g]} - \frac{\langle a|BC|a \rangle}{s_{BC}} \frac{[\eta|B+C|a]^2}{[\eta|B+C|f][\eta|B+C|g]} \right) + \mathcal{O}(B^2) + \mathcal{O}(C^2) \quad (\text{A13})$$

and the functions f and \bar{f} , partitions, summations etc. are those specified for (3.2).

The fact that τ_n^g as defined by (A12) satisfies conditions C.1 and C.2 follows from the identity

$$\frac{\langle Ca \rangle^2 \langle Ba \rangle^2}{\langle fB \rangle \langle BC \rangle \langle Cg \rangle} + \frac{\langle Ba \rangle \langle Ca \rangle}{\langle BC \rangle} \frac{[\eta|B+C|a]^2}{[\eta|B+C|f][\eta|B+C|g]} = \frac{\langle aB \rangle \langle aC \rangle \langle ag \rangle \langle fa \rangle}{\langle fB \rangle \langle Cg \rangle \langle gf \rangle} + \frac{\langle aB \rangle \langle af \rangle^2}{\langle Bf \rangle \langle gf \rangle} \frac{[\eta|C|a]}{[\eta|B+C|f]} + \frac{\langle aC \rangle \langle ag \rangle^2}{\langle Cg \rangle \langle gf \rangle} \frac{[\eta|B|a]}{[\eta|B+C|g]} \quad (\text{A14})$$

which holds for arbitrary null momenta and when B and C are massive, if λ_B etc. are understood to refer to their nullified forms. (A14) can be rewritten as

$$\frac{\langle aB \rangle \langle aC \rangle \langle ag \rangle \langle fa \rangle}{\langle fB \rangle \langle Cg \rangle \langle gf \rangle} + \frac{\langle aB \rangle \langle af \rangle^2}{\langle Bf \rangle \langle gf \rangle} \frac{[\eta|C|a]}{[\eta|B+C|f]} + \frac{\langle aC \rangle \langle ag \rangle^2}{\langle Cg \rangle \langle gf \rangle} \frac{[\eta|B|a]}{[\eta|B+C|g]} - \frac{\langle a|BC|a \rangle}{s_{BC}} \frac{[\eta|B+C|a]^2}{[\eta|B+C|f][\eta|B+C|g]} = \frac{\langle Ca \rangle^2 \langle Ba \rangle^2}{\langle fB \rangle \langle BC \rangle \langle Cg \rangle} + \mathcal{O}(B^2) + \mathcal{O}(C^2). \quad (\text{A15})$$

The right hand side is precisely the factor required for (A13) to reproduce the Parke-Taylor amplitude in the $B^2, C^2 \rightarrow 0, s_{BC} \neq 0$ limit. Thus in this limit (A12) is the standard KLT relation involving Yang-Mills amplitudes which reproduces the gravity amplitude as required by condition C.2.

Condition C.1 involves the $s_{BC} \rightarrow 0$ limit. τ_n has s_{BC}^{-1} singularities arising from diagrams of the form shown in fig. 8.




FIG. 8: Diagrams contributing to the s_{BC}^{-1} poles in τ_n .

There appear to be double poles in τ_n^g when \mathcal{P}_{KLT} leaves C adjacent to B in the first Yang-Mills factor and $i_k = C$ in the second. However, when $i_k = C$ there is a factor of s_{BC} in f which lowers the order of the pole. In fact these terms are the only ones that are singular in the $s_{BC} \rightarrow 0$ limit. When B and C are adjacent in just one of the Yang-Mills factors there is always a factor of s_{BC} in f which removes the singularity in the current. If B and C are adjacent in the second Yang-Mills factor, $i_k = C$ and there is an explicit factor of s_{BC} in f as previously. If B and C are adjacent in the first Yang-Mills factor, we are considering a term in \mathcal{P}_{KLT} as explicitly shown in (3.2) where $C = i_q$ with $q < k$. As \mathcal{P}_{KLT} hasn't touched C , we can still use the definition of $g(i, j)$ given in (3.5). In this case the product piece of f contains a factor with $m = q$ which takes the form

$$s_{Bi_q} + \sum_{p=q+1}^k g(i_q, i_p) = s_{BC} + \sum_{p=q+1}^k g(C, i_p). \quad (\text{A16})$$

As C is the left most leg in the argument list of τ_n^{grav} and in these cases is untouched by \mathcal{P}_{KLT} , $g(C, i_p) = 0$ and we again have a factor of s_{BC} .

In the $s_{BC} \rightarrow 0$ limit the momentum $B + C$ becomes null and setting $B + C = k$ the singular parts of the Yang-Mills current become

$$\begin{aligned} \tau_n^{\text{YM:sing}}(\hat{a}^-, b^+, \dots, f^+, B^-, C^+, g^+, \dots, n^+) & \\ &= \frac{\langle Ba \rangle^2}{\langle Ca \rangle^2} \frac{1}{\langle ab \rangle \dots \langle ef \rangle \langle gh \rangle \dots \langle na \rangle} \frac{1}{s_{BC}} \left(- \frac{\langle a|BC|a \rangle}{[\eta|B+C|f][\eta|B+C|g]} \frac{[\eta|B+C|a]^2}{[\eta|B+C|f][\eta|B+C|g]} \right) \\ &= \frac{\langle Ba \rangle^2}{\langle Ca \rangle^2} \frac{1}{\langle ab \rangle \dots \langle ef \rangle \langle fk \rangle \langle kg \rangle \langle gh \rangle \dots \langle na \rangle} \left(- \frac{\langle a|BC|a \rangle}{s_{BC}} \langle ka \rangle^2 \right) \\ &= \frac{\langle Ba \rangle^2}{\langle Ca \rangle^2} A_{n-1}^{\text{YM}}(\hat{a}^-, b^+, \dots, f^+, k^-, g^+, \dots, n^+) \left(- \frac{\langle a|BC|a \rangle}{s_{BC} \langle ka \rangle^2} \right) \end{aligned} \quad (\text{A17})$$

The singular part of the current thus becomes

$$\begin{aligned} \tau_n^{g:\text{sing}} &= i(1)^{n+1} \left[\tau_n^{\text{YM:sing}}(B, C, \dots, n, \hat{a}) \times \right. \\ &\quad \left. \sum_{\mathcal{P}'_1, \mathcal{P}'_2} f(i_1, \dots, i_k; B) \bar{f}(j_1, \dots, j_{k'}) \tau_n^{\text{YM:sing}}(i_1, \dots, i_k, B, n, j_1, \dots, j_{k'}, \hat{a}) \right] \\ &\quad + \mathcal{P}'_{KLT}(d, \dots, n-1), \end{aligned} \quad (\text{A18})$$

Where \mathcal{P}'_{KLT} and \mathcal{P}'_1 denote the subset of the original permutation sums that leave C adjacent to B in both Yang-Mills factors, i.e. C is excluded from \mathcal{P}'_{KLT} and $i_k = C$. The momentum factor f is then

$$f(i_1, \dots, i_{k-1}, C; B) = s_{BC} \prod_{m=1}^{k-2} \left(s_{Bi_m} + \sum_{p=m+1}^k g(i_m, i_p) \right) \left(s_{Bi_{k-1}} + g(i_{k-1}, C) \right) \quad (\text{A19})$$

As we are only interested in those permutations that leave leg C unmoved, C remains to the left of all possible legs i_{k-1} and $g(i_{k-1}, C) = 0$. Thus

$$f(i_1, \dots, i_{k-1}, C; B) = s_{BC} f(i_1, \dots, i_{k-1}; B) \quad (\text{A20})$$

The summations and momentum factors are thus precisely those of the $(n-1)$ -point KLT expansion.

$$\tau_n^{g:sing} = - \frac{\langle Ba \rangle^4 \langle a|BC|a \rangle^2}{\langle Ca \rangle^4 \langle ka \rangle^4} \frac{1}{s_{BC}} M_{n-1}(\hat{a}^-, b^+, \dots, f^+, k^-, g^+, \dots, n^+) \quad (\text{A21})$$

Taking the three-point gravity vertex to be the square of its Yang-Mills counter part, in axial gauge with $\lambda_\eta = \lambda_a$, we have

$$V(B^-, C^+, k^+) = \frac{[Ck]^2 \langle Ba \rangle^4}{\langle Ca \rangle^2 \langle ka \rangle^2} = \frac{\langle Ba \rangle^4 \langle a|Ck|a \rangle^2}{\langle Ca \rangle^4 \langle ka \rangle^4} = \frac{\langle Ba \rangle^4 \langle a|CB|a \rangle^2}{\langle Ca \rangle^4 \langle ka \rangle^4} \quad (\text{A22})$$

In the B, C co-linear limit $\tau_n^{g:sing}$ precisely reproduces the factorisation of the gravity current and hence satisfies condition C.1.

Appendix B: Using Soft Theorems to Construct Amplitudes

If the soft-limit theorems hold they could be a substitute for the rather cumbersome augmented recursion techniques. If we take

$$\lambda_n = t\lambda_s \quad \bar{\lambda}_n = \bar{\lambda}_s \quad (\text{B1})$$

together with a specific choice for implementing momentum conservation,

$$\bar{\lambda}_1(t) = \bar{\lambda}_1 - t \frac{\langle 2s \rangle}{\langle 21 \rangle} \bar{\lambda}_s \quad \bar{\lambda}_2(t) = \bar{\lambda}_2 - t \frac{\langle 1s \rangle}{\langle 12 \rangle} \bar{\lambda}_s \quad (\text{B2})$$

and treat t as a complex parameter, then we have the complexified amplitude $M_n(t)$. We would like to construct the n -point amplitude $M_n(t = t_0)$ recursively from M_{n-1} . This is possible by evaluating

$$\int dt \frac{M_n(t)}{t - t_0}. \quad (\text{B3})$$

Providing $M_n(t) \rightarrow 0$ as $t \rightarrow \infty$ we have

$$M_n(t_0) = - \sum_i \text{Residue} \left[\frac{M_n(t)}{t - t_0}, t_i \right] \quad (\text{B4})$$

At values of t (say t_i) where $M_n(t)$ has simple poles

$$\text{Residue}\left[\frac{M_n(t)}{t-t_0}, t_i\right] = \frac{1}{t_i-t_0} \text{Residue}[M_n(t), t_i] \quad (\text{B5})$$

which can be determined from the factorisation theorems as usual. When $M_n(t)$ has a non-simple pole, the residue of $M_n(t)/(t-t_0)$ in general involves all the singular terms. For example if $M_n(t)$ has a multiple singularity at $t=0$ and the expansion about $t=0$ is

$$M_n(t) = \frac{a_{-N}}{t^N} + \frac{a_{-N+1}}{t^{N-1}} \cdots \frac{a_{-1}}{t} + \text{non-singular} \quad (\text{B6})$$

then

$$\text{Residue}\left[\frac{M_n(t)}{(t-t_0)}, 0\right] = -\frac{a_{-N}}{t_0^N} - \frac{a_{-N}}{t_0^N} + \cdots - \frac{a_{-1}}{t_0^1} \quad (\text{B7})$$

Provided explicit formulae exist for the sub-leading singularities we may in principle use Cauchy to compute the result. The soft theorems thus might be useful in constructing these amplitudes.

Taking leg n to be of positive helicity and scaled according to (B1), for large t we find

$$A : M_n(1^-, 2^+, 3^+, \dots, n^+)(t) \longrightarrow t^{n-4} \quad (\text{B8})$$

While taking leg 1 to have negative helicity and scaled according to the complex conjugate of (B1), at large t we find

$$B : M_n(1^-, 2^+, 3^+, \dots, n^+)(t) \longrightarrow t^{-3} \quad (\text{B9})$$

There is an unfortunate conspiracy here. In case A, the shifted amplitude has single poles except at $t=0$ AND at $t=0$ the soft-limit is well known. However the large- t behaviour generates a contribution at infinity which can not be directly obtained from a factorisation of the amplitude. In case B the large t limit is good however the amplitude does not satisfy the soft limit theorems AND the amplitude has double poles at some points t_i which would require extra information to determine.

Appendix C: Mathematica form of Amplitudes

At <http://pyweb.swan.ac.uk/~dunbar/graviton.html> explicit forms of the single-minus amplitudes are available in `Mathematica` format. These are presented as polynomials in λ_i and $\bar{\lambda}_i$, specifically as `lambda[x, i]` and `barlambda[x, i]`. The expressions satisfy:

- symmetry under interchange of positive legs.
- independence of the choice of positive shift leg. This is actually quite a strenuous test which ties together all the different terms in the amplitude. If we view amplitudes as the area of some polyhedron [39] then the BCFW shift corresponds to a triangulation of the area.
- The amplitudes all satisfy the leading soft behaviour: this checks (or fixes) the overall normalisations.
- The amplitudes have the correct collinear limits. The collinear limit occurs when legs k_a and k_b are collinear, $k_a \cdot k_b \longrightarrow 0$. Unlike Yang-Mills amplitudes, gravity amplitudes are not singular in the collinear limit, but acquire a ‘‘phase-singularity’’ [33, 40]. If $k_a \longrightarrow zK$

and $k_b \rightarrow (1-z)K$, the singularity is

$$M_n(\cdots, a^{h_a}, b^{h_b}) \xrightarrow{a||b} \sum_{h'} \text{Sp}_{-h'}^{h_a h_b} M_{n-1}(\cdots, K^{h'}) \quad (\text{C1})$$

where the h 's denote the various helicities of the gravitons. The non-zero ‘‘splitting functions’’ are

$$\text{Sp}_-^{++} = -\frac{[ab]}{z(1-z)\langle ab \rangle}, \quad \text{Sp}_+^{-+} = -\frac{z^3 [ab]}{(1-z)\langle ab \rangle}. \quad (\text{C2})$$

An important result of ref. [33] is that the splitting functions do not obtain loop corrections.

We have made the amplitudes available in a very explicit, if rather cumbersome, polynomial form. The size of these polynomials grows rapidly with the number of points. In Table I we present the `LeafCount` of these `Mathematica` expressions as a rough indicator of the growth. The MHV tree and all-plus one-loop amplitudes are included for comparison. None of these has been optimised and they are all equally inefficient.

Amplitude	$n = 4$	$n = 5$	$n = 6$	$n = 7$	$n = 8$
$M_n^{\text{tree}}(-, -, +, \cdots, +)$	144	512	2,318	13,244	95,000
$M_n^{1\text{-loop}}(+, +, +, \cdots, +)$	216	4,085	84,470	553,145	3,814,189
$M_n^{1\text{-loop}}(-, +, +, \cdots, +)$	252	4,959	150,474	4,462,636	220,910,118
Δ_n	—	—	42,750	943,051	47,265,553

TABLE I: `LeafCount` of some rational amplitudes expressed as polynomials in λ and $\bar{\lambda}$.

-
- [1] Amplitudes 2015 at <http://amp15.itp.phys.ethz.ch/> provides an overview of the subject at the time of writing of this article
 - [2] R.J. Eden, P.V. Landshoff, D.I. Olive, J.C. Polkinghorne, *The Analytic S Matrix*, (Cambridge University Press, 1966).
 - [3] E. Witten, Commun. Math. Phys. **252** (2004) 189 [hep-th/0312171].
 - [4] R. Britto, F. Cachazo, B. Feng and E. Witten, Phys. Rev. Lett. **94** (2005) 181602 [hep-th/0501052].
 - [5] Z. Bern and G. Chalmers, Nucl. Phys. B **447** (1995) 465 [hep-ph/9503236].
 - [6] Z. Bern, L. J. Dixon, D. C. Dunbar, D. A. Kosower, Nucl. Phys. **B435** (1995) 59 [hep-ph/9409265].
 - [7] Z. Bern, L. J. Dixon, D. C. Dunbar and D. A. Kosower, Nucl. Phys. B **425** (1994) 217 [hep-ph/9403226].
 - [8] N. E. J. Bjerrum-Bohr, D. C. Dunbar, H. Ita, W. B. Perkins and K. Risager, JHEP **0612** (2006) 072 [arXiv:hep-th/0610043].
 - [9] R. E. Cutkosky, J. Math. Phys. **1** (1960) 429.
 - [10] Z. Bern, L. J. Dixon and D. A. Kosower, Nucl. Phys. B **513** (1998) 3 [hep-ph/9708239].
 - [11] R. Britto, F. Cachazo and B. Feng, Nucl. Phys. B **725** (2005) 275 [hep-th/0412103].

- [12] D. C. Dunbar, W. B. Perkins and E. Warrick, JHEP **0906** (2009) 056 [arXiv:0903.1751 [hep-ph]].
- [13] S. J. Bidder, N. E. J. Bjerrum-Bohr, D. C. Dunbar and W. B. Perkins, Phys. Lett. B **612** (2005) 75 [hep-th/0502028].
- [14] D. Forde, Phys. Rev. D **75** (2007) 125019 [arXiv:0704.1835 [hep-ph]].
- [15] N. E. J. Bjerrum-Bohr, D. C. Dunbar and W. B. Perkins, JHEP **0804** (2008) 038 [arXiv:0709.2086 [hep-ph]].
- [16] P. Mastrolia, Phys. Lett. B **644** (2007) 272 [arXiv:hep-th/0611091].
- [17] Z. Bern, L. J. Dixon and D. A. Kosower, Phys. Rev. D **71** (2005) 105013 [hep-th/0501240].
- [18] A. Brandhuber, S. McNamara, B. Spence and G. Travaglini, JHEP **0703** (2007) 029 [hep-th/0701187].
- [19] G. Mahlon, Phys. Rev. D **49** (1994) 4438 [hep-ph/9312276].
- [20] D. Vaman and Y. P. Yao, arXiv:0805.2645 [hep-th].
- [21] Z. Bern, D. C. Dunbar and T. Shimada, Phys. Lett. B **312** (1993) 277 [hep-th/9307001];
D. C. Dunbar and P. S. Norridge, Nucl. Phys. B **433** (1995) 181 [hep-th/9408014].
- [22] D. C. Dunbar, J. H. Eittle and W. B. Perkins, JHEP **1006** (2010) 027 [arXiv:1003.3398 [hep-th]].
- [23] Steven Weinberg, Phys.Rev. **140** (1965) B516-B524.
- [24] C. D. White, JHEP **1105** (2011) 060 [arXiv:1103.2981 [hep-th]].
- [25] F. Cachazo and A. Strominger, arXiv:1404.4091 [hep-th].
- [26] H. Bondi, M. G. J. van der Burg and A. W. K. Metzner, Proc. Roy. Soc. Lond. A **269** (1962) 21;
R. K. Sachs, Proc. Roy. Soc. Lond. A **270** (1962) 103.
- [27] Z. Bern, S. Davies and J. Nohle, Phys. Rev. D **90** (2014) 8, 085015 [arXiv:1405.1015 [hep-th]].
- [28] S. He, Y. t. Huang and C. Wen, JHEP **1412** (2014) 115 [arXiv:1405.1410 [hep-th]].
- [29] S. D. Alston, D. C. Dunbar and W. B. Perkins, Phys. Rev. D **86** (2012) 085022 [arXiv:1208.0190 [hep-th]].
- [30] C. Schwinn and S. Weinzierl, JHEP **0505**, 006 (2005) [hep-th/0503015].
- [31] D. A. Kosower, Nucl. Phys. B **335** (1990) 23.
- [32] H. Kawai, D. C. Lewellen and S. H. H. Tye, Nucl. Phys. B **269** (1986) 1.
- [33] Z. Bern, L. J. Dixon, M. Perelstein and J. S. Rozowsky, Nucl. Phys. B **546** (1999) 423 [hep-th/9811140].
- [34] J. Broedel, M. de Leeuw, J. Plefka and M. Rosso, Phys. Rev. D **90** (2014) 6, 065024 [arXiv:1406.6574 [hep-th]].
- [35] B. U. W. Schwab and A. Volovich, Phys. Rev. Lett. **113** (2014) 10, 101601 [arXiv:1404.7749 [hep-th]].
- [36] T. Klose, T. McLoughlin, D. Nandan, J. Plefka and G. Travaglini, JHEP **1507** (2015) 135 [arXiv:1504.05558 [hep-th]].
- [37] M. Zlotnikov, JHEP **1410** (2014) 148 [arXiv:1407.5936 [hep-th]].
- [38] C. Kalousios and F. Rojas, JHEP **1501** (2015) 107 [arXiv:1407.5982 [hep-th]].
- [39] N. Arkani-Hamed and J. Trnka, JHEP **1410** (2014) 30 [arXiv:1312.2007 [hep-th]].
- [40] D. C. Dunbar, J. H. Eittle and W. B. Perkins, Phys. Rev. D **86** (2012) 026009 [arXiv:1203.0198 [hep-th]].