



Swansea University
Prifysgol Abertawe



Cronfa - Swansea University Open Access Repository

This is an author produced version of a paper published in :
CORTEX

Cronfa URL for this paper:

<http://cronfa.swan.ac.uk/Record/cronfa26786>

Paper:

Kremláek , J., Kreegipuu , K., Tales, A., Astikainen, P., Pöldver, N., Näätänen, R. & Stefanics, G. Visual mismatch negativity (vMMN): A review and meta-analysis of studies in psychiatric and neurological disorders.. *CORTEX*

<http://dx.doi.org/10.1016/j.cortex.2016.03.017>

This article is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Authors are personally responsible for adhering to publisher restrictions or conditions. When uploading content they are required to comply with their publisher agreement and the SHERPA RoMEO database to judge whether or not it is copyright safe to add this version of the paper to this repository.

<http://www.swansea.ac.uk/iss/researchsupport/cronfa-support/>

Accepted Manuscript

Visual Mismatch Negativity (vMMN): A review and meta-analysis of studies in psychiatric and neurological disorders

Jan Kremláček, Kairi Kreegipuu, Andrea Tales, Piia Astikainen, Nele Pöldver, Risto Näätänen, Dr. Gábor Stefanics



PII: S0010-9452(16)30055-7

DOI: [10.1016/j.cortex.2016.03.017](https://doi.org/10.1016/j.cortex.2016.03.017)

Reference: CORTEX 1712

To appear in: *Cortex*

Received Date: 30 June 2015

Revised Date: 1 March 2016

Accepted Date: 17 March 2016

Please cite this article as: Kremláček J, Kreegipuu K, Tales A, Astikainen P, Pöldver N, Näätänen R, Stefanics G, Visual Mismatch Negativity (vMMN): A review and meta-analysis of studies in psychiatric and neurological disorders, *CORTEX* (2016), doi: 10.1016/j.cortex.2016.03.017.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Jan Kremláček^{a,b}, Kairi Kreegipuu^c, Andrea Tales^d, Piia Astikainen^e, Nele Pöldver^{c,f}, Risto Näätänen^{c,g}, Gábor Stefanics^{h,i*}

**Corresponding author:*

Dr. Gábor Stefanics,
Translational Neuromodeling Unit (TNU),
Institute for Biomedical Engineering, University of Zurich & ETH Zurich,
Wilfriedstrasse 6
CH-8032 Zürich, Switzerland.

Email: stefanics@biomed.ee.ethz.ch

Phone: +41 (0)44 634 9121

Fax: +41 (0)44 634 9125

^a Department of Pathological Physiology, Faculty of Medicine in Hradec Kralove, Charles University in Prague, Hradec Kralove, Czech Republic. Simkova 870, 500 38 Hradec Králové. E-mail: jan.kremlacek@lfhk.cuni.cz

^b Department of Neurology, Faculty of Medicine, Charles University in Prague, University Hospital Hradec Kralove, Hradec Kralove, Czech Republic

^c Institute of Psychology, University of Tartu, Tartu, Estonia. Näituse 2, Tartu 50409. E-mail: kairi.kreegipuu@ut.ee

^d Department of Psychology, College of Human and Health Sciences, Swansea University, Swansea, Wales, UK. Singleton Park, Swansea SA2 8PP. E-mail: a.tales@swansea.ac.uk

^e Department of Psychology, University of Jyväskylä, Jyväskylä, Finland. Ylistönmäentie 33, building Y33, PO Box 35, 40014 Jyväskylä. E-mail: piia.astikainen@jyu.fi

^f Doctoral School of Behavioural, Social and Health Sciences, University of Tartu, Tartu, Estonia. Näituse 2, Tartu 50409. E-mail: nele.poldver@ut.ee

^g Center of Functionally Integrative Neuroscience (CFIN), University of Aarhus, Aarhus, Denmark
Cognitive Brain Research Unit (CBRU), Institute of Behavioural Sciences, University of Helsinki, Helsinki, Finland.
Näituse 2, Tartu 50409, Estonia. E-mail: risto.naatanen@helsinki.fi

^h Translational Neuromodeling Unit (TNU), Institute for Biomedical Engineering, University of Zurich & ETH Zurich, Zurich, Switzerland. Wilfriedstrasse 6, CH-8032 Zürich. E-mail: stefanics@biomed.ee.ethz.ch

ⁱ Laboratory for Social and Neural Systems Research, Department of Economics, University of Zurich, Zurich, Switzerland

experiments use visual stimulus repetition to induce predictions, and vMMN is obtained by subtracting the response to rare unpredicted stimuli from those to frequent stimuli. One increasingly popular interpretation of the mismatch response postulates that vMMN, similar to its auditory counterpart (aMMN), represents a prediction error response generated by cortical mechanisms forming probabilistic representations of sensory signals. Here we discuss the physiological and theoretical basis of vMMN and review thirty-three studies from the emerging field of its clinical applications, presenting a meta-analysis of findings in schizophrenia, mood disorders, substance abuse, neurodegenerative disorders, developmental disorders, deafness, panic disorder and hypertension. Furthermore, we include reports on aging and maturation as they bear upon many clinically relevant conditions. Surveying the literature we found that vMMN is altered in several clinical populations which is in line with aMMN findings. An important potential advantage of vMMN however is that it allows the investigation of deficits in predictive processing in cognitive domains which rely primarily on visual information; a principal sensory modality and thus of vital importance in environmental information processing and response, and a modality which arguably may be more sensitive to some pathological changes. However, due to the relative infancy of research in vMMN compared to aMMN in clinical populations its potential for clinical application is not yet fully appreciated. The aim of this review and meta-analysis therefore is to present, in a detailed systematic manner, the findings from clinically-based vMMN studies, to discuss their potential impact and application, to raise awareness of this measure and to improve our understanding of disease upon fundamental aspects of visual information processing.

Highlights

1. VMMN is an ERP component, a correlate of automatic predictive processes.
2. VMMN relies on stimulus repetition similarly to fields that use repetition suppression.
3. Altered vMMN has been reported in several clinical populations.
4. A systematic review and meta-analysis of 33 studies is presented.
5. VMMN might be a useful tool for studying impairments of visual predictive processes.

Keywords

Visual mismatch negativity (vMMN), repetition suppression (RS), stimulus specific adaptation (SSA), schizophrenia, effect size

are repeated to induce an automatic prediction pertaining to the probability of an event. This prediction is often thought about as 'regularity' extracted from the stimulus stream (Winkler, 2007) and its presence is usually demonstrated indirectly by showing that stimuli that deviate from the frequent stimuli evoke a mismatch response (e.g., Stefanics et al., 2011, 2012, 2014). MMN studies thus investigate the effect of stimulus repetition by focusing on neural responses specific to rare (deviant) events embedded in a stream of frequent (standard) stimuli. The emphasis in many MMN studies therefore falls on automatic change detection, which is crucial for survival as unexpected environmental changes may carry important information that might trigger an orientation reaction (Sokolov, 1963). Nevertheless, this is only possible if probabilities of environmental events are continuously monitored and updated.

The MMN component, originally described in the auditory domain (aMMN) (Näätänen, Gaillard, & Mantysalo, 1978), has its analogues in the other sensory modalities as well and has been systematically explored in the visual domain for more than twenty years (for the first review, see Pazo-Alvarez et al., 2003, for more recent reviews, see Kimura et al., 2011; Stefanics et al., 2014). At the time of writing this article in January 2016, Pub Med (www.pubmed.com) returned **100** entries for the term "*visual mismatch negativity*". Similar to auditory MMN, that can be elicited by unexpected changes in different stimulus features, such as pitch, duration, intensity, etc. (for a review, see Näätänen et al., 2010), there is evidence that visual MMN can be elicited by rare changes in features such as line orientation (Astikainen et al., 2004, 2008; Czigler & Pató, 2009; Czigler & Sulykos, 2010; Flynn et al., 2009; Kimura et al., 2009, 2010), spatial frequency (Heslenfeld, 2003; Kenemans et al., 2003, 2010; Maekawa et al., 2005; Sulykos & Czigler, 2011), color (Czigler et al., 2002, 2004; Grimm et al., 2009; Horimoto et al., 2002; Kimura et al., 2006; Liu and Shi, 2008; Mazza et al., 2005; Mo et al., 2011; Müller et al., 2010; Stefanics et al., 2011; Thierry et al., 2009), luminance (Stagg et al., 2004), illusory brightness (Sulykos & Czigler, 2014), or motion (Amenedo et al., 2007; Kremláček et al. 2006; Kuldkepp et al., 2013; Lorenzo-López et al.; 2004). Furthermore, again similar to aMMN that can be elicited by rare changes in more abstract attributes of the acoustic stimuli than simple physical features, vMMN is also sensitive to higher-level attributes such as sequential regularities (Stefanics et al., 2011), object structure (Müller et al., 2013), symmetry (Kecskés-Kovács et al., 2013a), laterality of body parts (Stefanics & Czigler, 2012), or attributes of socially more relevant stimuli such as facial emotions (Astikainen & Hietanen, 2009; Fujimura & Okanoya, 2013; Susac et al., 2004, 2010; Zhao and Li, 2006; Stefanics et al., 2012), as well as facial gender (Kecskés-Kovács et al., 2013b). Thus, there is ample evidence that automatic perceptual predictive mechanisms operate in the visual modality too, which can be probed by experimentally manipulating statistical properties of a wide variety of stimulus attributes or their relationships.

Early accounts of the aMMN suggested that MMN generation relied on a strong auditory sensory memory trace encoding the repeating stimulus (Winkler, 2007). More recent theories suggest that the perceptual system extracts environmental regularities and represents expected events; i.e. the formation of predictions has been suggested as the primary function of the neural processes underlying the MMN (Winkler and Czigler, 1998). The basis of both auditory and visual MMN studies is the ubiquitous phenomenon that brain response properties to frequent events differ

might also depend on the experience of the subjects with the given stimulus category (Grotheer & Kovács, 2014; Sulykos et al., 2015), i.e., perceptual expertise. Often referred to as ‘functional magnetic resonance imaging adaptation’ (fMRIa), the method is particularly popular in neuroimaging to explore functional properties of neuronal populations (Krekelberg et al., 2006). While studies on RS or SSA investigate response attenuation over repetitions, MMN studies focus on the relative difference between suppressed and unsuppressed brain responses to predicted and unpredicted stimuli, respectively. Adaptation research and the MMN field thus focus on effects of manipulation of stimulus probabilities from different angles, although they use similar experimental techniques to study closely related phenomena.

Stimulus repetition often results in faster reaction times or improvement in stimulus detection and RS of the hemodynamic response has been associated with behavioral phenomenon of priming (Schacter & Buckner, 1998; Wiggs & Martin, 1998; Buckner et al., 2000; Henson & Rugg, 2003; Wig et al., 2005). Adaptation studies in psychophysics tend to use continuous stimulus exposure instead of stimulus repetition to study perceptual and behavioral effects. From the perspective of predictive coding longer exposure and stimulus repetition both provide the visual system with more sensory evidence about the probable external cause of the perceived stimulus. Psychophysics defines visual adaptation in terms of brief or long exposures and the ensuing aftereffects (Webster, 2011). Thus, in psychophysics the term adaptation is used to describe perceptual changes that follow exposure to recently viewed stimuli. Adaptation is primarily characterized by a loss in sensitivity to the adapting stimulus which is accompanied by a boost in neural responses to unexpected events. At the behavioral level adaptation can facilitate discrimination (e.g., McDermott et al., 2010; Kristjansson, 2011) similar to priming paradigms, and unexpected stimuli that are associated with MMN are also detected more easily (Garrido et al., 2013; Tiitinen et al., 1994; Solomon & Kohn, 2014).

In electrophysiology simpler forms of use-dependent adaptation of firing rates are distinguished from more sophisticated, context-dependent SSA (Kohn, 2007; Nelken, 2014; Pérez-González & Malmierca, 2014). Using the term ‘adaptation’ to describe stimulus-specific attenuation of neural activity in electrophysiology is somewhat unfortunate (Nelken & Ulanovsky, 2007). According to a widely accepted definition by Dudai (2002), adaptation is a “Use-dependent response decrement that occurs because of sensory and peripheral processes”. SSA is not use-dependent, that is, the reduction in the response to a stimulus does not generalize, or only partially generalizes, to other, rare stimuli (Movshon & Lennie, 1979; Nelken, 2014). A better term for SSA would be “habituation”, defined by Dudai (2002) as the “gradual diminution of the response to a stimulus following the repeated presentation of the same, or a similar, stimulus”. Habituation shows stimulus-specificity, rate-sensitivity, and other characteristics of SSA. The nomenclature is problematic (Nelken & Ulanovsky, 2007), and without looking at the details of the phenomena in question, one might easily get confused and/or swamped in semantics. Since different fields use the terms RS, SSA, habituation, adaptation or refractoriness to describe several related but distinct concepts and phenomena, it is difficult to pin one concept on one term (O’Shea 2015, Stefanics et al. 2016).

afterhyperpolarization, 2) synaptic depression due to the depletion of vesicles from the presynaptic terminal, and 3) synaptic (network) mechanisms (Kohn, 2007). Since the 'refractory' state of a neuron after spiking is too short to be responsible for the ERP amplitude decrease after repeated stimulation and synaptic depletion also occurs presumably only at higher stimulation rates than in MMN experiments, RS in MMN experiments likely results from network mechanisms. The proposed mechanisms involve increased inhibition or decreased excitation (Ibbotson, 2005). Adaptation effects are substantially more complex than suggested by traditional fatigue-based descriptions which focused on mechanisms in individual cells. Our emerging understanding is that adaptation modifies neural population coordination and its effects cascade through the processing stages, affecting networks further down the processing stream (Solomon & Kohn, 2014).

Several attempts have been made to identify the single-cell correlates of the scalp-recorded aMMN. SSA is the closest known single-neuron phenomenon of aMMN (for reviews see Nelken and Ulanovsky, 2007; Escera and Malmierca, 2014; Pérez-González and Malmierca, 2014; Nelken, 2014). The magnitudes of SSA and aMMN are both negatively correlated with the probability of the deviant but positively correlated with the difference between standard and deviant. However, an important difference is the earlier timing of SSA relative to aMMN, therefore Nelken and Ulanovsky (2007) suggested that SSA is a correlate of change detection in the primary auditory cortex upstream of MMN. This is in line with a recent EEG study where SSA was found to depend critically on statistical context in the auditory cortex (Herrmann et al., 2015), and that aMMN itself is a compound response of primary and higher-level cortical areas with longer response latencies. Adaptation has been observed in electrophysiological studies throughout the visual system including the retina (Hosoya et al., 2005), thalamus (Solomon et al., 2004), superior colliculus (Boehnke et al., 2011), and several cortical areas (Müller et al., 1999; Motter, 2006; Kremláček et al. 2007, Meyer et al., 2014; Kaliukhovich & Vogels, 2014; Ramachandran et al., 2016; for a review, see Vogels, 2015). Although the exact relationship of adaptation at different levels of the visual processing hierarchy to change detection as observed in vMMN studies (e.g., Kimura et al., 2011; Stefanics et al., 2014) is not clear, there is general agreement that the function of adaptation is to match response properties of the sensory system to the current environment (Clifford et al., 2007; Webster, 2011) and thus improve stimulus discrimination or detection of improbable stimuli (Kohn, 2007; Solomon & Kohn, 2014; Carandini & Heeger, 2012; Benucci, Saleem & Carandini, 2013; Wark et al., 2007).

Three models of network mechanisms have been put forward (Grill-Spector et al., 2006) to explain RS in terms of population dynamics. The model of *neural fatigue* is similar to the idea of "refractoriness" in the MMN field, and explains RS by firing rate attenuation, i.e., where the initial high neural response rate of spiking to a constant stimulus is not maintained but instead declines over time; a common feature of many sensory neurons (Hille, 1992; Pérez-González & Malmierca, 2014). The second model proposes that RS involves *sharpening* of the neural populations that generate the initial response, such that fewer neurons respond to repeated stimuli (Wiggs & Martin, 1998; Henson & Rugg, 2003). Specifically, neurons responding to features that are not essential for recognizing the object attenuate their responses, thus the network becomes sparser and more selective (Wiggs & Martin, 1998), thus according to this view, RS is a by-product of sharpening

in the decreased cumulative hemodynamic response (James et al., 2006; Grill-Spector et al., 2006).

The facilitation model shows a strong similarity to a more recent adaptation-based model of the MMN, which proposed that the MMN is generated by fresh-afferent activity of cortical neurons, the latency and amplitude of which is modulated during stimulus repetition (May & Tiitinen, 2010). However, other recent studies using neurobiologically informed computational models (Garagnani & Pulvermüller, 2011; Wacongne et al., 2012) found that the MMN is likely to be generated by active cortical predictive mechanisms rather than passive adaptation. The dynamics of the network that is thought to generate the aMMN has been extensively investigated with large-scale models which incorporate hypotheses of both adaptation and change detection (Garrido et al., 2007, 2008, 2009; Kiebel et al., 2007). Results of these dynamic causal modeling (DCM) studies show that RS is associated with repetition-dependent plasticity in connections within and between generating structures suggesting a conjoint role of adaptation (May & Tiitinen, 2010) and active model-adjustment (Winkler et al., 1996) processes in the aMMN response. The potential of predictive coding theory to provide a comprehensive explanation of MMN phenomenology is demonstrated by a recent modeling study which suggest that the aMMN reflects approximate Bayesian learning of sensory regularities (Lieder et al., 2013a), where prediction errors (i.e., aMMN responses) are used to adjust a probabilistic model of the environment (Lieder et al., 2013b). Assuming that vMMN has a functionally similar role in the visual system to that of aMMN in the auditory system, the implication of these modeling studies is that visual perception is an active process relying on an internal probabilistic representation of the world (cf. Wolfe, 1999) which serves to predict the incoming sensory signal and vMMN is a neural correlate of the update process when there is a mismatch between the predicted and incoming signal.

The predictive coding theory elegantly accommodates adaptation effects observed in psychophysics, behavioral priming, RS, as well as ERP components including the MMN (Friston, 2005). Predictive coding posits that the brain actively generates probabilistic models of the causes of the sensory input (for a comprehensive review, see Clark, 2013), and provides a framework which explains both response attenuation to repeated events and a larger response to unpredicted events. Predictive coding emphasizes the active nature of perceptual inference: instead of being a passive analyzer of bottom-up sensory information (Egner et al., 2010), the brain is thought to actively predict the sensory signal by generating a probabilistic model of the causes of the sensory signals. Thus, according the theory, perception is actively using a set of probabilistic assumptions to infer the most likely cause of the sensory signals by matching incoming sensory information to prediction of that signal (Rao & Ballard, 1999; Lee & Mumford, 2003; Friston, 2005). Repetition effects are thought to reflect ongoing statistical learning during which the generative model is updated. In MMN studies RS often manifests as decrease of the negative component elicited by rare events, i.e., as a relative positivity to repeated stimuli, which was suggested to represent rapid SSA underlying sensory memory formation (Haenschel et al., 2005). Predictive coding suggests that RS is brought about by fast changes in synaptic connections (Baldeweg, 2006, 2007; Garrido et al., 2009) within and between hierarchical levels of neural populations. Top-down predictions based on prior events are thought to explain away prediction errors at lower hierarchical levels, thus RS effects have been proposed to

violation of the model's prediction by an unexpected event, i.e., a deviant stimulus. These generative models can be viewed as hierarchical representations of observed events of the environment, similar to predictive perceptual object representations suggested by Winkler & Czigler (2012).

While RS is a tool used relatively seldom in clinical research, the aMMN has a long track record in studying clinical populations. Besides its important contribution to the understanding of automatic sensory processing, the aMMN is a promising technique for clinical research and possibly also for clinical applications, as it might help to better understand disease mechanisms and dissect spectrum diseases into well-defined subgroups to guide diagnosis, predict disease trajectory and response to treatments (Luck et al., 2011; Light & Näätänen, 2013). Recording the aMMN is a non-invasive and cost-effective method, and in the case of patients with limited cooperation, it allows assessment of brain function without using a behavioral task. The applications of aMMN as a clinical research tool have been extensively reviewed (Näätänen & Kähkönen, 2009; Näätänen et al., 2011, 2012a, 2014; Näätänen, Paavilainen, Rinne, & Alho, 2007a). However, the literature is still lacking a comprehensive review of vMMN studies with clinical relevance. Among articles concerning vMMN, we found **33** studies describing its use in a clinical context, and in the following we review these articles and compare their findings to those obtained in aMMN studies. In particular, we summarize the findings in schizophrenia, mood disorders, substance abuse, neurodegenerative disorders, developmental disorders, deafness, panic disorder and hypertension and present a meta-analysis of the reviewed studies in the Discussion section. Furthermore, we include reports on aging and maturation as these processes impact on several clinically relevant conditions.

2 Visual MMN studies in clinical populations

In general, visual MMN has been used to study similar neuropsychiatric and neurological disorders as the auditory MMN. We list studies in Table 1 for each disorder that we overview in the following sections.

Table 1 – vMMN clinical studies according to clinical topics

Source/ Diagnosis/ number in figures	Group	vMMN stimuli	Results	Intergroup vMMN differences	
				Temporal interval (Effect size, Power)	Area/Electrode
Schizophrenia and schizoaffective disorder					
(Csukly, Stefanics, Komlósi, Czigler, & Czobor, 2013) Schizophrenia 1	patients 24 evaluated, 4 excluded (11F, 34.2 ± 10.3 y), ASEm CG 24	emotional faces (happy, fearful)	NO significant vMMN response for patients	170 – 220 ms ^{PD} (<i>eCd</i> =0.87, <i>Pw</i> =0.84)	<u>temporal left ROI (happy)</u> ; central ROI (happy) ^{PH} ; (6 ROIs) (128)
				250 – 360 ms ^{PD} (<i>eCd</i> =0.74, <i>Pw</i> =0.71)	<u>temporal right ROI (fearful)</u> ; central ROI (happy) ^{PH} ; (6 ROIs) (128)
(Farkas, Stefanics, Marosi, & Csukly, 2015) 2	patients 28 (12 F, ϕ 37.7 \pm 8.4 y) CG 27 (12 F, ϕ 38.2 \pm 10.6 y)	shape: horizontal and vertical high frequency grating	smaller vMMN in patients	90 – 200 ms ^{PD} (<i>eCd</i> =0.76, <i>Pw</i> =0.79)	bilateral and sagittal occipito- parietal ROIs (negativity vMMN) and right and sagittal prefrontal ROIs ^{PD} ; (6 ROIs) (128)
(Neuhaus, Brandt, Goldberg, Bates, & Malhotra, 2013) Schizophrenia 3	patients 22 (10F, ϕ 40.67 \pm 11.3 y), CG 24 (11F, ϕ 37.96 \pm 7.3 y)	shape: X or O	smaller vMMN in patients	250 – 350 ms, ^{PD} (<i>eCd</i> =1.05, <i>Pw</i> =0.93)	bilateral inferior temporo- occipital areas ^{PD} ; (2 ROIs) (64)
(Urban, Kremláček, Masopust, & Libiger, 2008) Schizophrenia 4	patients 24 (5F, 27.90 ± 9.25 y), ASm CG	motion direction: periphery of visual field	smaller vMMN in patients	100-200 ms ^{PH} (<i>eCd</i> =0.76, <i>Pw</i> =0.73)	Oz, O2, Pz, Fz ^{PH} (6)
Mood disorders					
(Y. Chang, Xu, Shi, Zhang, & Zhao, 2010) Major Depressive Disorder 5	exp. 1 patients 15 (9F, ϕ 40.3 \pm 11.2 y), Am CG 15 exp. 2 patients 10 (7F, ϕ 44.0 \pm 15.8 y), ASm CG 10	faces: schematic – neutral: standard; happy, and sad: deviants	smaller or absent vMMN for patients; no modulation by inverted face position in depression;	120-200 ms ^{PD} (<i>eCd</i> =0.97, <i>Pw</i> =0.73 for exp. 1)	TP7/TP8, M1/M2, P7/P8, O1/O2 ^{PD} ; (32)
				220-320 ms ^{PD} (<i>eCd</i> =1.20, <i>Pw</i> =0.89 for exp. 1)	TP7/TP8, M1/M2, P7/P8, O1/O2 ^{PD} ; (32)
(Y. Chang et al., 2011) Major Depressive Disorder 6	patients 14 medication-free (5F, 41.4 \pm 12.6 y), ASm CG	shape: 1 or 2 bars	NO reduction in the oddball-vMMN (deviant- standard); decreased <u>deviant- control</u> response for patients	150-250 ms ^{PD} (<i>eCd</i> =1.06, <i>Pw</i> =0.77)	T5/T6, O1/O2; ^{PD} (32)
				250-320 ms ^{PD} (<i>eCd</i> =0.87, <i>Pw</i> =0.60)	T5/T6, O1/O2; ^{PD} (32)

condition: windmill deviants of different deviant frequency	smaller and less lateralized vMMN response for patients	200-350 ms ^{PD} (<i>eCd</i> =2.05, <i>Pw</i> =1.00)	right temporo-occipital area ^{PD} ; (128)	YES: smaller vMMN at Oz was related to higher lithium dosage (mood stabilizer)
condition: presentation time of deviants	smaller vMMN amplitude to <u>long duration deviants</u> in patients, NO difference for short deviants	200-250 ms ^{PD} (<i>eCd</i> =0.71, <i>Pw</i> =0.59)	O1, O2, Oz; ^{PD} (32)	NO: no correlation to depression severity
condition: bars	larger vMMN for nicotine in O1 and O2	Whole interval (up to 900 ms) (<i>eCd</i> =0.82, <i>Pw</i> =0.84)	O1, O2 ^{PD} (15)	not assessed
condition: location and duration: two color patches	alcohol reduced vMMN amplitude for location and duration deviants, not for color deviant	150-230 ms ^{PD} (location) (<i>eCd</i> =1.08, <i>Pw</i> =0.43) 150-280 ms ^{PD} (duration) (<i>eCd</i> =0.94, <i>Pw</i> =0.34)	left parietal-occipital electrode cluster (P1, P3, P5, PO3, PO5) and the right parietal-occipital electrode cluster (P2, P4, P6, PO4, PO6) ^{PD} (64)	not assessed
condition direction: robustness of visual deviants	abuse duration dependency: vMMN was higher for short term abusers and lower for long term abusers	120-240 ms ^{PD} (<i>eCd</i> =0.86, <i>Pw</i> =0.68)	Pz (6)	YES: vMMN was smaller with increase of abuse duration (O _L , O _R , Oz, Pz) or age (Oz, Pz)
condition: deviant frequency of deviants	vMMN reduction for alcohol compared to placebo condition (3 subjects excluded)	150-170 ms ^{PD} (<i>eCd</i> =0.98, <i>Pw</i> =0.36)	Oz ^{PD} (62)	NO: vMMN unrelated to post- experiment blood alcohol concentration

(Iijima, Osawa, Nageishi, Ushijima, & Iwata, 1995) ¹ Dementia of various etiologies 13	young 20 (20-29y) elder 20 (60-79y) patients 18 (58-78y)	shape: X or O	NO differences in vMMN among groups	No difference (<i>eCd Pw NA</i>)	No difference in Fz, Cz, Pz ^{PD} (3)	N
(Kremláček et al., 2011) SCA type 2 14	patients 10 (5F, 25-72 y) CG 54 (16 F, 19-57 y)	motion direction: periphery of visual field	NO pathological vMMN	100-200 ms ^{PD} (<i>eCd Pw NA</i>)	Oz ^{PD} (6)	Y m o C d
(Tales & Butler, 2006) AD 15	probable AD 8 (8F, ϕ 73; 52-84 y) aged controls 12 (10F, ϕ 73.2; 51-84 y) young controls 11 (8F, ϕ 28; 20-43 y)	shape: 1 or 2 bars	NO differences in vMMN among AD and aged control groups vMMN absent in first 16 trials for patients, larger in amplitude for <u>last 16 trials in the block</u> (as compared to healthy elder)	250-400 ms ^{PD} (<i>eCd=1.93, Pw=0.98</i>)	T5,T6, O1,O2 ^{PD} (13)	n
(Tales, Haworth, Wilcock, Newton, & Butler, 2008) AD, MCI 16	probable AD 10 (8F, ϕ 75.2; 67-81 y) amnesic MCI 8 (2F, ϕ 74.5; 65-82 y) Am CG 10 (6F, ϕ 71.2; 65-81 y)	shape: 1 or 2 bars	early vMMN larger for patients (similar for AD and MCI, second half of testing – not tested for the intergroup difference)	140-250 ms ^{PD} (<i>eCd Pw NA</i>)	T5,T6, O1,O2,Oz, Pz ^{PD} (14)	n
(Stoohart, Kazanina, Näätänen, Haworth, & Tales, 2014) AD, amnesic MCI 17	AD patients 20 (13F, ϕ 79.2; 60-91 y) amnesic MCI patients 25 (9F, ϕ 77.3; 62-91 y) controls 26 (12F, ϕ 76.0; 62-88 y)	shape: 1 or 2 bars	smaller vMMN (or even positivity) in <u>aMCI compared to controls</u> NO differences in vMMN between AD and controls	147-213 ms ^{PH} (AD) 146-234 ms ^{PH} (old, <u>aMCI</u>) (<i>eCd=0.91, Pw=0.89</i>)	occipital ROI ^{PD} (averaged O1, Oz, O2, PO9, PO10, PO7, PO8) (64)	Y lo
Developmental disorders (autism, dyslexia, mental retardation)						
(Cléry, Bonnet-Brilhault, et al., 2013) ASD 18	patients 12 (2F, 11.6 \pm 1.8 , 8-14 y) ASm CG	shape, motion: circle deformation	earlier vMM in patients, In CG occipito-parieto-temporal negativity around 330 ms and positivities around 280 and 450ms, in ASD several positivities in 50-300 ms	300-410 ms ^{PH} (<i>eCd Pw NA</i>)	occipito-parieto-temporal sites ^{PH} (29)	n
				400-600 ^{PH} (<i>eCd Pw NA</i>)	T4 ^{PH} (29)	

¹ Only abstract is available

			followed by a occipito-parietal positivity around 400 ms	500-600 ms ^{PH} (eCd Pw NA)	T4, C4, CO2, M2, Iz, T6 ^{PH} (29)	
(Cléry, Roux, et al., 2013) ASD 19	patients 13 (2F, 26.2 ± 5.0 y) CG 13 (5F, 24.3 ± 2.0 y)	shape, motion: circle deformation	NO vMMN in ASD, late positivity instead	180-240 ms ^{PH} (negativity in CG) (eCd Pw NA)	occipito-parietal electrodes ^{PH} (29)	n
				210-250 ms ^{PH} (negativity in CG) (eCd Pw NA)	fronto-central sites ^{PH} (29)	
				350-600 ms ^{PH} (positivity in AD) (eCd Pw NA)	occipito-temporo-parietal sites ^{PH} (29)	
(Gayle, Gal, & Kieffaber, 2012) Autism Spectrum Quotient in controls 20	CG 45 (16 F, 19.8 ± 1.7 y)	faces: happy and sad compared to neutral	not assessed	150-425 ms ^{PD} (eCd=0.73, Pw=0.52)	Parieto-occipital electrodes (PO3, PO4, PO7, PO8) ^{PD} (72)	Y O su fa
(Horimoto, Inagaki, Yano, Sata, & Kaga, 2002) Children with mental retardation 21	patients 10 (5F, ϕ 11.2; 9-15 y) Am CG 12 (5F, ϕ 10.0; 7-13 y) adult CG 11 (9F, ϕ 28.5 y)	color: blue, red or greenish blue square	different distribution of vMMN	160 -400 ms ^{PH} (eCd Pw NA)	Pz, Oz in control children, Pz, Cz in patients ^{PH} (4)	n
(Maekawa et al., 2011) High-functioning ASD 22	patients 11 (3F, ϕ 28, 18-40 y) CG 11 (7F, ϕ 28,9, 20-38 y)	shape: windmill patterns of different spatial frequency	NO differences in vMMN between groups	150-350 ms ^{PD} (eCd Pw NA)	Oz ^{PD} (128)	n
(Wang, Bi, Gao, & Wydell, 2010) Developmental Dyslexia (DD) 23	patients 11 (4F, ϕ 10.8 ± 0.8 y) age matched CG 12 (6F, ϕ 10.5 ± 0.5 y) Reading-level matched CG 13 (9F, ϕ 9.2 ± 0.3 y)	motion direction: low contrast, low spatial frequency (magnocellular) or high contrast, high spatial frequency (control)	smaller vMMN for DD as compared to both control groups	150-250 ms ^{PH} (magnocellular motion) (eCd=1.15, Pw=0.77)	Oz ^{PD} (32)	n
			NO difference for control motion between groups	200-300 ms ^{PH} (control motion) (eCd Pw NA)	Oz ^{PD} (32)	
Aging/Maturation						
(Cleary, Donkers, Evans, & Belger, 2013) Maturation 24	children 22 (13F, ϕ 10.4 ± 1.43, 8 - 12 y), 21 in the final analysis adults 20 (10F, ϕ 26.6 ± 5.65, 18 - 42 y)	pattern: spatial frequency of horizontal gratings	Lower vMMN amplitude in children	Around 150 ms (eCd=0.72, Pw=0.61)	O1, O2 ^{PH} (O1, O2, F3, F4 out of 13)	n
			Higher positive amplitude in children	Around 250 ms (eCd=0.70, Pw=0.59)	F3, F4 ^{PH} (O1, O2, F3, F4 out of 13)	

(Cléry et al., 2012) Maturation 25	children 12 (2F, ϕ 11.3 \pm 1.6; 8-14 y) adult controls 12 (5F, ϕ 24 \pm 2.2; 20-30 y)	shape, motion: circle deformation	vMMN with three peaks (150-350 ms, posteriorly) and later positivity in children, in adults occipito-parietal negativity (180-240 ms) followed by fronto-central negativity (210-250 ms) ^{PH}	130-160 ms ^{PH} (more negativity in children)	Temporo-parietal sites ^{PH} (29)
				250-310 ms ^{PH} (more positivity in children)	Fronto-central sites ^{PH} (29)
				350-400 ms	FC1, FT3, FC2, FT4, C3, T3, TP3 ^{PH} (29)
				400-600 ms	C4, T4, CP2, TP4, P4, T5 ^{PH} (29)
(Horimoto et al., 2002) Maturation 26	control children 12 (5F, ϕ 10.0; 7-13 y) adult controls 11 (9F, ϕ 28.5 y)	color: blue, red or greenish blue square	more often 2 peaks in children, early peak with larger amplitude in children	180-400 ms ^{PH} (<i>eCd Pw NA</i>)	Fz, Cz, Pz, Oz ^{PD} (4)
(Tanaka, Okubo, Fuchigami, & Harada, 2001) Maturation 27	newborns 83 (35-43 weeks of conceptional age, incl 11 potentially with cognitive dysfunction), vMMN recorded from 63	depth/shape: Ramachandran patterns (convex/concave)	vMMN latency shortened during maturation	Latency decreased from 519 ms to 418 ms between 35 and 37 week of conceptional age (<i>eCd=2.15, Pw=0.83</i>)	Pz ^{PD} (4)
(Tomio, Fuchigami, Fujita, Okubo, & Mugishima, 2012) Maturation 28	107 normal subjects (52 F, 2 - 27y)	depth/shape: Ramachandran patterns (convex/concave)	vMMN latency shortened during maturation	Latency decreased from 2-3 y (394 \pm 58 ms) until 16 y (273 \pm 32 ms) (<i>eCd=2.73, Pw=0.99</i>) stable (275 \pm 44 ms) for age >18 y	Pz ^{PD} (4)
(Iijima, Osawa, Nageishi, Ushijima, & Iwata, 1996) Aging 29	elders 20 (12F, 60 - 79 y) young 20 (11F, 20 - 29 y)	shape: X or O	NO vMMN difference (peak, latency); later N2b difference in latency considered as attention related;	NO difference in 70-250 ms ^{PH}	No difference in Fz, Cz, Pz ^{PD} (3)
(Lorenzo-López, Amenedo, Pazo-Alvarez, & Cadaveira, 2004) Aging 30	elders 9 (5F, ϕ 62.0 \pm 3.0, 58 - 67 y) middle-aged 5 (2F, ϕ 49.0 \pm 4.0, 45 - 54 y) young 7 (4F, ϕ 32.0 \pm 6.0, 24 - 38 y)	motion direction: up or down	Smaller average amplitude with age	165-205 ms ^{PH} (<i>eCd=0.94, Pw=0.41</i>)	Regardless of age maximal at occipital and temporal sites (20)
			Smaller amplitude in older group	145-165 ms ^{PH}	No vMMN in O1, Oz, T5, P3 ^{PH} in older group (8 posterior electrodes)
(Stothart, Tales, & Kazanina, 2013) Aging 31	elders 17 (8F, ϕ 76.8 \pm 6.1, 66 - 86 y) young 17 (10F, ϕ 20.8 \pm 3.1, 18 - 31 y)	shape: 1 or 2 bars	NO vMMN amplitude or duration difference	138-264 ms (<i>eCd=0.17, Pw=0.08</i>)	No vMMN differences in occipital ROI (7 posterior electrodes from 64)
(Tales, Troscianko, Wilcock, Newton, & Butler, 2002) Aging	elders 12 (8F, ϕ 77, 69 - 88 y) young 24 (16F, ϕ 30.5 y)	shape: 1 or 2 bars	smaller average amplitude in elders	250 - 400 ms ^{PD} (<i>eCd=0.97, Pw=0.81</i>)	T5, T6, O1, Oz, O2 ^{PH} (14)

32					
(Tales & Butler, 2006) Aging 33	aged controls 12 (10F, ϕ 73.2; 51-84 y) young controls 11 (8F, ϕ 28; 20-43 y)	shape: 1 or 2 bars	smaller average amplitude in elders	250-400 ($eCd=1.18$, $Pw=0.77$)	T5,T6, O1,O2 ^{PD} (13)
Miscellaneous					
(Bottari et al., 2014) Deafness 33	patients 12 (7F, 35.9 \pm 3.0 y) Controls 12 (6F, 31.6 \pm 2.2 y)	shape, motion: circle deformation	smaller amplitude for patients in posterior and larger in central area	153 – 292 ms ^{PH} ($eCd=1.93$, $Pw=0.99$)	left-right x frontal, central, posterior clusters ^{PD} (34)
(Si et al., 2014) Hypertension 34	patients 15 (7F, 55.3 \pm 7.5 y) ASEm CG	duration: onset of squares rectangles in periphery	vMMN smaller in patients vMMN delayed in patients	cca150 -300 ms ^{PD,PH} ($eCd=0.90$, $Pw=0.66$) 216 ms for CG and 240.8 ms for patients ($eCd=0.90$, $Pw=0.66$)	TP7, A1, A2, O1, Oz ^{PH} (main effect for 7 PD electrodes out of 32) temporal-occipital regions, O1 ^{PH} (main-effect for 7 PD electrodes out of 32)
(Tang et al., 2013) Panic disorder 35	patients 12 (8F, 47.8 \pm 10.7 y) Controls 17 (11F, 39.4 \pm 14.7 y) comparable in ASE and handedness	faces: schematic - positive, negative, neutral	vMMN smaller for positive/negative emotions in patients	220-330 ms ^{PD,PH} ($eCd=0.92$, $Pw=0.67$)	TP7/TP8 and O1/O2 ^{PH} (also M1/M2 and T5/T6 contributed to main-effect, altogether 32 electrodes)

Notes: AD - Alzheimer disease, Am – age matched, ASD - Autism Spectrum Disorder, ASE - age, sex, education, respectively, ASEm – age and sex matched, CAG - cytosine–adenine–guanine, CG – control group, DD – developmental dyslexia, eCd – equivalent Cohen’s *d* (see *vMMN studies*), MCI – Mild Cognitive impairment, MMSE- Mini-Mental State Examination, NA – not available, PD – pre-determined in difference was looked at, total number of electrodes used in parentheses, PH – post-hoc difference in interval or electrodes, Pw – statistical ataxia type 2, vMM – visual mismatch.

positive symptoms like delusions, hallucinations and thought disturbances, and negative symptoms such as blunted emotions, social withdrawal, catatonic behavior and lack of spontaneity. Furthermore, cognitive impairments are widely considered to lie at the core of the illness (Buchanan et al., 2005) and include symptoms of deficits in attention, memory, and executive functions (Heinrichs & Zakzanis, 1998; Barch & Ceaser, 2012). Prevalence of schizophrenia is 0.6 % in the general population (McGrath, Saha, Chant, & Welham, 2008), it requires long-term treatment and is associated with a decreased life expectancy. In spite of the advances in neurobiological research (structural, functional or genetic) the pathophysiology of this heterogeneous disease is not fully understood. For over 40 years of research on the neurobiology of schizophrenia, dopamine dysregulation emerged as one of the most robust findings, which is currently attributed to elevated dopamine synthesis capacity in the disease (Fusar-Poli & Meyer-Lindenberg 2012)². However, the dopamine dysregulation hypothesis does not explain cognitive deficits and negative symptoms. The current view, beside a dysregulation of the dopaminergic system, also incorporates misbalanced glutamatergic and GABAergic (Gamma-aminobutyric acid) neurotransmitters (Marsman et al., 2013)³ mediating cortical processing (pyramidal cells/interneurons) and supporting the notion of aberrant sensory processing (Lee et al., 2012; Lewis, 2014). In fact, interactions of glutamatergic NMDAR (N-Methyl-D-Aspartate receptor) and dopamine receptors are a key mechanism in multiple pathophysiological theories of schizophrenia (Frohlich & Van Horn, 2014).

NMDARs play a key role in synaptic transmission and in the synaptic plasticity underlying fundamental cognitive functions such as learning and memory (Zorumski & Izumi, 2012) and it has been hypothesized that modulation of plasticity underlying predictive representations is abnormal in schizophrenia (Javitt, 2004; Stephan et al., 2006, 2009; Moghaddam & Javitt, 2012). Beyond predictive processes, there is now much evidence for a general hypofunction of NMDARs in schizophrenia (Umbricht & Krljes, 2005; Stahl, 2007; Heekeren et al., 2008; Coyle, 2012; Javitt, 2012). Despite much research over the past decades, we still lack sensitive and non-invasive laboratory tests in psychiatry that predict whether an at-risk individual will transition to psychosis. There is a general consensus (Luck et al., 2011) that new biomarkers providing reliable and sensitive measures of neuro-cognitive functioning could be used to predict clinical course in prodromal individuals before the onset of psychosis. Thus, ideally, early intervention would become possible. Furthermore, such biomarkers would dramatically facilitate the development of treatments for cognitive dysfunction in mental illnesses.

One of the promising candidate ERP components is the auditory mismatch negativity (MMN) (Näätänen, et al., 2011) which has emerged as one of the most reliable electrophysiological

² Support for the dopamine hypothesis is mostly based on an effect of dopamine receptor antagonists in treatment of positive schizophrenia syndromes – i.e. delusions, hallucinations, disorganized speech/behavior, and on an observation that drugs increasing dopamine (amphetamines, levodopa) levels induced psychotic symptoms in healthy volunteers.

³ The main support for the glutamate hypothesis comes from experiments with ketamine, NMDAR antagonist widely used in subanesthetic doses as pharmacological model of schizophrenia due to its psychomimetic properties (Pomarol-Clotet et al., 2006; Corlett et al., 2011; Kocsis et al., 2013).

forming probabilistic representations of sensory signals (Friston, 2010; Stefanics & Czigler, 2012; Stefanics et al., 2014). MMN deficits are one of the features in schizophrenia that indicate severe abnormalities in fundamental brain processes of prediction and inference (Stephan et al., 2006; Corlett et al., 2007; Friston et al., 2014). This is further corroborated by parallel evidence for a key role of NMDAR in MMN generation and in the pathophysiology of schizophrenia (Coyle, 2006; Umbricht & Krljes, 2005; Javitt, 2009).

Although vision *per se* is not predominantly deteriorated in schizophrenia, numerous visual deficits have been identified, such as deficits in lateral inhibition, smooth eye movements, motion detection, contrast sensitivity, and recognition of faces and facial-expressions (Butler, Silverstein, & Dakin, 2008; Yoon, Sheremata, Rokem, & Silver, 2013). Furthermore, visual hallucinations are present in about 27% of schizophrenia patients (Waters et al., 2014). Magnetic resonance spectroscopy showed that increased levels of glutamine and decreased levels of glutamate in schizophrenic patients were not restricted to the frontal regions, but were evident also in the occipital lobe (Chang et al., 2007; Keshavan et al., 2009). Structural changes in the primary visual cortex have been found post-mortem (Dorph-Petersen et al., 2009). The relationship between the glutamate dysregulation and vision was confirmed in a human study administering the NMDA-receptor inhibitor ketamine. During ketamine infusion, subjects had a decreased aMMN amplitude and also impaired processing of visual context (Umbricht et al., 2000).

In the first vMMN report in schizophrenia, Urban et al. (2008) used motion-onset stimuli in the visual periphery while subjects attended to a motion-detection task in the central visual field. The authors reported a smaller vMMN to changes in motion direction in patients with schizophrenia (for details see Tab. 1) than in age- and gender-matched controls. The cumulative vMMN amplitude at the 100-200 ms interval was smaller in the occipital, parietal and frontal derivations than that of controls. The vMMN deficit was observed only in patients with deficit symptoms (assessed by Schedule for the Deficit Syndrome), a finding which corresponds to the glutamatergic hypothesis and early sensory deficit. However, further analyses showed that there was an association of the vMMN impairment with a higher dose of medication, a lower global assessment of functioning (GAF) score or the duration of the illness. Furthermore, there were no differences between patients with illness duration shorter than 3 years and controls. These findings support the view of the motion-onset vMMN as a state marker of schizophrenia.

In a recent study Farkas et al. (2015) used rare changes in the orientation of Gabor patches to elicit vMMN in a group of patients with schizophrenia, and in age- and education-matched controls. Controls had significant vMMN in the occipito-temporal region and showed a positive deviance detection activity (vMMP) in prefrontal region, both within an interval 90–200 ms after stimuli onset. In contrast to the control group, patients did not exhibit significant vMMN or vMMP and showed a lower absolute response amplitude in parieto-occipital and prefrontal areas (Fig. 1). The authors did not find any relationship between patient's deviance detection activity and age, illness duration, equivalent dose medication, and their functional status. While the impairment of an early deviance processing resembles results reported by Urban et al. (2008), the absence of any relationship to the

Beside changes in low-level visual paradigms, more complex natural stimuli such as facial expressions have been used in order to assess automatic change detection in schizophrenia (Csukly et al., 2013). The authors compared a group of patients with age-, gender- and education-matched controls. Four different faces were simultaneously displayed in the visual periphery showing either fearful or happy expressions. To minimize attentional effects in the processing of the face stimuli participants engaged in a centrally presented task which involved detection of changes in the orientation of a fixation cross. VMMN elicited by changes in facial emotions was observed in two time intervals, at 170-220ms and 250-360ms post-stimulus, over occipital, right temporal and central scalp regions. Patients had a significantly smaller vMMN in the left temporal area in the earlier interval for the happy deviant faces and in the right temporal area in the later interval for the fearful deviant faces. The groups showed different responses also in the central area for the happy deviant faces in both intervals, however, the mismatch activity (“deviant – standard”) observed here was positive and larger in controls. The authors found no correlation between the mismatch response and medication or symptoms severity, which supports the view that the impairment in the automatic change detection of peripherally presented emotional expressions might be a trait rather than state marker. A significant correlation was found between the activity in the central electrode sites for the happy deviant faces and emotion recognition performance.




Fig. 1. Scalp areas of a significant amplitude difference in the deviance-related activity between schizophrenic patients and controls. Numbers represent the following studies according to Table 1: 1-Csukly et al. (2013), 2-Farkas et al. (2015), 3-Neuhaus et al. (2013), and 4-Urban et al. (2008). Highlighted areas represent ERP effects as measures on the scalp. About the limited correspondence between scalp potential maps and cortical sources, see the Technical notes section at the end of the paper.

Thus the stimuli eliciting the MMN were also task-relevant, which renders interpretation of the results ambiguous, as differential activity observed in this study might be related to predictive processes, attention, or executive processes (for the role of attention in the vMMN generation, see Czigler, 2007; Stefanics et al., 2014).

Summary of vMMN studies in schizophrenia

While there are only four vMMN studies yet in patients with schizophrenia, their results unanimously point to impaired predictive processes in vision. This is well in line with deficits in auditory MMN which have been widely explored for more than three decades, and since the first report of an aMMN deficit (Shelley et al., 1991), there have been **256** articles published addressing this issue (source PubMed, "*mismatch negativity*" schizophrenia, May 2015). Most of these studies showed an aMMN amplitude reduction in schizophrenia (Todd, Michie, Schall, Ward, & Catts, 2012; Umbricht & Krljes, 2005) and, moreover, this sensory deficit was associated with the impaired daily functioning of the patients (Todd, Harms, Schall, & Michie, 2013). The aMMN sensitivity to the sensory deficit depends on the type of deviance used, e.g. pitch, intensity, or duration, with the duration violation being the most sensitive test as shown by a meta-analysis of 32 reports (Umbricht & Krljes, 2005). Correlations between aMMN and clinical measures have been mostly weak when prodromal population and patients with the first hospitalization criteria or patients' healthy relatives have been studied (Todd et al., 2013). Generally, the numerous reports on aMMN elicited by duration or pitch deviance show a consistent picture, whereas vMMN findings are mosaic-like, since in the visual domain there have only been a few studies, each using a specific design.

The NMDAR hypofunction hypothesis is perhaps the most powerful model of schizophrenia, with a strong translational potential. Currently the aMMN component is widely used as a proxy for NMDAR deficits in schizophrenia. Although we still lack experimental evidence that deficits in visual MMN are mediated by NMDAR hypofunction similarly to aMMN deficits, there is compelling evidence that vMMN is also impaired in schizophrenia. Direct comparison between the aMMN and vMMN in schizophrenia is thus warranted. Future studies should investigate heterogeneity in schizophrenia and consider dividing patients into subgroups along behavioral, genetic or molecular endophenotypes as well as variability in vMMN. Using experimental paradigms with appropriate control for attention and stimulus probability effects (Stefanics et al. 2014) will help reduce variability among studies due to confounding variables. In summary, exploration of vMMN as a research tool, or as a potential biomarker in schizophrenia is highly justified.

2.2 Mood disorders

Major depression disorder (MDD) is one of the most common forms of psychopathology. It affects approximately one in six men and one in four women at least once during their lifetimes (Kessler, 2012; Kessler et al., 2003), and its 12-month prevalence is approximately 5.5 % (Bromet et al., 2011). The most frequent symptoms in MDD include depressed mood, sleeping problems, fatigue and suicidal thoughts or intentions, whereas bipolar disorder (BP), another diagnosis with disturbance in mood, is characterized by recurring depressive and manic periods (American Psychiatric Association, 2013). Both theoretical (Beck, 1967, 1976) and empirical (Mineka & Sutton, 1992) work suggests that a cognitive bias is an important feature of mood disorders. According to Beck's cognitive theory of depression (1967, 1976), dysfunctional attitudes, negative schemata, and faulty information processing are critical factors that predispose individuals to experience episodes of depression. The neural mechanisms underlying disturbances in cognitive functions are poorly understood, and they may be present even at the level of automatic information processing (for a review, see Beck, 2008). Therefore several studies used aMMN (or its magnetic counterpart, MMNm) to study automatic sensory-cognitive processes in depression (Kähkönen et al., 2007; Lepistö et al., 2004; Iv, Zhao, Gong, Chen, & Miao, 2010; Pang et al., 2014; Takei et al., 2009) and bipolar disorder (Andersson, Barder, Hellvin, Løvdahl, & Malt, 2008; Takei et al., 2010).

Takei et al. applied magnetoencephalography to measure information processing deficit in depression (2009) and bipolar disorder (2010). In the study of depressive patients, MMNm was elicited in response to duration and frequency changes of pure-tone stimuli and in response to an across-category change in a vowel. The magnetic global field power (mGFP) of the MMNm was significantly smaller in the group of depressive disorder patients than in the healthy control group. Kähkönen et al., (2007) found that the aMMN amplitude in the EEG (but not its magnetic counterpart) was increased in depression compared to controls. It is not clear why both amplitude increase and decrease of MMN have been found related to depression. One possibility is that the contradictory results can be explained by differences in stimulus conditions or patient populations.

When patients with bipolar disorder were studied using an aMMN paradigm, no amplitude differences were found, but mGFP of MMNm in the right hemisphere for pure-tones was delayed in patients with bipolar disorder compared to healthy participants (Takei et al., 2010). In addition, the MMNm dipole in the left hemisphere was located inferiorly in patients with bipolar disorder compared to that in healthy group.

Alterations of MMN did not correlate with clinical symptoms in either of the patients groups. The lack of correlation between aMMN (Kähkönen et al., 2007; Pang et al., 2014; Takei et al., 2009) and depression symptoms suggests that sensory-cognitive deficit in depression is a trait- not a state-dependent phenomenon.

attended to changes in the color of a square in the fixation point. The vMMN was compared between the groups at two latency ranges: 150-250 ms and 250-320 ms after stimulus onset. No group differences were found in response amplitude. On the other hand, the late vMMN amplitude correlated with the Hamilton rating scale for depression in the depressed group. However, although vMMN did not differ between the groups, they showed differential responses to the deviant and control stimulus (the same stimulus presented in an equiprobable control condition). This pattern of results does not provide a straightforward interpretation, but presumably there is a deficit in the processing of low-level visual features, and perhaps also in the detection of regularity violations, in depression.

In a study with depressed participants, Qiu et al. (2011) presented two black squares simultaneously for either 50 ms or 150 ms in duration. Both an increment and a decrement in duration were used in separate stimulus blocks as the deviant stimulus. Participants attended to changes in the size of a black cross in the center of the screen. The vMMN was calculated as a difference between responses to physically identical stimuli presented in the two stimulus blocks (i.e. responses to the 150- ms stimulus as deviant minus responses to 150- ms standard stimulus). In the 200-250 ms post stimulus analysis time window the mean amplitude of the differential response for the 150 ms duration stimulus was smaller in the depressed patients than in the controls. However, details of the analysis of standard and deviant responses were not reported, leaving it open whether the standard and deviant responses differed from one another in either of the groups.

Maekawa et al. (2013) used changes in spatial frequency of circular black-white windmill patterns as vMMN evoking stimuli in a group of bipolar disorder patients and healthy controls. Participants were instructed to listen to a story presented through earphones, and also attend to visual stimuli and to press a button whenever a target stimulus appeared. It is worth noting, since the visual stimuli were also attended, that the study was not a typical vMMN study where visual stimuli are usually ignored. A six-vane and 24-vane stimuli were assigned as deviant and standard stimuli counterbalanced across the stimulus blocks. The vMMN at 200-350 ms latency range (mean amplitude values at the occipital electrodes) was right lateralized in the control group, but not in the bipolar disorder group (Fig. 2). This finding suggests that the source of the activity is different between the groups. However, it was not reported whether the observed vMMN response was statistically significant in either of the groups. The response amplitude for the differential response at the 200-350 ms latency range correlated with the lithium dosage, but not with the symptoms.




Fig. 2. Scalp areas of a significant difference in the deviance-related activity between patients with mood disorders and controls. Numbers represent the following studies according to Table 1: 5-Chang et al. (2010), 6-Chang et al. (2011), 7-Maekawa et al. (2013), and 8-Qiu et al. (2011).

Both depression and bipolar disorder are also associated with abnormalities in emotion processing (for a review, see Phillips et al., 2003). Visual MMN to changes in facial expressions has been reported several times in healthy participants (Astikainen, Cong, Ristaniemi, & Hietanen, 2013; Astikainen & Hietanen, 2009; Kimura, Kondo, Ohira, & Schröger, 2012; Li, Lu, Sun, Gao, & Zhao, 2012; Stefanics et al., 2012; Susac, Ilmoniemi, Pihko, & Supek, 2003; Zhao & Li, 2006). Because both depression and bipolar disorder are associated with abnormalities in emotion processing, and since depression affects social cognition (Wolkenstein, Schönberg, Schirm, & Hautzinger, 2011), vMMN seems to be a feasible tool to study altered emotion processing in this disorder. It is thus surprising that there is only one report on vMMN to facial expressions in depressed participants (Chang et al., 2010).

Chang et al. (2010) presented schematic faces to depressed and control participants. Neutral standard faces were infrequently replaced by sad or happy faces, both drawn in red. Participants were instructed to count the target faces drawn in green. In two separate stimulus blocks, upright and inverted faces were presented. In the control group, emotional faces elicited larger responses relatively to neutral faces at posterior electrode sites. These differential responses were observed at the latencies of the N170 (early vMMN) and P250 (late vMMN) components. Both early and late vMMN were decreased in the depressive group indicating a deficit in automatic change detection in facial emotions in depression. Face inversion decreased the amplitude of vMMN in control group, but not in the patient group. This may indicate difference in high-level perceptual processing in depression and/or different connectivity in the face processing-related neural network (Matsuyoshi et al., 2015).

In addition to emotional visual stimuli, emotional auditory stimuli have been used to study information processing dysfunction in depression. Pang et al. (2014) found that sad prosodies in meaningless words did not elicit aMMN in comparison to neutral prosody in depression patients while no differences were found for happy or angry prosody in comparison to healthy controls. The aMMN did not correlate with depressive symptoms.

disorders. These alterations are observed as decreased response amplitude (e.g. Kähkönen et al., 2007; Takei et al., 2009), delayed latency and different source location of the change detection-related activity (Takei et al., 2010). The previous studies in depressive participants have not found relationship between clinical symptoms and aMMN (or aMMNm) amplitude or latency (Kähkönen et al., 2007; Takei et al., 2009).

One general aspect comes forth in all the vMMN studies related to mood disorders. Namely, standard and deviant stimulus responses were not compared directly (optimally by applying stimulus type factor in a multivariate statistical model). It is thus unclear whether the vMMN, i.e. differential response, was statistically significant. This methodological issue should be taken into account in future studies. In addition, future (vMMN) studies should explore whether processing deficit in the visual modality in depressed patients is related to visual feature processing, deviance detection, and/or emotional processing deficits.

Studies on vMMN in mood disorders could be useful in the future to disentangle different aspects of the cognitive dysfunction in these disorders. Especially studies using elementary visual features (e.g. orientation or color changes), abstract or rule-like changes (e.g. Kecskés-Kovács et al., 2013) and perhaps also emotional changes like changes in facial expressions in same participant groups would be very informative. In a long run these vMMN studies can be applied for developing diagnostic tools.

2.3 Substance abuse

While the auditory MMN has been extensively used to study different forms of substance abuse (Näätänen et al., 2012), there are only four papers so far concerning substance abuse effects on the visual MMN. In auditory modality, tens of different substances have been investigated, starting from alcohol, caffeine, nicotine and opioids, ending in various drugs for treatment of psychiatric disorders (benzodiazepines, antipsychotic/neuroleptic drugs, GABA agonists/antagonists, adrenergic drugs, dopamine, serotonin, NMDA/glutamate), antihistamines and hormones/neuropeptides (Näätänen et al., 2012). In relation to vMMN, there are two studies using alcohol (Kenemans, Hebly, van den Heuvel, & Grent-'T-Jong, 2010; He, Hu, Pakarinen, Li, & Zhou, 2014), one study on nicotine effects (Fisher, Scott, Shah, Prise, Thompson, & Knott, 2010), and one study investigating metamphetamine (MAP) effects on visual processing (Hosák, Kremláček, Kuba, Libiger, & Čížek, 2008). Two robust patterns emerge from these studies in accordance with previous studies on the enhancing or attenuating effects of those substances on visual information processing (see e.g. Newman et al., 1997 for alcohol; Houlihan et al., 2001 for nicotine; and Kremláček et al., 2008 for metamphetamine). First, moderate acute dosage of alcohol diminishes the vMMN response to unattended and unpredicted stimuli. Second, both nicotine and MAP use enhance the vMMN response (for MAP, less than 5 years of use, see below for more details).

supported by studies showing reduced inhibition to visual stimuli, e.g., in a stop-signal task (Fillmore & Vogel-Sprott, 1999); when the subject is asked to perform more difficult dual-tasks (Fillmore & Van Selst, 2002); or when the central visual target stimulus is being flanked by response-incongruent stimuli (Eriksen flanker task, see Bartholow et al., 2003) (cf. Kenemans et al., 2010). Impairments of attention processing after alcohol consumption have been shown to be reflected in diminished amplitudes of ERP components N1, N2 (see Bijl et al., 2005, for further references for alcohol-dependent subject groups) and P3 (e.g. Houlihan et al., 2001). The vMMN recordings may therefore yield insight into the source of the increased risk: the automatic attention-switching mechanism for visual change detection is dampened even by minor alcohol doses.

Kenemans, Hebly, van den Heuvel, & Grent-'T-Jong (2010) used a visual oddball paradigm (9:1 standard to deviant ratio) with spatial frequency change as a peripheral deviant, while the subjects performed a visual discrimination task in the center of the visual field. They found that in young healthy subjects (university students) with BAC varying from 0.04 to 0.07 % (average 0.05% right before and 0.033% right after the ERP experiment), the vMMN amplitude for peripheral spatial frequency change was considerably reduced in most subjects. In contrast, the exogenous spatial-frequency dependent difference at 80 ms (Kenemans, Baas, Mangun, Lijffijt, & Verbaten, 2000) was unaffected by alcohol. Kenemans et al. (2010) concluded that even a moderate alcohol dosage considered legally acceptable in many countries reduced the sensitivity of the visual cortex to automatic detection of unexpected changes in the visual periphery, whereas alcohol effects are far less dramatic on low-level sensory processes. This is also supported by their behavioral results (as well as the behavioral results of the second vMMN study on alcohol effects by He et al., 2014) showing diminished reaction speed, but intact reaction accuracy in the central task, possibly reflecting stronger focus on the central task-relevant events while peripheral unexpected changes go unnoticed.

However, it is not yet clear, whether the abovementioned conclusions apply to all types of visual stimuli or if there is some feature-specificity to be considered, since another study investigating alcohol effects on the vMMN suggests that stimulus features also plays a role. He et al. (2014) used a new multi-feature vMMN paradigm (Qian et al., 2014; Shi, Wu, Sun, Dang, & Zhao, 2013) with three types of deviant stimuli – duration (60 or 100 ms deviants vs. 20 ms standards), location (45° and 90° change in respect to standard stimulus' placement), and color (red and green deviants vs. sea-green standards). The subjects were attending to the task that involved detecting changes of the size of a cross in the center of the screen, while the vMMN eliciting stimulus sequences were presented in locations surrounding the central stimulus (distance 2.8°). Twelve healthy subjects attended both alcohol consumption (dosage of 0.65 g/kg resulting in average BAC level $0.06 \pm 0.005\%$ before, and $0.048 \pm 0.006\%$ as measured after the experiment) and placebo conditions (separated by two weeks). The results showed that alcohol considerably decreased the vMMN amplitude for changes in peripheral stimulus location and duration whereas that for color change was not affected. The latter might be due to the physical stimulus properties – no reliable vMMN emerged to the green color deviant (compared to the sea-green standard), implying a relatively high level of change is required to elicit the color vMMN at first place (Czigler et al., 2002).

sample (e.g., Ahveninen et al., 1999). Otherwise, He et al., (2014) results are in line with the Kenemans et al. (2010) results, showing decreased the vMMN amplitude after acute alcohol consumption.

Previously, studies of alcohol effects on the aMMN have yielded an analogous data pattern as suggested by the vMMN results of Kenemans et al. (2010) and to some extent the results of He et al. (2014). Even minor doses of alcohol reduced the frontal aMMN subcomponent presumably reflecting attenuated involuntary attention-switching function, whereas the sensory-specific aMMN subcomponent (recorded as polarity reversal in mastoid recordings) was unaffected (Jääskeläinen, Pekkonen, Hirvonen, Sillanaukee, & Näätänen, 1996). More specifically the authors found that the attenuation of the aMMN amplitude even by a small amount of ethanol (0.55 g/kg) resulted from the decrement of the prefrontal rather than that of the supratemporal aMMN subcomponent. Kenemans et al. (2010) also replicated the experiment and the results of Jääskeläinen et al. (1996) showing that alcohol consumption (compared to placebo condition) reduced the amplitude of the frontal component of the aMMN in the 100-200 ms latency window. Consistent with this, a small amount of ethanol decreased the detrimental effect of the task-irrelevant auditory changes on performance accuracy in a visual forced-choice discrimination task, i.e., alcohol improved task performance (Jääskeläinen, Alho, et al., 1996).

Another relatively widely accepted legal substance people use for its effects on mood, arousal and cognition is nicotine. It is proposed that nicotine enhances cognitive performance in smokers and non-smokers probably due to sharpening of primary encoding of sensory and temporal stimuli presented outside the attentional focus (e.g., Fisher et al., 2010, Martin et al., 2009). Nicotine is a cholinergic agonist that may also enhance the glutamatergic system and presumably through this alleviate some symptoms of schizophrenia (see also chapter 2.1 on schizophrenia). For example, it has been shown that nicotine may normalize duration (but not frequency) aMMN in schizophrenics (Dulude et al., 2010). This promising result is already a good justification for any attempts to test the nicotine effects on change detection in other modalities. There is a positive report from a study where acute nicotine effects on the vMMN were investigated by Fisher et al. (2010) in a randomized, double-blind, placebo controlled design in nonsmoking population (n=27). The experimental paradigm had an intermodal design, where the vMMN-eliciting stimuli were vertical bars that differed in their length (long bars being standards, short ones deviants; standard to deviant ratio 3:1). The subjects' task was to attend to and respond to certain tones. Contrary to the authors' hypothesis, nicotine (administered as gum with 6 mg nicotine dosage that typically causes blood nicotine levels to rise 16-26 ng/ml in 25 minutes) did not affect response speed and accuracy measures of a concurrent auditory task. In contrast, the vMMN amplitude for visual deviant events in the center of the field of fixation was dramatically enhanced by nicotine. The authors proposed that nicotine enhances the vMMN amplitude by increasing the ability to encode and process information outside the attended (auditory) modality, such as visual deviant events, while protecting against any decrement in primary task performance.

consequences of acute intoxication (Hart et al., 2001). MAP abuse is related to damage in dopamine and serotonin systems with direct effects on cognition, most notably difficulties in suppressing irrelevant task information but also longer decision times or increased impulsivity (reviewed by Nordahl et al., 2003). Kremláček et al. (2008) tested MAP effects at the cortical level and reported slowing, attenuation, and reorganization of visual evoked potentials (VEPs) in response to pattern reversal of high spatial frequency checkerboards and visual motion in a group of 23 MAP users (with approximately 5 years of MAP abuse). In accordance with these effects, a vMMN study with low-contrast sinusoidal gratings moving fast (50°/s) into opposite directions presented in the visual periphery for 200ms also found that the vMMN amplitude for the motion direction was increased for short-time (less than 5 years) abusers, and decreased only for those who had abused MAP for more than 5 years (Hosak et al., 2008). It is unknown whether similar pattern occurs with aMMN as well, as we found no corresponding study with the auditory mismatch response, but we suggest that it is rather reasonable to look for it in a magnocellular task dealing with rapid temporal information. From a clinical point of view, as vMMN appears to reflect the severity of abuse it makes sense to examine it in addiction, in particular with respect to the possibility of monitoring the extent of sensory functional damage and recovery in cases of withdrawal,. It should be noted that beside automatic irregularity detection also other sensory mechanisms might be of interest, as it is evident from the study investigating VEPs (Kremláček et al. 2008).




Fig. 3. Scalp areas of a significant difference in the deviance-related activity in substance abuse compared to controls or control condition. Numbers represent the following studies according to Table 1: **9**-Fisher et al. (2010), **10**-He et al. (2014), **11**-Hosák et al. (2008) (note that vMMN decrease in >5yr MAP use is not shown), and **12**-Kenemans et al. (2010).

advantage of using vMMN rather than aMMN for studying alcohol effects might be its higher ecological validity, since in everyday situations (e.g. in traffic) people rely on and use visual input more than auditory input. One has to take into consideration, though, that the two existing vMMN studies report the dampening effects of acute alcohol consumption. However, similar changes in vMMN amplitude would be expected in case of chronic alcohol consumption as well, since studies using visual tasks on alcohol-dependent groups show diminished N1, N2 and P3 ERP components (see Bijl et al. (2005) for overview).

The neurobiological and genetic background of addiction and vulnerability has been thoroughly investigated. Using the vMMN component in the above conditions can provide information about the pharmacological mechanisms underpinning visual predictive processes and change detection.

2.4 Neurodegenerative disorders (Alzheimer's disease, mild cognitive impairment and spinocerebellar ataxia)

Alzheimer's disease (AD) is the most common form of dementia (Thies & Bleiler, 2013). It is an insidious, progressive neurodegenerative disorder accounting for 60% to 80% of cases. According to the World Health Organization there were an estimated 25 million people worldwide aged 65 years and older living with the disease in 2012. Characteristic brain pathology includes amyloid plaques, tau tangles, neuronal damage and death, while behavioral characteristics include inexorable decline in cognitive integrity, social interaction, self-care and quality of life. Although drugs cannot yet arrest disease progression, active management and behavioral intervention during the early stages of AD may confer some preservation of function, although short-lived, and improvement in quality of life (Thies & Bleiler, 2013). However, using current clinical test regimes, the very early stages of AD can be difficult to differentiate from cognitively healthy ageing and other factors influencing cognition. Consequently, research is increasingly focusing upon identifying changes in a wider range and level of brain functions (including fundamental levels of processing) than those relatively high-level processes measured at present.

One area of such research involves determining the functional integrity of visual information processing. Research increasingly reveals pathological and functional change in the visual cortex and disruption to a wide range of basic visual processes (see Tales et al., 2008; Stothart et al., 2014, for reviews). Evidence of pathological change in the visual extra-striate cortex (a region closely associated with the vMMN) indicates that abnormality in vMMN and its associated operations in AD should be considered a possibility.

Iijima et al. (1995), in an early study reported in a published abstract the preservation of the vMMN in older adults and in a group of patients with dementing illness of various etiologies. In a later study, Tales and Butler (2006) examined vMMN in cognitively healthy ageing and AD.

appeared with a randomized inter-stimulus interval of 612-642ms for a duration of 200ms. The vMMN amplitude was measured over a 240-400ms period after stimulus onset.




Fig. 4. Scalp areas of a significant difference in the deviance-related activity between patients with neurodegenerative diseases and controls. Numbers represent the following studies according to Table 1: **15**-Tales & Butler (2006), **16**-Tales et al. (2008), and **17**-Stothart et al. (2014), studies without group differences are ignored. Note that patients in studies 14 and 15 showed the vMMN increase only in the second part of the examination.

For the young adults, a significant vMMN was evident in response to both the first and second blocks of trials, whereas the older adult group failed to demonstrate a significant vMMN in response to both blocks. However, although the AD group resembled the older group over the first block of trials, in that they exhibited no significant vMMN, they differed significantly from the older adults in the second block of trials by revealing a robust and significant vMMN (an effect resulting from an AD-related selective increase in the amplitude of the responses to deviants during the second set of trials). The ageing-related reduction in vMMN was described by Tales and Butler (2006) as representative of a reduction in the efficiency of change detection, with the resurgence of the effect in AD possibly indicative of the known AD-related tendency for over-distractibility in response to novel or distracting stimuli and events within the environment. Thus unlike the efficient processing it represents in healthy young individuals, the significant vMMN in AD may reveal dysfunction at the level of automatic vision-related processing. However, the vMMN is not always found to be reduced in cognitively healthy ageing (see the section 2.6 in this review) and although this effect was repeated to some extent in a later study examining AD and mild cognitive impairment (MCI; a prodromal stage of AD for some individuals; Tales et al., 2008) the effects were dependent upon the epoch over which the vMMN was measured. Furthermore, Stothart and colleagues (2014), using the same stimulus paradigm, did not detect statistically significant difference between the

together with inter-study variation in measurement parameters and analysis. Research on AD-specific effects of the vMMN has been further restricted by a lack of longitudinal analysis, which has precluded verification of disease etiology and resulted in a failure to determine within a given group for whom MCI represented the prodromal stage of AD. Other limiting factors include the difficulty of controlling the drug status of participants (both in relation to AD-specific intervention and medication for concurrent illnesses); the relatively small number of participants recruited and tested, the failure to take into account potential covariates such as cognitive reserve (education, intelligence), age, gender and disease stage, participant demographics, and treatment status.

Another disorder affecting visual areas and accompanied by abnormality in some aspects of visual information processing is spinocerebellar ataxia (SCA). This autosomal dominant neurodegenerative disorder is characterized by progressive movement abnormalities and cerebellar atrophy, with the common SCA genotype 2 (SCA2) typically associated with saccadic slowing, tremor and deterioration in attention, short-term memory and executive dysfunction (Kremláček et al., 2011). SCA-related abnormalities in visual processing lead to the prediction that an abnormal vMMN will also be evident in this disorder. Using a motion-related vMMN paradigm (in which the vMMN was calculated as an integral of the difference between the deviant minus the standard ERP within an interval 100-200 ms after stimulus onset), Kremláček et al. (2011) found generally preserved vMMN in SCA2, but also a positive correlation between the vMMN and age and age at SCA onset and a negative correlation between the vMMN and pathological load (CAG repeats - cytosine-adenine-guanine triplet coding protein ataxin-2). As highlighted by Kremláček et al. (2011), and similarly to studies of vMMN in AD, patient numbers were relatively small; a reflection of how difficult recruitment can be in such populations when rigorous inclusion and exclusion criteria are observed. However, despite these limitations, such studies show proof of concept in that such patients can be tested successfully using typical vMMN paradigms and that there is some evidence of potential abnormality in the vMMN in neurodegenerative disorders.

Summary in neurodegenerative disorders

Compared to the vMMN, the auditory mismatch negativity has been widely applied to the study of clinical populations, although once again, there are relatively few studies examining aMMN in neurodegenerative disorders such as AD and in other forms of dementia (see Näätänen et al., 2012 for a review and Riekkinen et al., 1997). Evidence in relation to AD again reveals a somewhat heterogeneous study outcome, which in part appears to reflect the choice of auditory features used to elicit the aMMN (see Näätänen et al., 2012 for a review) but note however that abnormal aMMN has also been reported in the behavioral variant of fronto-temporal dementia (Rowe, Hughes, & Nestor, 2009) and in dementia related to Parkinson's disease (Brønneck, Nordby, Larsen, & Aarsland, 2010).

While there is a relative paucity of studies, there is evidence to suggest that with further investigation and development, the vMMN may represent an important functional characteristic of AD and related disorders. Although in the early studies by Tales and colleagues (see above), the relationship between a measure of cognitive integrity and the vMMN was not attempted, Stothart et

general cognitive integrity as well as reflecting early sensory processing (Näätänen et al., 2011).

It is clear that despite the substantial clinical and social importance of neurodegenerative disorders relatively few studies examining both visual and auditory MMN function have been performed. As well as improving participant numbers, future studies need to examine even earlier stages of AD and MCI, namely subjective cognitive decline (Jessen et al., 2014), together with other forms of dementia. Despite these limitations, such studies show proof of concept in that such patients can be tested successfully using typical vMMN paradigms and that there is some evidence of potential abnormality in the vMMN in neurodegenerative disorders which warrants further investigation. Only when more studies are performed can we work towards a consensus for what constitutes normative MMN data and what factors characterize specific pathological change.

2.5 Developmental disorders

Among the 33 vMMN studies⁴ reviewed here six focused on developmental disorders, including autism spectrum disorders (ASD), dyslexia, and mental retardation. The ASD prevalence in school children is about 2% (Blumberg, Ph, & Bramlett, 2013) and etiology of the disease is not fully understood. In addition to impairment in cognitive and social interaction, sensory activation and habituation are also affected. The sensory impairment is frequently connected with a fascination in stimuli (e.g. visual motion) or fear of them (e.g. loud sound) and is related to a repeated movement pattern or stiffness in behavior. Since prevailing cognitive neuropsychological models cannot explain the afore-mentioned symptoms, hypotheses on sensory over- and/or hypo-arousal were developed. However, they do not have unequivocal experimental support yet (Rogers & Ozonoff, 2005b), therefore vMMN studies might give contribute to the discussion.

The study of high-functioning ASD adults found an intact vMMN for the onset of a windmill pattern, however, the sensory response (P1) had a lower amplitude in standard and deviant stimuli, longer P300 latency and, interestingly, faster target detection compared to controls (Maekawa et al., 2011). The authors emphasized that they used non-social stimuli and interpreted their findings as an impairment of low-level processing and top-down modulation with preserved bottom-up attentional mechanism in the adult ASD patients. The protocol did not control the subject's attention that was oriented toward the deviant stimulus; therefore the measured vMMN might be confounded with attentional effects.

Another study employed socially relevant stimuli, such as happy or sad faces as deviant and neutral faces as standard stimuli to elicit the vMMN in a group of adults in whom Autism Spectrum Quotient (AQ) was measured (Gayle et al., 2012). Happy faces elicited vMMN within the 150-425ms time interval at the occipito-parietal area. The vMMN showed a significant positive correlation with the AQ score: more negative vMMN amplitude was associated with lower AQ

⁴ Table 1. contains 35 experiments because different parts of two studies were used twice in our review: Horimoto et al. (2002) in Developmental disorders, and Aging and maturation, and Tales & Butler (2006) in Neurodegenerative disorders, and Aging and maturation.

A further study of adults with ASD (Cléry, Rodax, et al., 2013) used two types of change in the shape of a circle as standard and deviant stimuli. The paradigm also included a novel, always different, circle deformation and the occasional disappearance of the fixation cross, to engage the subject's attention. Adults with ASD processed deviants differently compared to the control group. While controls showed vMMN in occipito-parietal (180-240ms) and fronto-central areas (210-250ms), the patients showed a late vMMP around 460ms in occipito-parieto-temporal sites. Interestingly, the topography of the late vMMP was similar to that of the novelty P3. In addition patients with ASD also showed a significantly smaller negative component at around 160ms for the standard and deviant stimuli, and a decrease in amplitude and a shift in latency of the later positive peak (240-310ms). In contrast, the novel stimuli elicited quite similar ERP components in both groups including the novelty P3. Regarding behavioral results, there were no differences between groups in the accuracy or reaction time (RT). This study indicates dissociation between the less effective automatic detection of deviancy and augmented processing of the attention orientation in patients with ASD. The authors conclude that their findings support a higher distractibility and a lower selectivity of the attention of patients with ASD reflected as patients' intolerance to a change.

Cléry et al. (2013a) investigated 12 school age children with ASD and typically developing children with the same protocol as in their above study. The groups matched in chronological age showed a different pattern of deviance processing. While the controls had an early (280-340ms) vMMP at fronto-central electrodes and a late vMMP at 450ms at the occipito-parieto-temporal electrodes, children with ASD only showed the late vMMP that appeared significantly earlier than that of controls. The intergroup differences were restricted to the late vMMP only. The children with ASD also showed significantly prolonged early (obligatory) sensory responses and increased RT together with decreased accuracy of responses. The authors interpreted their findings reflecting an impaired sensitivity of ASD children to saliency and unselective processing of environmental changes.

An early study on visual mismatch responses compared autistic children to three control groups including typically developing children, children with dyslexia, and children with attention disorder (Kemner, Verbaten, Cuperus, Camfferman, & Van Engeland, 1994). The study used the pattern-onset of standard (rectangular shapes in the central part of a screen), deviant (180°-rotated standard) and novel (full screen letter "&") stimuli in two task conditions. The subjects fixated on the center of the screen in the passive condition, and counted the deviants in the active condition. At the time of the study, there was no agreement concerning the vMMN existence and as a visual alternative to the aMMN, a difference between the second positive and negative peaks (P2-N2) was used, and described as sensitive to the deviancy, independent of a task and without habituation (Kenemans, Verbaten, Melis, & Slangen, 1992). The authors found a significant interaction between groups (autistic vs. the controls), task (active and passive) and stimulus (standard vs. deviant or novel). The P2-N2 amplitudes of the autistic group were larger for the deviant stimuli in active condition compared to the passive condition; this result was not observed in control groups.

The afore-mentioned reports of adult patients do not yield a consistent picture of alterations of automatic visual deviance detection in the ASD. Whereas Maekawa et al. (2011) did not find any vMMN difference between patients and controls, Cléry et al. (2013) found that deviance processing is distinct in controls and the ASD patients, and in the latter group it resembles novelty detection. Gayle et al. (2012) described a relationship between autistic quotient and the deviance processing of socially relevant stimuli in healthy subjects. The only report of children with ASD (Cléry, Bonnet-Brilhault, et al., 2013) showed different characteristic of the visual deviance processing similarly to adults with ASD (Cléry, Andersson, et al., 2013) when compared to control group similarly

facial images RS was reduced in adults with ASD the RS for geometric shapes did not differ compared to controls (Ewbank et al., 2015a). RS for faces in patients with ASD was predicted from correlation analysis of RS and AQ in healthy controls (Ewbank et al., 2015b). Another prediction of a diminished RS for geometric shapes was not confirmed.

Considering the various symptoms of autism and inter-study methodological diversity, such a finding is not surprising. In spite of that, a dissociation between the sensory processing, the deviance detection, and the behavioral response was observed between controls and patients with ASD. Whereas studies in other disorders reported lower amplitudes or longer latencies of the early sensory response for standard or deviant stimuli, deviance detection was either similar (Maekawa et al., 2011) or even earlier (Cléry, Andersson, et al., 2013; Cléry, Bonnet-Brilhault, et al., 2013) in patients with ASD than that in controls. Such dissociation suggests 1) sensory impairment, and 2) disorganization of higher information processing in patients with ADS.




Fig. 5. Scalp areas of a significant difference in the deviance-related activity between patients with developmental disorders and controls. The numbers correspond to studies in Table 1: **18**-Cléry, Bonnet-Brilhault, et al. (2013), **19**-Cléry, Andersson, et al. (2013), and **23**-Wang et al. (2010). Studies without intergroup difference are omitted.

Developmental dyslexia, a reading disability, has an estimated prevalence of 7 % (Peterson & Pennington, 2012). Although the etiology of dyslexia is not well understood, the current view supports an impairment of phonological processing (Norton, Beach, & Gabrieli, 2014), as well as sensory deficits (e.g. Schulte-Körne & Bruder, 2010; Kubová et al., 2015; Stefanics, Fosker, et al., 2011). Auditory MMN has been extensively used to investigate the development of the auditory and speech systems (Näätänen et al., 1993; Cheour et al., 1998; Stefanics et al., 2007, 2009; Håden et al., 2009, 2015), as well as their disorders (for a review, see Kujala & Näätänen, 2010). The phonological deficit hypothesis is also supported by aMMN studies (for a review, see Näätänen et al., 2012a, 2014).

logographic orthographies like Chinese, and there the etiology might be in favor of the visual disturbances. The only study of visual deviance detection was conducted among Mandarin Chinese readers and was oriented to discussion of the magnocellular deficit (Wang et al., 2010). The authors used a motion sequence of a low contrast and spatial frequency grating (a magnocellular stimulus) and a high contrast and spatial frequency grating (a control condition) and compared the vMMN between children with dyslexia, calendar age matched normal readers and reading level matched controls. An auditory task with postponed response was used to capture the attention. Visuo-spatial attention was oriented to the standard/deviant stimuli. The authors found a significant difference of the vMMN amplitude (150-250ms) for the magnocellular condition between the dyslexia and both control groups at the Oz electrode. Such a difference was not present in the control condition (high contrast and spatial frequency). Together with behavioral results from a separate experiment assessing the choice reaction time to the same motion stimuli, the authors concluded that the magnocellular impairment is closely related to the orthographic processing impairment in the Chinese dyslectics. The aMMN for pitch deviance was also recorded in this study and no difference was found among the groups, which authors interpreted as a support for absence of the auditory deficit in Chinese dyslexics. The carefully prepared experiments in this study however suffered from an unclear selection of intervals for the vMMN evaluation.

Visual deviance detection was also evaluated among groups of children with idiopathic mental retardation (mean IQ=66.7), adults, and control children (Horimoto et al., 2002). The stimuli were presented for 1 s in the screen center. A green square served as deviant, and a blue square as standard stimulus. The subjects fixated at the center and focused their attention to an auditory oddball task with tones presented between visual stimuli. Whereas for adult controls the vMMN was evoked at 250ms at the Pz electrode, the vMMN in control children had a different morphology and was very difficult to identify in patients. Nevertheless, the authors described the dominant vMMN in the occipito-parietal region for control children and in the centro-parietal region for patients; however it is not clear whether the differences in the vMMN topography were statistically significant. Their experiment revealed a lower accuracy in the behavioral task and an unusually large early sensory response for the deviant stimulus in the patients. The authors concluded that children with mental retardation show an impairment of an early automatic change detection affecting the level of attention (Horimoto et al., 2002).

Summary of vMMN studies in developmental disorders

Current results support the use of the vMMN for examining sensory processing in developmental disorders. Specifically, the vMMN provides evidence of selective involvement of visual processing with intact auditory processing aMMN, in orthographic dyslexia (Wang et al. 2003). The vMMN showed differences in topography in children with ASD, while comparison between adults with ASD and control subjects failed to do so; a result possibly related to the varying use of shape or motion stimuli.

An inspiration for future vMMN studies is the aMMN study of Ceponiene et al. (2003) showing that the MMN was normal for non-social stimuli, and it was considerably reduced in amplitude for speech-sound stimuli in children with ASD. In the same vein, Kuhl et al. (2005) found that children with ASD showed a normal-size aMMN to change in non-speech sound stimuli but showed no aMMN in response to change in speech syllables. It seems that future study designs should benefit from using more socially oriented stimuli like faces and emotions, however they

Newborns are not fully equipped with a perfect sensory system and cognitive abilities. Different abilities develop at different rates, probably due to diverse experience, rate of myelination and growth. For example, basic perceptual processes mature relatively quickly showing that visual acuity is at the adult level for 8 month old babies (Norcia & Tyler, 1985), and working memory in about the early twenties (Brockmole & Logie, 2013). Once at the peak, cognitive and sensory functioning will not last forever. Although growing older is not necessarily a clinical condition, a decrement in sensory (e.g., Cheng and Lin, 2012; Lindenberger and Baltes, 1994) and cognitive (Nyberg, Bäckman, Erngrund, Olofsson, & Nilsson, 1996; Park et al., 2002) processing may accompany an otherwise healthy aging (Shimamura, Berry, Mangels, Rusting, & Jurica, 1995). Both sensory discrimination and memory processes play a crucial role in automatic deviance detection indicated by MMN (Näätänen, Paavilainen, Rinne, & Alho, 2007b). Thus, MMN may be used to assess cognitive maturation or decline in aging.

There are four studies that assessed automatic change detection in the visual modality in otherwise healthy aged participants compared to young adults. Different stimuli and tasks have been used including shape discrimination (Iijima et al., 1996; Stothart et al., 2013; Tales et al., 2002) and motion direction changes (Lorenzo-López et al., 2004). Lorenzo-López et al. (2004) used three age groups: young adults (32 ± 6 years), middle-aged (49 ± 4 years) and old (62 ± 3 years). The vMMN peak latency (measured in an oddball paradigm with 8:2 standard to deviant ratio) emerged between 145-165 ms in all age groups, with the only difference being that the older subjects showed vMMN in a smaller number of electrodes than young and middle-aged. A significant and progressive age-related reduction of the vMMN amplitude was reported between 165 and 205 ms. Tales et al. (2002) also showed a significant age-related decrease in the vMMN amplitude between young (mean age 30.5 years) and old (mean age 77, range 69-88 years) subjects, using an oddball paradigm (standard to deviant to target ratio 16:1:1; the experimental paradigm was the same as in Tales & Butler (2006) described in section 2.4 *Neurodegenerative disorders*). The group differences in amplitude occurred principally at posterior electrodes and the latency range observed was 250-400 ms. In the third study (Iijima et al., 1996) which used X and O stimuli (standard to deviant ratio 8:2), vMMN was measured from Fz, Cz and Pz electrodes, and showed similar latencies at all sites. The difference in amplitude or latency of the vMMN between young (20-29 years) and elderly (60-79 years)⁵ group was not significant but surprisingly, elderly subjects tended to elicit vMMN more frequently than did young participants. The frequency of vMMN among subjects was not statistically evaluated, unfortunately; a later vMMP emerging at around 200 ms is visible in Fig. 1 of Iijima et al. (1996) and might show age group differences which were not reported. Differences between the two age groups emerged in more frequent N2b presence for the young subjects and delayed N2b for the elderly subjects showing that attentional processing was more sensitive to age. Similarly to Iijima et al., Stothart et al. (2013) did not find vMMN differences between young (20.8 ± 3.1 years) and old (76.8 ± 6.1 years)

⁵ Iijima et al. (1996) reported different age ranges (22-29 and 62-72 years) in the full-text and in the abstract (20-29 and 60-79 years). Ranges from their abstract are used in this review.


windows according to the actual latencies (those being slightly earlier for the elderly subjects).

Even if no full-head montage of electrodes has been used, results of the above studies indicate that the main loci of the vMMN tend to emerge at O1, Oz, and O2 (Lorenzo-López et al., 2004; Stothart et al., 2013; Tales et al., 2002). These studies report a negative deflection peaking around 145-165 ms or later in averaged ERPs related to deviant detection. Different vMMN latencies may be related to the stimulus features. The control of attention may not have been an optimal in studies presenting the targets in a close vicinity of the standard/deviant stimuli (Stothart et al., 2013; Tales et al., 2002). As the ISIs used have been relatively short (1 s or less), a contribution of conscious processing in the visual working memory has probably not been decisive as it has been shown that the retention depends on time (Baddeley, 2012). This is supported also by the absence of attention related components in the ERP waveforms.

Altogether, two out of four studies suggest that there is an age-related drop of the deviance-related potential, whereas other two report an intact vMMN. Since no aging effects on vMMN were present in the Iijima et al. (1996) and Stothart et al. (2013) studies, it still is unsettled whether healthy aging is necessarily accompanied by a decrement in the change detection mechanism in the visual modality. Stothart and his coworkers propose that instead there is an age-related compensatory neural response to impoverished sensory input in aging, reflected by reduced early sensory processing (decrease in P1 amplitude), but maintained object perception (increase in N1 amplitude) and change detection processes (intact vMMN amplitude). The decrease in the vMMN amplitude (Lorenzo-López et al., 2004; Tales et al., 2002) might, however, reflect a general decrease in cognitive performance as was reported for the aMMN (Gaeta, Friedman, Ritter, & Cheng, 2001; Kisley, Davalos, Engleman, Guinther, & Davis, 2005). This is also supported by the aforementioned studies on neurodegenerative disorders (section 2.4) that do not focus on aging processes per se, while using elderly subjects as a control group for patients. Tales & Butler (2006) showed an age-related decline in the vMMN to shape discrimination between young (mean age 28, age range 20-43) and old (mean age 73 years and 2 months, age range 51-84) healthy adults in the 250-400 latency range, confirming their previous results (Tales et al., 2002). In the Tales et al. (2008 and Stothart et al. (2014) studies, only older adults (mean age 71.2 and 76, respectively) were used as the control group for the two patient groups. Similar experimental stimuli and paradigms were used as in Tales et al. (2002) and Tales & Butler (2006) studies, which makes the results of the control groups comparable to the two previous studies. Indeed, Tales et al. (2008) showed the absence of a significant vMMN response in the 250-400 and also in the 140-250 ms latency ranges, confirming their previous results. Stothart et al. (2014) showed a vMMN response in the 146-234 ms latency range, but no later vMMN and point out a rather large inter-individual variation in the vMMN response. Overall, the results of the elderly control groups of the three papers support the notion of an attenuated vMMN response in healthy aging.

In early development, opposite trends emerge than in the case of aging-related decline, but methodological diversity and possible developmental trajectories make comparing studies complicated. One of the earliest studies taking a developmental look at the vMMN (Tanaka et al.,

decrease of about 100 ms of the MMN latency (measured from Pz) emerged between 36 and 37 weeks of conceptional age, being 519 ± 17 ms for 35 weeks old babies and 418 ± 58 ms for 37 week old babies (and staying rather stable from there on). The authors suggest that it is related to the cognitive status of newborns, specifically due to myelination and qualitative and quantitative changes in neurotransmitters in the brain. The study of Tanaka et al. logically led to a study with the same visual stimuli comparing subjects from 2 to 27 years (Tomio et al., 2012). The vMMN latency was again measured from Pz and showed a gradual decrease with an increasing age, being 394 ± 58 ms at the age of 2-3 and 273 ± 32 ms at 16 years, the latter reaching the same level as the vMMN latency for adults. Based on this study, it can be concluded that for depth perception, the perceptual system seems to work as effectively at about the age of 16 years as in adulthood.



For color discrimination, the perceptual system of children seems to be comparable to that of the adults' one earlier – at about 10 years of normal development (Horimoto et al., 2002). Horimoto et al. compared normal school children (mean age 10, range 7-13 years) with adults (mean age 28.5 years), using color change as a deviant (1:4 ratio to standards). Both groups showed similar posterior vMMN peaking highest at Pz electrode in the observed latency range (180-400 ms): the vMMN amplitude was $-4.8 \pm 3.0 \mu\text{V}$ (at 250 ± 41 ms) in the adult group and only slightly (but not statistically significantly) higher in the children group. The vMMN in the children group had two peaks in the latency range of 180-400 ms. The authors conclude that the latency of the vMMN to color change reached the adult level in the 7-13 age-group and that the vMMN amplitudes were fairly similar.⁶

Cléry et al. (2012) used dynamic stimuli (deformation of a circle into an ellipse in one or another direction) in an oddball paradigm with healthy children (mean age 11.3, range 8-14 years) and adults (mean age 24, range 20-30 years). Both groups showed vMMN responses in occipito-parietal sites. The vMMN response in adults peaked at around 210 ms, while in children it seemed to have a longer duration between 150-350 ms with a broader peak, and was followed by a positive component (MMP450) around 450 ms. The same experimental paradigm was used in the Cléry et al., 2013a study, where children with ASD were compared to healthy control group (mean age 11.3, chronological age match to the ASD group). Similarly to Cléry et al. (2012), the control group showed a vMMN response peaking at around 330 ms in the occipito-parieto-temporal areas with a concurrent fronto-central positive component peaking at around 280 ms followed by a large positive wave peaking at around 450 ms. The authors suggest that this sequential change-detection process might be related to immature attentional abilities, that longer time is needed for children to process visual deviancy, and that slight variations in topographies reflect non-integrative processing of form and motion in children. Differently from Horimoto et al.(2002), the studies of Cléry et al. (2012, 2013a) indicate that the vMMN to dynamic stimuli has not yet matured by the age of 11. The latter also applies to spatial frequency changes, shown by Cleary et al. (2013) comparing healthy children (mean age 10.4, range 8-12) and adults (mean age 26.6, range 18-42) with each other. The vMMN in children had a second peak at around 250 ms which was absent in the adult group, with a concurrent frontal positivity. The results support the idea that the change detection process is still under maturation in the 8-12 age group.

Summary in aging and maturation

Studies comparing the visual deviance detection between adults and children strongly support the modularity of such a system as the vMMN matures differently for separate visual functions. This is consistent with different maturation of electrophysiological markers of visual sensory detection (Langrová, Kuba, Kremláček, Kubová, & Vít, 2006) and corresponds to the deviance detection in the

⁶ They also compared the normal children group with children that have mental retardation; those results are discussed in *Developmental disorders* section of this article.

Todd et al., 2008). Thus, the vMMN may be used as a means to assess the opposing trends of cognitive maturation or aging.

There is a strong evidence of reduced amplitude and prolonged latency of the aMMN in healthy aging (for a review, see Näätänen et al. 2012). These age changes correlated to a decline of sensory memory maintenance, interestingly, its encoding was relatively preserved (Ruzzoli et al. 2012). For this reason some studies using a short interstimulus interval (less than 2 s) showed little effect of age on the aMMN (Cheng et al. 2012). Consequently the contradictory findings between young and elderly subject observed in vMMN might be caused/explained by the short interstimulus interval (below 1 s) and therefore the age effect in the vMMN might be small and perturbable. Possible experiment addressing this issue should vary the interstimulus interval and relate it to the vMMN changes and to participants' age.

Since knowledge of the vMMN changes in healthy aging/maturation is crucial for potential search of biomarkers among various CNS disorders normative studies are of a substantial interest. As age changes of various electrophysiological components are distinct for separate visual subdomains (e.g. Kuba et al. 2012, Langrová et al. 2002), disproportional aging/maturation of the vMMN can be also expected. Research in this area will bring original findings.

As previously described in this review, no direct comparison between aMMN and vMMN have been performed and in future experiments evaluating the sensitivity/specificity of vMMN and aMMN should be balanced for saliency and attentional involvement. From the current perspective of vMMN research, there is not enough support for direct application of the vMMN in situations when healthy and pathological ageing/maturation should be separated for clinical purposes.

2.7 Miscellaneous

In this section we discuss clinically relevant vMMN studies which do not fit into the disorders grouped above. Bottari et al. (2014) compared vMMN between a group of subjects with early deafness and a hearing control group. The authors used motion stimuli in the near periphery of the visual field and allocated the subject's attention centrally by a simple detection task. The authors showed that the vMMN was evoked in the same interval (153-292ms) in deaf and hearing subjects, but with a different distribution on the scalp. While in the control group, the vMMN dominated in parietal areas, whereas in the deaf group it was most prominent centrally. The authors suggested that visual deviance is processed by supramodal brain areas that, and in case of a deafferentation of auditory input in early childhood, they might shift from visual areas toward the auditory cortex, caused by cross-modal plasticity (Kujala, Alho, & Näätänen, 2000). Interestingly, the early ERP activity around 100 ms after stimulus onset was similar across groups.

versa was examined to reveal origins of the cognitive impairment. The authors found that the temporo-parietal and occipital vMMN in the 220-330ms interval was significantly smaller in amplitude in patients compared to that of the controls. The authors suggested that the ability to automatically process facial expressions is impaired in these patients.

Moreover, Si et al. (2014), studying patients with hypertension, found that their vMMN to duration changes in stimuli randomly presented in both peripheral fields, was considerably attenuated in amplitude and delayed in peak latency compared with those of controls.

3 Discussion

As a result of the relatively narrow concept of the vMMN, the variability among experimental designs (see Fig. 7), and manifold diseases etiology thereby creating a wide spectrum of hypotheses, an individual summary for the specific clinical condition was presented at the end of each chapter. Here we summarize general views of the vMMN across all clinical conditions laying emphasis on observed effect size, correlations to other clinical parameters, and the spectrum of the experimental paradigms used. The discussion is closed by recommendations for further research.

3.1 Meta-analysis: Effect size and power in the reviewed vMMN studies

The amplitude of vMMN is usually in the range of a few microvolts, and signal to noise ratio of such a small response, as well as small sample sizes, might be an issue in some studies. To evaluate whether the reported effects have a relevant magnitude the effect size of clinical studies reviewed here was used. As the effect size was not always reported, we evaluated an equivalent Cohen's *d* effect (*eCd*) in this review. We derived the *eCd* using reported *p*, *F* or *r*-values and the group size. When a paired test was used in an original study then the *eCd* corresponds to Cohen's *d* effect measured between groups with an equal variability. The median of estimated *eCds* was 0.97 (0.84; 1.17 – the first and third quartiles respectively), showing a large effect (Cohen, 1992). The largest *eCds* per study with their respective 95% confidence intervals are plotted in Fig. 7. A considerable effect was a latency shift observed in three studies assessing the vMMN during maturation. This remarkable change in latency over maturation seems to be closely related to different projections of the vMMN on scalp in developmental and neurodegenerative disorders (Cléry, Andersson, et al., 2013; Cléry, Bonnet-Brilhault, et al., 2013; Horimoto et al., 2002), suggesting that the automatic deviance detection is likely accomplished by different neural networks.

Because of missing parameters for the *eCd* calculation, 6 studies were not included in the meta-analysis (Cléry, Andersson, et al., 2013; Cléry, Bonnet-Brilhault, et al., 2013; Iijima et al., 1995; Kremláček et al., 2011; Maekawa et al., 2011; Tales et al., 2008), therefore we evaluated 27 studies. Considering that 25 studies in Fig. 7 found a statistically significant group difference and usually there

neuroscience (Button et al., 2013). In our meta-analysis a corresponding power (PW) was calculated from the eCd , respective group size and the first type error of 0.05. We found median power of 0.77 (0.60; 0.87 - first; third quartile). Button et al. (2013) meta-analysis of neuroscience field showed that median power of 730 primary studies was around 0.21, which is much lower compared to 0.77 obtained for the vMMN studies reviewed here.

Our meta-analysis of effect and power gives a reasonable expectation that observed vMMN changes have a solid statistical background. A possible limitation of such a conclusion is that that our meta-analysis was not corrected for publication bias, and several studies reviewed here might have suffered from circular inference (Kriegeskorte, Simmons, Bellgowan, & Baker, 2009), e.g., due to a pre-selection of the intervals for statistical analysis based on visual inspection of the data. Such circular inference, also known as double dipping, might have led us to overestimate the observed effect. In future studies, it should also be taken into account that an initially reported effect is prone to extreme fluctuations, known as the Proteus phenomenon (Ioannidis & Trikalinos, 2005). In case of an original study where the effect was boosted by noise, overestimation might reduce reliability of the effect size. Considering such possible tendencies, the size of the above $eCds$ should be judged with caution, and if possible, the lower ends of the effect estimates should be used. Since many of the studies reviewed have an explorative character we have to wait for replication studies to verify the observed effects.




Fig. 7. Equivalent effect size, study power, interval of difference and type of stimulation among selected studies (see *Effect size and power of the vMMN studies*). Forest plot in the left panel shows the estimate of equivalent Cohen's effect size (black squares for significant and opened squares for not significant observations), horizontal black lines represent 95 percentile intervals. Cohen's d was computed from the f value (marked as *), from the p value (#), or from correlation coefficient (§), as it is marked in the column *Power*. The power was calculated from the *eCd* and group size. The blue bar marks the interval of Cohen's d for a medium effect size (0.5 – 0.8). The dotted line delineates the *eCd* median. Forest plot in the right panel shows intervals of the vMMN difference between groups. The *Interval* column lists intervals in milliseconds for amplitude evaluation. In case when latency was assessed, the *lat* precedes the indicated interval of intergroup means. In the right side column *Domain* lists stimulus types used in the studies. The green bars mark a task with a spatial encoding of deviancy, and the orange ones represent the deviancy encoded by a temporal stimulus aspect.

search for a trait or state marker or to address behavioral properties of the vMMN (Stelzner et al., 2014). Mostly a non-causal measure like Pearson or Spearman correlation is applied with a coefficient varying in interval from -1 to 1 based on the association of large and small values of the two analyzed parameters (Delorme, 2005).

The correlation analysis was conducted for 17 out of 33 reviewed studies and in 10 of them a significant relationship to various clinical parameters was reported (see Table 1). The vMMN latency was explored only in two developmental studies (Tanaka et al., 2001; Tomio et al., 2012), where the vMMN latency decreased with maturation. The remaining studies analyzed vMMN amplitude. Smaller visual mismatch responses were linked to more severe negative symptoms (Urban et al., 2008), and more positive vMMN was associated with better facial emotion recognition skills (Csukly et al., 2013) in schizophrenia. In a study focusing on healthy adults more negative vMMN amplitude was associated with lower Autism Spectrum Quotient (Gayle et al., 2012), however, in some subjects with a higher AQ the mismatch response was in the positive range. In general, the interpretation of such results is not straightforward.

Reports regarding the MMSE are equivocal and while Stothart et al. (2014) found that a lower MMSE corresponds to a less pronounced vMMN, such a relationship was not observed in a study of patients with dementia (Iijima et al., 1995) or hypertension (Si et al., 2014). Stothart et al. (2014) showed a positive relationship between the vMMN amplitude and the MMSE. They aggregated measures across three groups (elderly controls, patients with AD, and patients with MCI). For this particular case predominantly patients with the MCI had the positive amplitudes and therefore a compensatory mechanism, analogous to changes accompanying aging in vision (Stothart et al., 2013), might explain the vMMP. Furthermore, depression severity, assessed by the Hamilton Rating Scale of Depression, was found to be associated to the vMMN in the study by Chang et al. (2011) but not in the study by Qui et al. (2011).

Regarding studies on substance abuse, while a longer methamphetamine abuse (Hosák et al., 2008) or a higher antipsychotic dose (Urban et al., 2008) corresponded to a lower vMMN amplitude, the level of alcohol consumption did not (Kenemans et al., 2010). However, these results should not be taken as a sign of inconsistency, because an acute intoxication was considered in case of the alcohol study and a long term effect of use/abuse was followed in the other two studies. Moreover the antipsychotics have diverse pharmacological effects on brain functions from sympathomimetic drugs like methamphetamine or alcohol.

3.3 Stimulus features used for vMMN elicitation

Various visual features (color, location, motion, shape, or facial expression) were used to establish regularity through stimulus repetition in the studies reviewed. In most studies an unexpected change in shape was used (Fig. 7) to violate the established regularity and elicit a mismatch response. The stimulus features used can be divided into two categories based on their presumed processing pathways. Specifically, we suggest that the ventral (what) and dorsal (where) visual stream

that there might be an earlier deviance processing deficit for temporal stimuli compared to stimuli which violates a spatial regularity. This is not surprising, considering that the spatial stimuli might also combine several features (location, color, shape, or emotion) which might be attributes of complex objects that elicit mismatch responses at higher levels of the processing hierarchy with a corresponding impairment at longer latencies.

3.4 Specificity to predictive processing

A current hypothesis links the vMMN to predictive coding theories and suggests that the vMMN response represents a bottom-up perceptual prediction error signal as well as its ensuing top-down update process. Since the majority of the reviewed studies reported some kind of alteration of the deviance processing in various disorders, is it safe to say that an impairment of automatic predictive perceptual processes seem to be present in the very heterogeneous disorders reviewed here?

It was suggested that two underlying processes contribute to the vMMN, 1) response attenuation to repeated stimuli, and 2) the prediction error elicited by the unexpected event (Czigler, Sulykos, & Kecskés-Kovács, 2014). For assessment of the prediction error a subtraction of response to equiprobable control stimulus from the deviant response was proposed (Kimura, 2012). Among the reviewed studies, Chang et al. (2011) used this approach to compare patients with depression to a control group. Interestingly, the only difference they found between the groups was for the deviant minus the equiprobable stimuli. Such observation suggests that processes underlying stimulus repetition effects and prediction error generation might be dissociated and separately impaired in depression. Concerning other effects listed in this review, it is not possible to exclude that impairment of response attenuation might contribute to deficient vMMN in patients.

Impaired predictive processing might also manifest as a general sensory impairment. Such impairment would be present in a sensory evoked potential, sometimes called an obligatory response, recorded in a so-called one stimulus paradigm, e.g. in response to a repeating standard stimulus. Out of 33 reviewed studies 16 evaluated obligatory sensory components beside the vMMN. The sensory processing was disturbed in 13 of them (Bottari et al., 2014; Cleary et al., 2013; Cléry et al., 2012; Cléry, Bonnet-Brilhault, et al., 2013; Cléry, Roux, et al., 2013; Fisher et al., 2010; Iijima et al., 1996; Kremláček et al., 2011; Lorenzo-López et al., 2004; Maekawa et al., 2013, 2011; Stothart et al., 2014, 2013), whereas deficits only in components related to processing of deviants was observed in two studies only (Tales & Butler, 2006; Urban et al., 2008). The almost ubiquitous impairment of sensory stimulus processing in reviewed studies suggests that vMMN impairment is not necessarily the only candidate marker for impairments of perceptual processes.

3.5 Limitations for clinical use

It is often stated that using the vMMN does not require the subject's behavioral response for evaluating higher functions than sensory detection, e.g. sensory discrimination, as this ERP component is usually obtained in a passive paradigm. This, however, does not mean that patient's

available in the auditory domain, such as testing predictive processing of information which can only be conveyed in the visual modality. The advantage of using vMMN to investigate specific hypotheses with appropriate control conditions for possible confounds arising from attentional modulation or specific sensory adaptation, comes at a cost of complex and time demanding protocols, which might impose limits on patient studies.

Auditory MMN studies suggest that some subjects, despite good stimulus discrimination and behavioral performance, do not show a reliable aMMN response. Bishop & Hardiman (2010) found that 82% of their healthy subjects (n=17) showed a statistically reliable MMN component to pitch deviants. Thus, the interpretation of absence of a reliable vMMN response in a single subject might not always be straightforward also in vMMN studies, assuming that a reliable vMMN response is not observable in a subset of healthy subjects with otherwise good behavioral performance in stimulus discrimination.

4 Conclusions and directions for further research

We reviewed 33 studies where visual MMN has been used to investigate impairments of automatic perceptual processes in neuropsychiatric conditions. Our survey suggests that vMMN - similarly to auditory MMN - is a potentially useful tool for research in several clinical populations.

The deficits in vMMN in different disorders are not restricted to a certain time interval or scalp location. This is presumably due to the fact that different studies used different experimental stimulus sets, and brain areas generating the mismatch response to different stimuli might vary across studies. Furthermore, it seems plausible that there are multiple distinct physiological mechanisms that serve a common function: computing prediction errors and predictions. It is conceivable that these mechanisms might be altered differentially in specific disorders, or in different subgroups of the same disease, factors that might have contributed to the observed variability of vMMN findings across studies. We must point out that an optimal vMMN paradigm with control for all aspects of balanced design (see Technical notes at the end of the paper) has not been used frequently, and reproducibility, specificity, and sensitivity were not always estimated or addressed. Clinical studies using vMMN would benefit from standardized protocols as they allow for a broader replicability/reproducibility evaluation as well as multicenter studies.

Although there are now increasingly stratified treatments for psychiatric and neurological conditions, in most cases no clear indicators which treatment would be the best for a given individual patient are available. ERPs, including visual and auditory MMN, hold promise to serve as biomarkers for individually tailored treatments (Luck et al., 2011). Furthermore, they might be useful to evaluate new compounds and as diagnostic tools. However, the use of ERPs in guiding diagnostic or treatment decisions requires further basic research. There are open questions such as, which are the optimal paradigms, stimuli, and ERP components for each specific purpose. More advanced signal processing methods should be used to allow an optimal evaluation of an individual patient's ERP data and how

subgroups. For example, a strong genetic component underlying depression might be observed as a distinct pattern of findings in ERPs compared to depression that has its roots in environmental stress. Unraveling these effects could allow a more accurate diagnosis and treatment guided by low-cost and non-invasive ERP recordings.

Future research would benefit from large-scale research projects comparing brain responses in different psychiatric and neurological disorders using similar experimental protocols. Also comparisons of brain responses in different stimulus conditions in the same group of participants are of interest. Future research should also test the predictive value of the vMMN for treatment outcome.

In clinical research both examination time and the patient's attention can be limited, therefore it is important to determine the most appropriate or potentially successful application of MMN. Regarding the optimal choice of a deviant feature to elicit the MMN in an oddball paradigm, the experimenter is not necessarily limited to using only one deviant stimulus type, as multi-feature paradigms that allow obtaining vMMN for several visual attributes in a short time are possible (Kreegipuu et al., 2013; Qian et al., 2014), some of which might prove more relevant in a particular population than the others. Another important choice is in which sensory modality should be the MMN investigated, e.g., visual and/or auditory. Although this review cannot give a clear answer as today there are only a few studies where both modalities have been tested, there are some situations where the disease directly influences the choice. While in schizophrenia the mean *eCD* vMMN was 0.86, meta-analysis of 32 aMMN studies showed comparably large Cohen's *d* effect of 0.99 (Umbricht & Krljes, 2005) and the impairment of the sensory processing seems to be general, only the vMMN and no aMMN was reduced in study of orthographic dyslexia (Wang et al. 2010). Furthermore, using aMMN is also not feasible in cases of severe auditory impairment, where, nevertheless, vMMN might be fruitful. For example, Bottari et al. (2014) used vMMN to assess plastic reorganization of the auditory cortex in early deafness and showed that cross-modally recruited auditory cortex participates in predictive processing of visual information..

Another field where vMMN might be potentially useful in clinical research is neuro-ophthalmology. Development of intraocular implants (Ahuja et al., 2011, Shepherd et. al., 2013) to artificially augment vision in blind people provides a unique opportunity for studying mechanisms of neural plasticity in the visual cortex. Furthermore, understanding plasticity could have major implications in the regenerative medicine of retinal diseases, such as age-related macular degeneration, retinitis pigmentosa, or other clinical conditions, e.g., long term visual deprivation, amblyopia, or stroke that involves cortical compensatory plasticity (Rosa et al., 2013). Since vMMN is relying on mechanisms of short-term plasticity, it might represent an important contribution to research in the above conditions.

VMMN opens new space for exploring automatic predictive processing of information with a substantial advantage over the aMMN where complex content have to be presented abruptly to elicit a brain response measurable with ERPs. A potential advantage of visual MMN over its auditory

automatic perceptual component of social cognition (Astikainen et al., 2009; Stefanics et al., 2012) in disorders where it is impaired, such as in schizophrenia (Kohler et al., 2003; Morris et al., 2009; Komlosi et al., 2012; Csukly et al., 2014). In general, VMMN might be a useful tool for studying deficits in predictive processing in cognitive domains where using visual rather than auditory stimuli is more adequate. Also, vMMN might be very useful when pathology is shown to potentially affect visual processing areas/or related neurotransmitters.

On the downside, using an appropriate visual distractor task to control attentional effects might be demanding for clinical populations. Controlling visual attention in MMN experiments is important to keep responses to unpredicted stimuli interpretable as prediction errors, free of potentially confounding effects of attention and task demands. Therefore using less demanding, but more entertaining, game-like tasks (Sulykos & Czigler, 2011; Kecskés-Kovács et al., 2013) is important, and may counteract this possible drawback.

Visual MMN research, both basic and clinical, might substantially benefit from adopting the predictive coding perspective. Predictive coding provides a framework for theoretical models of perceptual inference, cognition, learning and decision making (den Ouden et al., 2012; Friston, 2005, 2010), and it might provide visual MMN research with a principled probabilistic approach to test neurophysiologically grounded hypotheses in clinical conditions (Stephan et al., 2006, 2009, 2016a,b; Corlett et al., 2007, 2011; Adams et al., 2013; Friston et al., 2014). Thus, it might help to understand better disease mechanisms and dissect heterogeneous clinical groups into well-defined subgroups to guide diagnosis, predict response to treatments or conversion to psychosis, or track disease progression. We believe that visual MMN, along to its auditory counterpart, might contribute to this process.

Although the potential for vMMN to serve as a specific clinical marker remains to be fully determined, this field is in its infancy, and though there is much debate regarding theoretical and methodological approach, the outcome of the studies described here highlight potentially clinically-relevant applications of vMMN and call for further research investment, addressing current limitations and proposing standardized protocols.

5 Technical notes

5.1 MMN as differential activity

The determination of an original condition (standard / deviant activation) from difference wave (MMN = Deviant - Standard) is an ill-posed problem, there being endless possibilities to generate the same differential curve.




Fig. 8. Different scenarios of deviant and standard responses that result in similar difference waves. Besides the situation (a), where the rare deviant stimulus elicits a larger negative potential than the repeated standard, there are also plausible scenarios, where (b) the standard stimulus elicits a more positive potential, which might correspond to potentiation in response to repeated stimuli described for example in migraine (de Tommaso et al., 2014), or where (c) responses with similar magnitude but with opposite polarity are evoked. For this reason it is advisable to include data both from the standard and deviant conditions in the statistical analysis and not only test the differential waveforms. Including stimulus types as a factor might reveal differences which would otherwise remain undetected.

Another important point to consider is using a proper control condition (for a review, see Stefanics et al., 2014). Studies using an oddball paradigm often record a so-called “reverse-block” where probabilities of the stimuli that served as frequent standard and rare deviant are reversed, which allows comparing responses to physically identical stimuli that were presented with different probability. An elegant method to control for effects both for physical features and probability is to use an equal probability control condition (Schröger & Wolff, 1996; Schröger, 1997; Jacobsen & Schröger, 2001; Ruhnau et al., 2012). For each disorder in the following sections we show scalp plots of group differences indicating altered deviance processing in a variety of disorders. However, as the above examples show these differences do not necessarily mean that patients have a smaller negative response to the deviant stimuli. Furthermore, the interpretation of differences in MMN between groups should be done cautiously as generating structures might differ between the groups which might result in different scalp potential topographies for patient and control groups.

5.2 Limitation of the reported vMMN scalp distributions

In this review we report electrode positions or scalp areas where differences between study groups in mismatch activity have been found. It should be emphasized that these locations do not necessarily correspond to cortical sources of the MMN potential. The main reason is that distribution of scalp potentials depends on the reference electrode, and an electrically active source, which can project

It is worth to note that locations of the reported group differences could be affected by limited number of recording electrodes or by predetermined selection of the electrodes for statistical evaluation of the vMMN.

6 Acknowledgement

JK was supported by P37/07 (PRVOUK) program. PA was supported by the Academy of Finland (project no. 140126). RN, KK and NP were supported by Institutional Research Grant IUT 02-13 to Jüri Allik from Estonian Ministry of Education and Research and by funding from the "Norwegian financial mechanism 2009-2014" through the Norwegian-Estonian Research Cooperation Programme (project EMP180). AT was supported by BRACE-Alzheimer's Research [Registered Charity No. 297965] and Swansea University.

7 References

- Adams, R.A., Stephan, K.E., Brown, H.R., Frith, C.D., & Friston, K.J. (2013). The computational anatomy of psychosis. *Front. Psychiatry* 4:47. doi: 10.3389/fpsyt.2013.00047
- Ahuja, A. K., Dorn, J. D., Caspi, A., McMahon, M. J., Dagnelie, G., Dacruz, L., Greenberg, R. J., & Argus II Study Group. (2011). Blind subjects implanted with the Argus II retinal prosthesis are able to improve performance in a spatial-motor task. *British Journal of Ophthalmology*, 95, 539–543.
- Amenedo, E., Pazo-Alvarez, P., & Cadaveira, F. (2007). Vertical asymmetries in pre-attentive detection of changes in motion direction. *Int. J. Psychophysiol.* 64, 184-189.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Washington, DC: American Psychiatric Association.
- Andersson, S., Barder, H. E., Hellvin, T., Løvdahl, H., & Malt, U. F. (2008). Neuropsychological and electrophysiological indices of neurocognitive dysfunction in bipolar II disorder. *Bipolar Disorders*, 10, 888–899. doi:10.1111/j.1399-5618.2008.00638.x
- Andrade, G.N., Butler, J.S., Mercier, M.R., Molholm, S., & Foxe, J.J. (2015). Spatio-temporal dynamics of adaptation in the human visual system: a high-density electrical mapping study. *Eur. J. Neurosci.* 41, 925-939.
- Astikainen, P., & Hietanen, J.K. (2009). Event-related potentials to task-irrelevant changes in facial expressions. *Behav. Brain Funct.* 5:30. doi:10.1186/1744-9081-5-30

- Astikainen, P., Liistaring, E., & Ruusuvirta, T. (2008). Visual mismatch negativity for changes in orientation – a sensory memory-dependent response. *Eur. J. Neurosci.* 21, 2319-2324.
- Astikainen, P., Ruusuvirta, T., Wikgen, J., & Korjonen, T. (2004). The human brain processes visual changes that are not cued by attended auditory stimulation. *Neurosci. Lett.* 368, 231-234.
- Baddeley, A. (2012). Working memory: theories, models, and controversies. *Annual Review of Psychology*, 63, 1–29. doi:10.1146/annurev-psych-120710-100422
- Baldeweg, T. (2006). Repetition effects to sounds: evidence for predictive coding in the auditory system. *Trends Cogn. Sci.* 10, 93-94.
- Baldeweg, T. (2007). ERP repetition effects and mismatch negativity generation - A predictive coding perspective. *J. Psychophysiol.* 21, 204-213. 10.1027/0269-8803.21.34.204.
- Barch, D.M., Ceaser, A. (2012). Cognition in schizophrenia: core psychological and neural mechanisms. *Trends Cogn. Sci.*, 16, 27-34.
- Barlow, H.B. (1990). A theory about the functional role and synaptic mechanisms of visual after-effects. In *Vision: Coding and Efficiency*, C. Blakemore, ed. (New York: Cambridge University Press), pp. 363–375.
- Barr, A.M., Panenka, W.J., MacEwan, G.W., Thornton, A.E., Lang, D. J., Honer, W.G., & Lecomte, T. (2006). The need for speed: an update on methamphetamine addiction. *Journal of Psychiatry and Neuroscience*, 31, 301–313.
- Bartholow, B. D., Pearson, M., Sher, K. J., Wieman, L. C., Fabiani, M., & Gratton, G. (2003). Effects of alcohol consumption and alcohol susceptibility on cognition: a psychophysiological examination. *Biological Psychology*, 64, 167-190. doi:10.1016/S0301-0511(03)00108-X
- Beck, A. T. (1967). *Depression: Clinical, Experimental and Theoretical Aspects*. New York: Harper & Row.
- Beck, A. T. (1976). *Cognitive Therapy and the Emotional Disorders*. New York: Int. Univ. Press.
- Beck, A. T. (2008). The evolution of the cognitive model of depression and its neurobiological correlates. *The American Journal of Psychiatry*, 165, 969–977. doi:10.1176/appi.ajp.2008.08050721
- Benucci, A., Saleem, A. B., & Carandini, M. (2013). Adaptation maintains population homeostasis in primary visual cortex. *Nature Neuroscience*, 16, 724–729.
- Bijl, S., Bruin, E. A., Kenemans, J. L., Verbaten, M. N., & Böcker, K. B. (2005). Effects of Chronic Alcohol Consumption in a Visual Attention Task and an Auditory Oddball Task: An Event-Related Potential Study. *Alcoholism: Clinical and Experimental Research*, 29, 2029-2038.

- Biau, V., N. Van Atteveldt, M. Erkebus, R. Goebel, & Blömert, L. (2009). Reduced neural integration of letters and speech sounds links phonological and reading deficits in adult dyslexia. *Current Biology* 19, 503–508, 2009.
- Blumberg, S. J., Ph, D., & Bramlett, M. D. (2013). Changes in Prevalence of Parent-reported Autism Spectrum Disorder in School-aged U. S. Children : 2007 to 2011 – 2012. *National Health Statistics Reports*, 20, 1–12.
- Boehnke, S.E., Berg, D.J., Marino, R.A., Baldi, P.F., Itti, L., & Munoz, D.P. (2011). Visual adaptation and novelty responses in the superior colliculus. *Eur J Neurosci*. 34, 766-779. doi: 10.1111/j.1460-9568.2011.07805.x.
- Bottari, D., Heimler, B., Caclin, A., Dalmolin, A., Giard, M.-H., & Pavani, F. (2014). Visual change detection recruits auditory cortices in early deafness. *NeuroImage*, 94, 172–84. doi:10.1016/j.neuroimage.2014.02.031
- Brockmole, J. R., & Logie, R. H. (2013). Age-related change in visual working memory: A study of 55,753 participants aged 8-75. *Front. Psychol.* 4:12. doi:10.3389/fpsyg.2013.00012
- Bromet, E., Andrade, L. H., Hwang, I., Sampson, N. A., Alonso, J., de Girolamo, G., de Graaf, R., Demyttenaere, K., Hu, C., Iwata, N., Karam, A.N., Kaur, J., Kostyuchenko, S., Lépine, J.P., Levinson, D., Matschinger, H., Mora, M.E., Browne, M.O., Posada-Villa, J., Viana, M.C., Williams, D.R., & Kessler, R. (2011). Cross-national epidemiology of DSM-IV major depressive episode. *BMC Medicine*, 9, 90. doi:10.1186/1741-7015-9-90
- Brønneck, K. S., Nordby, H., Larsen, J. P., & Aarsland, D. (2010). Disturbance of automatic auditory change detection in dementia associated with Parkinson’s disease: A mismatch negativity study. *Neurobiology of Aging*, 31, 104–13. doi:10.1016/j.neurobiolaging.2008.02.021
- Buchanan, R.W., Davis, M., Goff, D., Green, M.F., Keefe, R.S.E, Leon, A.C., Nuechterlein, K.H., Laughren, T., Levin, R., Stover, E., Fenton, W., & Marder, S.R. (2005). A Summary of the FDA-NIMH-MATRICES Workshop on Clinical Trial Design for Neurocognitive Drugs for Schizophrenia. *Schizophr Bull.* 31, 5-19.
- Buckner, R.L., Koutstaal, W., Schacter, D.L., & Rosen, B.R. (2000). Functional MRI evidence for a role of frontal and inferior temporal cortex in amodal components of priming. *Brain*. 123, 620-640.
- Butler, P. D., Silverstein, S. M., & Dakin, S. C. (2008). Visual perception and its impairment in schizophrenia. *Biological Psychiatry*, 64, 40–47. doi:10.1016/j.biopsych.2008.03.023
- Button, K. S., Ioannidis, J. P. a, Mokrysz, C., Nosek, B. a, Flint, J., Robinson, E. S. J., & Munafò, M. R. (2013). Power failure: why small sample size undermines the reliability of neuroscience. *Nat. Rev. Neurosci.* 14, 365–376. doi:10.1038/nrn3475

- Ceponiene, R., Lepistö, T., Shestakova, a, Vanhala, R., Alku, P., Näätänen, R., & Yaguchi, K. (2003). Speech-sound-selective auditory impairment in children with autism: they can perceive but do not attend. *Proceedings of the National Academy of Sciences of the United States of America*, 100, 5567–5572. doi:10.1073/pnas.0835631100
- Chang, L., Friedman, J., Ernst, T., Zhong, K., Tsopeles, N. D., & Davis, K. (2007). Brain metabolite abnormalities in the white matter of elderly schizophrenic subjects: implication for glial dysfunction. *Biological Psychiatry*, 62, 1396–404. doi:10.1016/j.biopsych.2007.05.025
- Chang, Y., Xu, J., Shi, N., Pang, X., Zhang, B., & Cai, Z. (2011). Dysfunction of preattentive visual information processing among patients with major depressive disorder. *Biological Psychiatry*, 69, 742–7. doi:10.1016/j.biopsych.2010.12.024
- Chang, Y., Xu, J., Shi, N., Zhang, B., & Zhao, L. (2010). Dysfunction of processing task-irrelevant emotional faces in major depressive disorder patients revealed by expression-related visual MMN. *Neuroscience Letters*, 472, 33–37. doi:10.1016/j.neulet.2010.01.050
- Cheng, C. H., & Lin, Y. Y. (2012). The effects of aging on lifetime of auditory sensory memory in humans. *Biological Psychology*, 89, 306–312. doi:10.1016/j.biopsycho.2011.11.003
- Cheng, C. H., Hsu, W. Y., & Lin, Y. Y. (2013). Effects of physiological aging on mismatch negativity: A meta-analysis. *International Journal of Psychophysiology*, 90, 165–171. doi:10.1016/j.ijpsycho.2013.06.026
- Cheour, M., Ceponiene, R., Lehtokoski, A., Luuk, A., Allik, J., Alho, K., & Näätänen, R. (1998). Development of language-specific phoneme representations in the infant brain. *Nat Neurosci.* 1, 351-353.
- Clark, A. (2013). Whatever next? Predictive brains, situated agents, and the future of cognitive science. *Behav. Brain Sci.* 36, 181-204.
- Cleary, K. M., Donkers, F. C. L., Evans, A. M., & Belger, A. (2013). Investigating developmental changes in sensory processing: visual mismatch response in healthy children. *Front. Hum. Neurosci.* 7:922. doi:10.3389/fnhum.2013.00922
- Cléry, H., Andersson, F., Bonnet-Brilhault, F., Philippe, A., Wicker, B., & Gomot, M. (2013). FMRI investigation of visual change detection in adults with autism. *NeuroImage: Clinical*, 2, 303–312. doi:10.1016/j.nicl.2013.01.010
- Cléry, H., Bonnet-Brilhault, F., Lenoir, P., Barthelemy, C., Bruneau, N., & Gomot, M. (2013). Atypical visual change processing in children with autism: An electrophysiological Study. *Psychophysiol.* 50, 240–52. doi:10.1111/psyp.12006

- Ciery, H., Roux, S., Houdy-Durand, E., Bonnet-Brihault, F., Bruneau, N., & Gomot, M. (2013). Electrophysiological evidence of atypical visual change detection in adults with autism. *Front. Hum. Neurosci.* 7:62. doi:10.3389/fnhum.2013.00062
- Cohen, J. (1992). A power primer. *Psychological Bulletin*, 112, 155–159.
- Corlett, P.R., Honey, G.D., Krystal, J.H., & Fletcher, P.C. (2011). Glutamatergic model psychoses: prediction error, learning, and inference. *Neuropsychopharmacology*. 36, 294-315.
- Corlett, P.R., Murray, G.K., Honey, G.D., Aitken, M.R., Shanks, D.R., Robbins, T.W., Bullmore, E.T., Dickinson, A., & Fletcher, P.C. (2007). Disrupted prediction-error signal in psychosis: evidence for an associative account of delusions. *Brain*. 130, 2387-400.
- Coyle, J.T. (2012). NMDA Receptor and Schizophrenia: A Brief History. *Schizophr Bull.*, 38, 920-926.
- Csukly, G., Stefanics, G., Komlósi, S., Czigler, I., & Czobor, P. (2013). Emotion-Related Visual Mismatch Responses in Schizophrenia: Impairments and Correlations with Emotion Recognition. *PLoS ONE*, 8(10), e75444. doi:10.1371/journal.pone.0075444
- Csukly, G., Stefanics, G., Komlósi, S., Czigler, I., Bitter, I., & Czobor, P. (2014). Event-related theta synchronization predicts deficit in facial affect recognition in schizophrenia. *Journal of Abnormal Psychology*, 123, 178-189.
- Czigler, I. (2007). Visual Mismatch Negativity. *J. Psychophysiol.* 21, 224–230. doi:10.1027/0269-8803.21.34.224
- Czigler, I., & Pató, L. (2009). Unnoticed regularity violation elicits change-related brain activity. *Biol. Psych.* 80, 339-347.
- Czigler, I., & Sulykos, I. (2010). Visual mismatch negativity to irrelevant changes is sensitive to task-relevant changes. *Neuropsychol.* 48, 1277-1282.
- Czigler, I., Balázs, L., & Pató, L. (2004). Visual change detection: event-related potentials are dependent on stimulus location in humans. *Neurosci. Lett.* 364, 149-153.
- Czigler, I., Balazs, L., & Winkler, I. (2002). Memory-based detection of task-irrelevant visual changes. *Psychophysiology*, 39, 869-873.
- Czigler, I., Balázs, L., & Winkler, I. (2002). Memory-based detection of task-irrelevant visual changes. *Psychophysiol.* 39, 869-873.
- de Gardelle, V., Waszczuk M, Egnér T, Summerfield C. (2013). Concurrent repetition enhancement and suppression responses in extrastriate visual cortex. *Cereb. Cortex*, vol. 23, pp. 2235-2244.

- Delorme, A. (2005). Statistical methods. *Encyclopedia of Medical Device and Instrumentation*, 6, 1–23. doi:10.1002/0471732877.emd318
- Desimone, R. (1996). Neural mechanisms for visual memory and their role in attention. *Proc. Natl. Acad. Sci. USA*, 93, 13494-13499.
- Dorph-Petersen, K.-A., Caric, D., Saghafi, R., Zhang, W., Sampson, A. R., & Lewis, D. A. (2009). Volume and neuron number of the lateral geniculate nucleus in schizophrenia and mood disorders. *Acta Neuropathologica*, 117, 369–84. doi:10.1007/s00401-008-0410-2
- Dudai, Y. (2002). *Memory from A to Z*. Oxford: Oxford University Press.
- Dulude, L., Labelle, A., & Knott, V. J. (2010). Acute nicotine alteration of sensory memory impairment in smokers with schizophrenia. *Journal of clinical psychopharmacology*, 30, 541-548.
- Egner, T., Monti, J. M., & Summerfield, C. (2010). Expectation and surprise determine neural population responses in the ventral visual stream. *Journal of Neuroscience*, 30, 16601-16608.
- Ewbank, M. P., Rhodes, G., von dem Hagen, E. a H., Powell, T. E., Bright, N., Stoyanova, R. S., & Calder, A. J. (2015b). Repetition Suppression in Ventral Visual Cortex Is Diminished as a Function of Increasing Autistic Traits. *Cerebral Cortex*. 25, 3381-3393. doi:10.1093/cercor/bhu149
- Ewbank, M., Pell, P., Powell, T., von em Hagen, E., Baron-Cohen, S., & Calder, A. (2015a). Reduced repetition suppression to faces in the fusiform face area of adults with autism spectrum conditions. *Journal of Vision*, 15, 1210.
- Farkas, K., Stefanics, G., Marosi, C., & Csukly, G. (2015). Elementary sensory deficits in schizophrenia indexed by impaired visual mismatch negativity. *Schizophrenia Research*, 166, 164-170. doi:10.1016/j.schres.2015.05.011
- Fillmore, M. T., & Van Selst, M. (2002). Constraints on information processing under alcohol in the context of response execution and response suppression. *Experimental and Clinical Psychopharmacology*, 10, 417-424. <http://dx.doi.org/10.1037/1064-1297.10.4.417>
- Fillmore, M. T., & Vogel-Sprott, M. (1999). An alcohol model of impaired inhibitory control and its treatment in humans. *Experimental and Clinical Psychopharmacology*, 7, 49-55. <http://dx.doi.org/10.1037/1064-1297.7.1.49>
- Fisher, D. J., Scott, T. L., Shah, D. K., Prise, S., Thompson, M., & Knott, V. J. (2010). Light up and see: enhancement of the visual mismatch negativity (vMMN) by nicotine. *Brain Research*, 1313, 162–71. doi:10.1016/j.brainres.2009.12.002

- Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). Mini-mental state. *Journal of Psychiatric Research*. doi:10.1016/0022-3956(75)90026-6
- Friston, K. (2005). A theory of cortical responses. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* 360, 815–836.
- Friston, K. (2010). The free-energy principle: a unified brain theory. *Nat. Rev. Neurosci.* 11, 127-138.
- Friston, K.J., Stephan, K.E., Montague, R., & Dolan, R.J. (2014). Computational psychiatry: the brain as a phantastic organ. *Lancet Psychiatry*. 1, 148-158. doi: 10.1016/S2215-0366(14)70275-5.
- Fujimura, T., & Okanoya, K. (2013). Event-related potentials elicited by pre-attentive emotional changes in temporal context. *PLoS One*. 8(5):e63703. doi: 10.1371/journal.pone.0063703.
- Fusar-Poli, P., & Meyer-Lindenberg, A. (2013). Striatal presynaptic dopamine in schizophrenia, Part II: meta-analysis of [18F/11C]-DOPA PET studies. *Schizophrenia bulletin*, 39, 33-42. doi: 10.1093/schbul/sbr180
- Gaeta, H., Friedman, D., Ritter, W., & Cheng, J. (2001). An event-related potential evaluation of involuntary attentional shifts in young and older adults. *Psychology and Aging*, 16, 55–68. doi:10.1037/0882-7974.16.1.55
- Garagnani, M., & Pulvermüller, F. (2011). From sounds to words: a neurocomputational model of adaptation, inhibition and memory processes in auditory change detection. *Neuroimage* 54, 170-181. doi: 10.1016/j.neuroimage.2010.08.031.
- Garrido, M.I., Friston, K.J., Kiebel, S.J., Stephan, K.E., Baldeweg, T., & Kilner, J.M. (2008). The functional anatomy of the MMN: A DCM study of the roving paradigm. *Neuroimage* 42, 936-944.
- Garrido, M.I., Kilner, J.M., Kiebel, S.J., Stephan, K.E., & Friston, K.J. (2007). Dynamic causal modelling of evoked potentials: a reproducibility study. *NeuroImage* 36, 571–580.
- Garrido, M.I., Kilner, J.M., Stephan, K.E., & Friston, K.J. (2009). The mismatch negativity: a review of underlying mechanisms. *Clin. Neurophysiol.* 120, 453-463. doi: 10.1016/j.clinph.2008.11.029.
- Garrido, M.I., Sahani, M., & Dolan, R.J. (2013). Outlier responses reflect sensitivity to statistical structure in the human brain. *PLoS. Comput. Biol.* 9, e1002999.
- Gayle, L. C., Gal, D. E., & Kieffaber, P. D. (2012). Measuring affective reactivity in individuals with autism spectrum personality traits using the visual mismatch negativity event-related brain potential. *Front. Hum. Neurosci.* 6, 334. doi:10.3389/fnhum.2012.00334
- Green, M. F., Kern, R. S., & Heaton, R. K. (2004). Longitudinal studies of cognition and functional outcome in schizophrenia: implications for MATRICS. *Schizophrenia Research*, 72, 41–51. doi:10.1016/j.schres.2004.09.009

- Grotheer, M., & Kovács, G. (2014). Repetition probability effects depend on prior experiences. *J. Neurosci.* 34, 6640-6646. doi: 10.1523/JNEUROSCI.5326-13.2014.
- Grotheer, M., & Kovács, G. (2015). The relationship between stimulus repetitions and fulfilled expectations. *Neuropsychologia.* 67, 175-182. doi: 10.1016/j.neuropsychologia.2014.12.017.
- Háden, G., Németh, R., Török, M., & Winkler, I. (2015). Predictive processing of pitch trends in newborn infants. *Brain Res.* 1626, 14-20. doi: 10.1016/j.brainres.2015.02.048.
- Háden, G., Stefanics, G., Huotilainen, M., Balázs, L., Sziller, I., Beke, A., Fellman, V., & Winkler, I. (2009). Timbre-independent extraction of pitch in newborn infants. *Psychophysiology*, 46, 69-74.
- Haenschel, C., Vernon, D.J., Dwivedi, P., Gruzelier, J.H., & Baldeweg, T. (2005). Event-related brain potential correlates of human auditory sensory memory-trace formation. *J. Neurosci.* 25, 10494-10501.
- Haijiang, Q., Saunders, J. A., Stone, R. W., & Backus, B. T. (2006). Demonstration of cue recruitment: Change in visual appearance by means of Pavlovian conditioning. *Proceedings of the National Academy of Sciences of the United States of America*, 103, 483-488.
- Hart, C. L., Ward, A. S., Haney, M., Foltin, R. W., & Fischman, M. W. (2001). Methamphetamine self-administration by humans. *Psychopharmacology*, 157, 75-81.
- He, J., Hu, Y., Pakarinen, S., Li, B., & Zhou, Z. (2014). Different effects of alcohol on automatic detection of colour, location and time change: A mismatch negativity study. *J. Psychopharmacol.* 28, 1109-1114. doi:10.1177/0269881114548294
- Heekeren, K., Daumann, J., Neukirch, A., Stock, C., Kawohl, W., Norra, C., Waberski, T.D., & Gouzoulis-Mayfrank, E. (2008). Mismatch negativity generation in the human 5HT2A agonist and NMDA antagonist model of psychosis. *Psychopharmacology (Berl)*. 199, 77-88.
- Heinrichs, R.W., & Zakzanis K.K. (1998). Neurocognitive deficit in schizophrenia: a quantitative review of the evidence. *Neuropsychology*, 12, 426-445.
- Heng, K., Hargarten, S., Layde, P., Craven, A., & Zhu, S. (2006). Moderate alcohol intake and motor vehicle crashes: The conflict between health advantage and at-risk use. *Alcohol and Alcoholism* 41, 451-454. DOI: <http://dx.doi.org/10.1093/alcalc/agh258>
- Henson, R.N., & Rugg, M.D. (2003). Neural response suppression, haemodynamic repetition effects, and behavioural priming. *Neuropsychologia.* 41. 263-270.
- Henson, R.N., Mouchlianitis, E., Matthews, W.J., & Kouider, S. (2008). Electrophysiological correlates of masked face priming. *Neuroimage*, 40, 884-895.

- Heslenfeld, D.I. (2003). Visual mismatch negativity. In: Polich, J.(ed.) *Detection of change: Event-related potential and fMRI findings*. Kluwer Academic Press, Boston, pp. 41-59.
- Horimoto, R., Inagaki, M., Yano, T., Sata, Y., & Kaga, M. (2002). Mismatch negativity of the color modality during a selective attention task to auditory stimuli in children with mental retardation. *Brain and Development*, 24, 703-709.
- Horimoto, R., Inagaki, M., Yano, T., Sata, Y., & Kaga, M. (2002). Mismatch negativity of the color modality during a selective attention task to auditory stimuli in children with mental retardation. *Brain & Development*, 24, 703–709.
- Hosák, L., Kremláček, J., Kuba, M., Libiger, J., & Čížek, J. (2008). Mismatch negativity in methamphetamine dependence: a pilot study. *Acta Neurobiologiae Experimentalis*, 68, 97–102.
- Hosoya, T., Baccus, S.A. & Meister, M. (2005). Dynamic predictive coding by the retina. *Nature*, 436, 71–77.
- Houlihan, M. E., Pritchard, W. S., & Robinson, J. H. (2001). Effects of smoking/nicotine on performance and event-related potentials during a short-term memory scanning task. *Psychopharmacology*, 156, 388-396.
- Ibbotson, M.R. (2005). Physiological mechanisms of adaptation in the visual system. In: *Fitting the Mind to the World: Adaptation and After-Effects in High-Level Vision*, edited by Clifford CWG, Rhodes G. New York: Oxford Univ. Press. p. 17–45.
- Iijima, M., Osawa, M., Nageishi, Y., Ushijima, R., & Iwata, M. (1995). PS-46-3 Visual mismatch negativity (MMN) in normal aging and dementing illness. *Electroencephalogr. Clin. Neuro.* 97, S205–S205.
- Iijima, M., Osawa, M., Nageishi, Y., Ushijima, R., & Iwata, M. (1996). Visual mismatch negativity (MMN) in aging. In: Ogura C. Koga Y. Shimokochi M. (Eds.) *Recent advances in event-related brain potentials research* (pp. 804–809). Amsterdam, The Netherlands: Elsevier.
- Ioannidis, J. P. A, & Trikalinos, T. A. (2005). Early extreme contradictory estimates may appear in published research: The Proteus phenomenon in molecular genetics research and randomized trials. *Journal of Clinical Epidemiology*, 58, 543–549.
doi:10.1016/j.jclinepi.2004.10.019
- Jääskeläinen, I. P., Alho, K., Escera, C., Winkler, I., Sillanauke, P., & Näätänen, R. (1996). Effects of ethanol and auditory distraction on forced choice reaction time. *Alcohol*, 13, 153–156.
doi:10.1016/0741-8329(95)02027-6

Jacobsen, T., & Schroger, E. (2001). Is there pre-attentive memory-based comparison of pitch? *Psychophysiol.* 38, 723-727.

James, T. W., & Gauthier, I. (2006). Repetition-induced changes in BOLD response reflect accumulation of neural activity. *Human Brain Mapping*, 27, 37–46.

Javitt, D.C. (2004). Glutamate as a therapeutic target in psychiatric disorders. *Mol Psychiatry*, 979, 984-997.

Javitt, D.C. (2009). Sensory Processing in Schizophrenia: Neither Simple nor Intact. *Schizophrenia Bulletin*, 35, 1059–1064. doi:10.1093/schbul/sbp110

Javitt, D.C. (2012). Twenty-five Years of Glutamate in Schizophrenia: Are We There Yet? *Schizophr. Bull.*, 38, 911-913.

Jessen, F., Amariglio, R. E., van Boxtel, M., Breteler, M., Ceccaldi, M., Chételat, G., ... Wagner, M. (2014). A conceptual framework for research on subjective cognitive decline in preclinical Alzheimer's disease. *Alzheimers Dement.* 10, 844-852. doi:10.1016/j.jalz.2014.01.001

Kähkönen, S., Yamashita, H., Rytsälä, H., Suominen, K., Ahveninen, J., & Isometsä, E. (2007). Dysfunction in early auditory processing in major depressive disorder revealed by combined MEG and EEG. *Journal of Psychiatry and Neuroscience*, 32, 316–322.

Kaliukhovich, D.A., & Vogels, R. (2014). Neurons in macaque inferior temporal cortex show no surprise response to deviants in visual oddball sequences. *Journal of Neuroscience*, 34, 12801–12815.

Kecskés-Kovács, K., Sulykos, I., & Czigler, I. (2013). Visual mismatch negativity is sensitive to symmetry as a perceptual category. *Eur J Neurosci.* 37, 662-667.

Kecskés-Kovács, K., Sulykos, I., & Czigler, I. (2013a). Visual mismatch negativity is sensitive to symmetry as a perceptual category. *Eur. J. Neurosci.* 37, 662-667.

Kecskés-Kovács, K., Sulykos, I., & Czigler, I. (2013b). Is it a face of a woman or a man? Visual mismatch negativity is sensitive to gender category. *Front. Hum. Neurosci.* 7:532. doi: 10.3389/fnhum.2013.00532.

Kemner, C., Verbaten, M. N., Cuperus, J. M., Camfferman, G., & Van Engeland, H. (1994). Visual and somatosensory event-related brain potentials in autistic children and three different control groups. *Electroencephalography and Clinical Neurophysiology/Evoked Potentials Section*, 92, 225–237. doi:10.1016/0168-5597(94)90066-3

Kenemans, J. L., Baas, J. M. ., Mangun, G., Lijffijt, M., & Verbaten, M. (2000). On the processing of spatial frequencies as revealed by evoked-potential source modeling. *Clinical Neurophysiology*, 111, 1113–1123. doi:10.1016/S1388-2457(00)00270-4

- Kenemans, J. L., Verbaten, M. N., Meelis, C. J., & Slagten, J. L. (1992). Visual stimulus change and the orienting reaction: event-related potential evidence for a two-stage process. *Biological Psychology*, 33, 97–114. doi:10.1016/0301-0511(92)90026-Q
- Kenemans, J.L., Jong, T.G., & Verbaten, M.N. (2003). Detection of visual change: Mismatch or rareness? *Neuroreport*, 14, 1239-1242.
- Keshavan, M. S., Dick, R. M., Diwadkar, V. a, Montrose, D. M., Prasad, K. M., & Stanley, J. A. (2009). Striatal metabolic alterations in non-psychotic adolescent offspring at risk for schizophrenia: a (1)H spectroscopy study. *Schizophrenia Research*, 115, 88–93. doi:10.1016/j.schres.2009.08.012
- Kessler, H., Roth, J., von Wietersheim, J., Deighton, R. M., & Traue, H. C. (2007). Emotion recognition patterns in patients with panic disorder. *Depression and Anxiety*, 24, 223–226. doi:10.1002/da.20223
- Kessler, R. (2012). The Costs of Depression. *Psychiatric Clinics of North America*, 35, 1–14. doi:10.1016/j.psc.2011.11.005
- Kessler, R., Berglund, P., Demler, O., Jin, R., Koretz, D., Merikangas, K. R., Rush, A.J., Walters, E.E., Wang, P.S., & National Comorbidity Survey Replication. (2003). The Epidemiology of Major Depressive Disorder: results from the National Comorbidity Survey Replication (NCS-R). *JAMA*, 289, 3095-3105. doi:10.1001/jama.289.23.3095
- Kiebel, S.J., Garrido, M.I., & Friston, K.J. (2007). Dynamic causal modelling of evoked responses: the role of intrinsic connections. *Neuroimage*. 36, 332-345.
- Kimura, M. (2012). Visual mismatch negativity and unintentional temporal-context-based prediction in vision. *Int. J. Psychophysiol.* 83, 144–155. doi:10.1016/j.ijpsycho.2011.11.010
- Kimura, M., Katayama, J., & Murohashi, H. (2006). Probability-independent and -dependent ERPs reflecting visual change detection. *Psychophysiol.* 43, 180-189.
- Kimura, M., Katayama, J., Ohira, H., & Schröger, E. (2009). Visual mismatch negativity: new evidence from the equiprobable paradigm. *Psychophysiol.* 46, 402-409.
- Kimura, M., Kondo, H., Ohira, H., & Schröger, E. (2012). Unintentional temporal context-based prediction of emotional faces: an electrophysiological study. *Cerebral Cortex.* 22, 1774–1785. doi:10.1093/cercor/bhr244
- Kimura, M., Schröger, E., & Czigler, I. (2011). Visual mismatch negativity and its importance in visual cognitive sciences. *Neuroreport.* 22, 669-673. doi: 10.1097/WNR.0b013e32834973ba.
- Kimura, M., Widmann, A., & Schröger, E. (2010). Human visual system automatically represents large-scale sequential regularities. *Brain Res.* 1317, 165-179.

- Rócsis, B., Brown, R.E., McCarley, R.W., & Hajos, M. (2013). Impact of ketamine on neuronal network dynamics: translational modeling of schizophrenia-relevant deficits. *CNS Neurosci Ther.* 19, 437-47.
- Kohler, C.G., Turner, T.H., Bilker, W.B., Brensinger, C.M., Siegel, S.J., Kanes, S.J., Gur, R.E., & Gur, R.C. (2003). Facial emotion recognition in schizophrenia: intensity effects and error pattern. *Am. J. Psychiatry* 160, 1768–1774.
- Kohn, A. (2007). Visual adaptation: physiology, mechanisms, and functional benefits. *J. Neurophysiol.* 97, 3155-3164.
- Kok, P., & de Lange, F.P. (2014). Shape perception simultaneously up- and downregulates neural activity in the primary visual cortex. *Curr. Biol.* 24, 1531-1535. doi: 10.1016/j.cub.2014.05.042.
- Kok, P., Jehee, J.F., & de Lange, F.P. (2012). Less is more: expectation sharpens representations in the primary visual cortex. *Neuron*, 75, 265–270.
- Komlosi, S., Csukly, G., Stefanics, G., Czigler, I., Bitter, I., & Czobor, P. (2013). Fearful face recognition in schizophrenia: an electrophysiological study. *Schizophr. Res.* 149, 135–140.
- Kreegipuu, K., Kuldkepp, N., Sibolt, O., Toom, M., Allik, J., & Näätänen, R. (2013). vMMN for schematic faces: automatic detection of change in emotional expression. *Front. Hum. Neurosci.* 7:714. doi: 10.3389/fnhum.2013.00714.
- Krekelberg, B., Boynton, G., & van Wezel, R.J.A. (2006). Adaptation: from single cells to BOLD signals. *Trends Neurosci.* 29, 250–256.
- Kremláček, J., Hosák, L., Kuba, M., Libiger, J., & Čížek, J. (2008). Visual information processing in recently abstaining methamphetamine-dependent individuals: evoked potentials study. *Documenta ophthalmologica*, 117, 245-255.
- Kremláček, J., Hulan, M., Kuba, M., Kubova, Z., Langrova, J., Vit, F., & Szanyi, J. (2012). Role of latency jittering correction in motion-onset VEP amplitude decay during prolonged visual stimulation. *Documenta Ophthalmologica. Advances in Ophthalmology*, 124, 211–23. doi:10.1007/s10633-012-9321-6
- Kremláček, J., Kuba, M., Kubová, Z., Langrová, J., Kubova, Z., & Langrova, J. (2006). Visual mismatch negativity elicited by magnocellular system activation. *Vision Research*, 46, 485–90.
- Kremláček, J., Kuba, M., Kubova, Z., Langrova, J., Vit, F., & Szanyi, J. (2007). Within-session reproducibility of motion-onset VEPs: effect of adaptation/habituation or fatigue on N2 peak amplitude and latency. *Doc Ophthalmol*, 115, 95–103. doi:10.1007/s10633-007-9063-z

- Kriegeskorte, N., Simmons, W. K., Bellgowan, P. S. F., & Baker, C. I. (2009). Circular analysis in systems neuroscience: the dangers of double dipping. *Nature Neuroscience*, 12, 535–540. doi:10.1038/nn.2303
- Kristjansson, A. (2011). The functional benefits of tilt adaptation. *Seeing and Perceiving*, 24, 37–51.
- Kuba, M., Kremláček, J., Langrova, J., Kubova, Z., Szanyi, J., & Vit, F. (2012). Aging effect in pattern, motion and cognitive visual evoked potentials. *Vision Research*, 62, 9–16. doi:10.1016/j.visres.2012.03.014
- Kubová, Z., Kuba, M., Kremláček, J., Langrová, J., Szanyi, J., Vít, F., & Chutná, M. (2015). Comparison of visual information processing in school-age dyslexics and normal readers via motion-onset visual evoked potentials. *Vision Research*, 111, 97–104. doi:10.1016/j.visres.2015.03.027
- Kuhl, P. K., Coffey-Corina, S., Padden, D., & Dawson, G. (2005). Links between social and linguistic processing of speech in children with autism: behavioural and electrophysiological measures. *Developmental Science*, 8, 1–12.
- Kujala, T., Alho, K., & Näätänen, R. (2000). Cross-modal reorganization of human cortical functions. *Trends in Neurosciences*, 23, 115–120
- Kuldkepp, N., Kreegipuu, K., Raidvee, A., Näätänen, R., & Allik, J. (2013). Unattended and attended visual change detection of motion as indexed by event-related potentials and its behavioral correlates. *Front. Hum. Neurosci.* 7:476. doi: 10.3389/fnhum.2013.00476
- Langrová, J., Kuba, M., Kremláček, J., Kubová, Z., & Vít, F. (2006). Motion-onset VEPs reflect long maturation and early aging of visual motion-processing system. *Vision Research*, 46, 536–544. doi:10.1016/j.visres.2005.06.024
- Lee, S.-H., Kwan, A. C., Zhang, S., Phoumthipphavong, V., Flannery, J. G., Masmanidis, S. C., Taniguchi, H., Huang, Z.J., Zhang, F., Boyden, E.S., Deisseroth, K., & Dan, Y. (2012). Activation of specific interneurons improves V1 feature selectivity and visual perception. *Nature*, 488, 379–383. doi:10.1038/nature11312
- Lee, T.S., & Mumford, D. (2003). Hierarchical Bayesian inference in the visual cortex. *Journal of Optical Society of America, A*, 20, 1434-1448.
- Lepistö, T., Soininen, M., Čeponiene, R., Almqvist, F., Näätänen, R., & Aronen, E. T. (2004). Auditory event-related potential indices of increased distractibility in children with major depression. *Clinical Neurophysiology*, 115, 620–627. doi:10.1016/j.clinph.2003.10.020
- Lewis, D. (2014). Inhibitory neurons in human cortical circuits: substrate for cognitive dysfunction in schizophrenia. *Current Opinion in Neurobiology*, 26, 22–26. doi:10.1016/j.conb.2013.11.003

- Light, G. A., & Naatanen, R. (2013). Mismatch negativity is a breakthrough biomarker for understanding and treating psychotic disorders. *Proceedings of the National Academy of Sciences of the United States of America*, 110, 15175–15176. doi:10.1073/pnas.1313287110
- Lindenberger, U., & Baltes, P. B. (1994). Sensory functioning and intelligence in old age: a strong connection. *Psychology and Aging*, 9, 339–355. doi:10.1037/0882-7974.9.3.339
- Liu, T., & Shi, J. (2008). Event-related potentials during preattentive processing of color stimuli. *Neuroreport* 19, 1221-1225.
- Lorenzo-López, L., Amenedo, E., Pazo-Alvarez, P., & Cadaveira, F. (2004). Pre-attentive detection of motion direction changes in normal aging. *NeuroReport*, 15, 2633-2636.
- Lorenzo-López, L., Amenedo, E., Pazo-Alvarez, P., & Cadaveira, F. (2004). Pre-attentive detection of motion direction changes in normal aging. *Neuroreport*, 15, 2633–2636.
- Luck, S. J., Mathalon, D. H., O'Donnell, B. F., Hmlinen, M. S., Spencer, K. M., Javitt, D. C., & Uhlhaas, P. J. (2011). A roadmap for the development and validation of event-related potential biomarkers in schizophrenia research. *Biological Psychiatry*, 70, 28–34. doi:10.1016/j.biopsych.2010.09.021
- Maekawa, T., Goto, Y., Kinukawa, N., Taniwaki, T., Hanbu, S., & Tobimatsu, S. (2005). Functional characterization of mismatch negativity to visual stimulus. *Clin. Neurophysiol.* 116, 2392-2402.
- Maekawa, T., Katsuki, S., Kishimoto, J., Onitsuka, T., Ogata, K., Yamasaki, T., Ueno, T., Tobimatsu, S., & Kanba, S. (2013). Altered visual information processing systems in bipolar disorder: evidence from visual MMN and P3. *Front. Hum. Neurosci.* 7:403. doi:10.3389/fnhum.2013.00403
- Maekawa, T., Tobimatsu, S., Inada, N., Oribe, N., Onitsuka, T., Kanba, S., & Kamio, Y. (2011). Top-down and bottom-up visual information processing of non-social stimuli in high-functioning autism spectrum disorder. *Research in Autism Spectrum Disorders*, 5, 201–209. doi:10.1016/j.rasd.2010.03.012
- Marsman, A., van den Heuvel, M. P., Klomp, D. W. J., Kahn, R. S., Luijten, P. R., & Hulshoff Pol, H. E. (2013). Glutamate in schizophrenia: a focused review and meta-analysis of 1H-MRS studies. *Schizophrenia Bulletin*, 39, 120–129. doi:10.1093/schbul/sbr069
- Martin, L. F., Davalos, D. B., & Kisley, M. A. (2009). Nicotine enhances automatic temporal processing as measured by the mismatch negativity waveform. *Nicotine & Tobacco Research*, 11, 698-706.

- Matsuyoshi, D., Morita, T., Kochiyama, T., Tanabe, H.C., Sadato, N., & Kakigi, R. (2015). Dissociable cortical pathways for qualitative and quantitative mechanisms in the face inversion effect. *J. Neurosci.* 35, 4268-4279. doi: 10.1523/JNEUROSCI.3960-14.2015
- May, P. J. C., & Tiitinen, H. (2010). Mismatch negativity (MMN), the deviance-elicited auditory deflection, explained. *Psychophysiol.* 47, 66–122. doi:10.1111/j.1469-8986.2009.00856.x
- Mazza, V., Turatto, M., & Sarlo, M. (2005). Rare stimuli or rare changes: What really matters for the brain? *Neuroreport* 16, 1061-1064.
- McDermott, K. C., Malkoc, G., Mulligan, J. B., & Webster, M. A. (2010). Adaptation and visual salience. *Journal of Vision*, 10:17, doi:10.1167/10.13.17.
- McGrath, J., Saha, S., Chant, D., & Welham, J. (2008). Schizophrenia: a concise overview of incidence, prevalence, and mortality. *Epidemiologic Reviews*, 30, 67–76. doi:10.1093/epirev/mxn001
- Meyer, T., Ramachandran, S., & Olson, C.R. (2014). Statistical learning of serial visual transitions by neurons in monkey inferotemporal cortex. *J. Neurosci.* 34, 9332-7. doi: 10.1523/JNEUROSCI.1215-14.2014.
- Mineka, S., & Sutton, S. K. (1992). Cognitive Biases and the Emotional Disorders. *Psychological Science*, 3, 65–69. doi:10.1111/j.1467-9280.1992.tb00260.x
- Mo, L., Xu, G., Kay, P., & Tan, L.H. (2011). Electrophysiological evidence for the left-lateralized effect of language on preattentive categorical perception of color. *Proc. Natl. Acad. Sci. USA* 108, 14026-14030.
- Moghaddam B., & Javitt D. (2012). From revolution to evolution: the glutamate hypothesis of schizophrenia and its implication for treatment. *Neuropsychopharmacology*, 37, 4-15.
- Morris, R.W., Weickert, C.S., & Loughland, C.M. (2009). Emotional face processing in schizophrenia. *Curr. Opin. Psychiatry.* 22, 140–146.
- Motter, B.C. (2006). Modulation of transient and sustained response components of V4 neurons by temporal crowding in flashed stimulus sequences. *J. Neurosci.*, 26, 9683–9694.
- Müller, D., Widmann, A., & Schröger, E. (2013). Object-related regularities are processed automatically: evidence from the visual mismatch negativity. *Front. Hum. Neurosci.* 7:259. doi: 10.3389/fnhum.2013.00259
- Müller, D., Winkler, I., Roeber, U., Schaffer, S., Czigler, I., & Schröger, E. (2010). Visual object representations can be formed outside the focus of voluntary attention: evidence from event-related brain potentials. *J. Cogn. Neurosci.* 22, 1179-1188.
- Müller, J.R., Metha, A.B., Krauskopf, J. & Lennie, P. (1999). Rapid adaptation in visual cortex to the structure of images. *Science*, 285, 1405–1408.

- Näätänen, R., & Picton, T. (1987). The N1 wave of the human electric and magnetic response to sound: a review and an analysis of the component structure. *Psychophysiology*, 24, 375–425. doi:10.1111/j.1469-8986.1987.tb00311.x
- Näätänen, R., Astikainen, P., Ruusuvirta, T., & Huotilainen, M. (2010). Automatic auditory intelligence: an expression of the sensory-cognitive core of cognitive processes. *Brain Res. Rev.* 64, 123-136.
- Näätänen, R., Gaillard, A. W., & Mantysalo, S. (1978). Early selective-attention effect on evoked potential reinterpreted. *Acta Psychologica*, 42, 313–329.
- Näätänen, R., Kujala, T., Escera, C., Baldeweg, T., Kreegipuu, K., Carlson, S., & Ponton, C. (2012). The mismatch negativity (MMN)-a unique window to disturbed central auditory processing in ageing and different clinical conditions. *Clin. Neurophysiol.* 123, 424–58. doi:10.1016/j.clinph.2011.09.020
- Näätänen, R., Kujala, T., Kreegipuu, K., Carlson, S., Escera, C., Baldeweg, T., & Ponton, C. (2011). The mismatch negativity: an index of cognitive decline in neuropsychiatric and neurological diseases and in ageing. *Brain*, 134, 3435-53.
- Näätänen, R., Kujala, T., Kreegipuu, K., Carlson, S., Escera, C., Baldeweg, T., & Ponton, C. (2011). The mismatch negativity: An index of cognitive decline in neuropsychiatric and neurological diseases and in ageing. *Brain*. 134, 3432–3450. doi:10.1093/brain/awr064
- Näätänen, R., Paavilainen, P., Rinne, T., & Alho, K. (2007). The mismatch negativity (MMN) in basic research of central auditory processing: A review. *Clin. Neurophysiol.* 118, 2544-2590. doi:10.1016/j.clinph.2007.04.026
- Näätänen, R., Schröger, E., Karakas, S., Tervaniemi, M., & Paavilainen, P. (1993). Development of a memory trace for a complex sound in the human brain. *Neuroreport*. 4, 503-506.
- Näätänen, R., Sussman, E. S., Salisbury, D., & Shafer, V. L. (2014). Mismatch negativity (MMN) as an index of cognitive dysfunction. *Brain Topography*, 27, 451–66. doi:10.1007/s10548-014-0374-6
- Nelken, I. (2014). Stimulus-specific adaptation and deviance detection in the auditory system: experiments and models. *Biol. Cybern.* 108, 655-663. doi: 10.1007/s00422-014-0585-7.
- Neuhaus, A. H., Brandt, E. S. L., Goldberg, T. E., Bates, J. a, & Malhotra, A. K. (2013). Evidence for impaired visual prediction error in schizophrenia. *Schizophrenia Research*, 147, 326–330. doi:10.1016/j.schres.2013.04.004
- Newman, D., Speake, D. J., Armstrong, P. J., & Tiplady, B. (1997). Effects on ethanol on control of attention. *Human Psychopharmacology*, 12, 235-241.

- Norton, E. S., Beach, S. D., & Gabrieli, J. De. (2014). Neurobiology of dyslexia. *Current Opinion in Neurobiology*, 30C, 73–78. doi:10.1016/j.conb.2014.09.007
- Nunez, P. L., & Srinivasan, R. (2006). *Electric Fields of the Brain*. Oxford University Press. doi:10.1093/acprof:oso/9780195050387.001.0001
- Nyberg, L., Bäckman, L., Erngrund, K., Olofsson, U., & Nilsson, L. G. (1996). Age differences in episodic memory, semantic memory, and priming: relationships to demographic, intellectual, and biological factors. *J. Gerontol. B. Psychol. Sci. Soc. Sci.* 51, P234-40. doi:10.1093/geronb/51B.4.P234
- O'Shea, R. P. (2015). Refractoriness about adaptation. *Front. Hum. Neurosci.* 9:38. doi: 10.3389/fnhum.2015.00038
- Pammer, K. (2013). Temporal sampling in vision and the implications for dyslexia. *Front. Hum. Neurosci.* 7:933. doi:10.3389/fnhum.2013.00933
- Pang, X., Xu, J., Chang, Y., Tang, D., Zheng, Y., Liu, Y., & Sun, Y. (2014). Mismatch negativity of sad syllables is absent in patients with major depressive disorder. *PLoS ONE*, 9(3):e91995. doi:10.1371/journal.pone.0091995
- Park, D. C., Lautenschlager, G., Hedden, T., Davidson, N. S., Smith, A. D., & Smith, P. K. (2002). Models of visuospatial and verbal memory across the adult life span. *Psychology and Aging*, 17, 299–320. doi:10.1037/0882-7974.17.2.299
- Pazo-Alvarez, P., Cadaveira, F., & Amenedo, E. (2003). MMN in the visual modality: a review. *Biological Psychology*, 63, 199–236.
- Pérez-González, D., & Malmierca, M.S. (2014). Adaptation in the auditory system: an overview. *Front. Integr. Neurosci.* 8:19. doi: 10.3389/fnint.2014.00019
- Peterson, R. L., & Pennington, B. F. (2012). Developmental dyslexia. *Lancet*, 379, 1997–2007. doi:10.1016/S0140-6736(12)60198-6
- Phillips, M. L., Drevets, W. C., Rauch, S. L., & Lane, R. (2003). Neurobiology of emotion perception II: Implications for major psychiatric disorders. *Biological Psychiatry*. 54, 515-28. doi:10.1016/S0006-3223(03)00171-9
- Pomarol-Clotet, E., Honey, G.D., Murray, G.K., Corlett, P.R., Absalom, A.R., Lee, M., McKenna, P.J., Bullmore, E.T., & Fletcher, P.C. (2006). Psychological effects of ketamine in healthy volunteers. Phenomenological study. *Br. J. Psychiatry*. 189, 173-179.

- Qiu, X., Yang, X., Qiao, Z., Wang, L., Ning, N., Shi, J., Zhao, L., Yang, Y. (2011). Impairment in processing visual information at the pre-attentive stage in patients with a major depressive disorder: A visual mismatch negativity study. *Neuroscience Letters*, 491, 53–57.
- Ramachandran, S., Meyer, T., & Olson, C.R. (2016). Prediction suppression in monkey inferotemporal cortex depends on the conditional probability between images. *J. Neurophysiol.* 115, 355–362. doi: 10.1152/jn.00091.2015.
- Rao, R.P.N., & Ballard, D.H. (1999). Predictive coding in the visual cortex: a functional interpretation of some extraclassical receptive-field effects. *Nature Neuroscience* 2, 79 – 87.
- Riekkinen, P., Pääkkönen, A., Karhu, J., Partanen, J., Soininen, H., Laakso, M., & Riekkinen, P. (1997). THA disrupts mismatch negativity in Alzheimer disease. *Psychopharmacology*, 133, 203–206. doi:10.1007/s002130050392
- Rogers, S. J., & Ozonoff, S. (2005). Annotation: What do we know about sensory dysfunction in autism? A critical review of the empirical evidence. *Journal of Child Psychology and Psychiatry*, 46, 1255–1268. doi:10.1111/j.1469-7610.2005.01431.x
- Rowe, J. B., Hughes, L., & Nestor, P. J. (2009). Abnormal auditory mismatch negativity (MMN) in behavioural variant frontotemporal dementia. *NeuroImage*, 47(Supplement 1), pS90.
- Ruhnau, P., Herrmann, B., and Schröger, E. (2012). Finding the right control: The mismatch negativity under investigation. *Clin. Neurophysiol.* 123, 507-512.
- Ruzzoli, M., Pirulli, C., Brignani, D., Maioli, C., & Miniussi, C. (2012). Sensory memory during physiological aging indexed by mismatch negativity (MMN). *Neurobiology of Aging*, 33, 625.e21–30.
- Schacter, D.L., & Buckner, R.L. (1998). Priming and the brain. *Neuron*. 20, 185-95.
- Schröger, E. (1997). Higher-order processes in auditory-change detection: a response to Näätänen and Alho. *Trends in Cognitive Sciences*, 1, 45-46.
- Schröger, E., & Wolff, C. (1996). Mismatch response of the human brain to changes in sound localization. *NeuroReport*. 7, 3005-3008.
- Schulte-Körne, G., & Bruder, J. (2010). Clinical neurophysiology of visual and auditory processing in dyslexia: A review. *Clinical Neurophysiology*, 121, 1794–1809. doi:10.1016/j.clinph.2010.04.028

- Shepherd, R.K., Srivivasani, M.N., Nayagam, D.A., Williams, C.E., & Blamey, P.J. (2013). Visual prostheses for the blind. *Trends in Biotechnology*, 31, 562–71.
- Shi, L., Wu, J., Sun, G., Dang, L., & Zhao, L. (2013). Visual mismatch negativity in the “optimal” multi-feature paradigm. *Journal of Integrative Neuroscience*, 12, 247–58.
doi:10.1142/S0219635213500179
- Shimamura, A. P., Berry, J. M., Mangels, J. A., Rusting, C. L., & Jurica, P. J. (1995). Memory and Cognitive Abilities in University Professors: Evidence for Successful Aging. *Psychological Science*, 6, 271–277. doi:10.1111/j.1467-9280.1995.tb00510.x
- Si, C., Ren, C., Wang, P., Bian, H., Wang, H., & Yan, Z. (2014). Impairment in Preattentive Processing among Patients with Hypertension Revealed by Visual Mismatch Negativity. *BioMed Research International*, 2014, 1–8. doi:10.1155/2014/945121
- Sokolov, E. N. (1963). Higher nervous functions; the orienting reflex. *Annual Review of Physiology*, 25, 545–580. doi:10.1146/annurev.ph.25.030163.002553
- Solomon, S.G., Peirce, J.W., Dhruv, N.T. & Lennie, P. (2004). Profound contrast adaptation early in the visual pathway. *Neuron*, 42, 155–162.
- Stagg, C., Hindley, P., Tales, A., & Butler, S. (2004). Visual mismatch negativity: the detection of stimulus change. *Neuroreport* 15, 659-663.
- Stahl, SM. (2007). The genetics of schizophrenia converge upon the NMDA glutamate receptor. *CNS Spectr.*, 12, 583-588.
- Steele, C. M., & Josephs, R. A. (1988). Drinking your troubles away. II: An attention-allocation model of alcohol's effect on psychological stress. *Journal of Abnormal Psychology*, 97, 196–205.
doi:10.1037/0021-843X.97.2.196
- Stefanics G, & Czigler I. (2012). Automatic prediction error responses to hands with unexpected laterality: An electrophysiological study. *Neuroimage*, 63, 253-261.
- Stefanics G, Csukly G, Komlosi S, Czobor P, & Czigler I. (2012). Processing of unattended facial emotions: a visual mismatch negativity study. *Neuroimage*, 59, 3042–3049.
- Stefanics, G., Fosker, T., Huss, M., Mead, N., Szucs, D., & Goswami, U. (2011). Auditory sensory deficits in developmental dyslexia: A longitudinal ERP study. *NeuroImage*, 57, 723–732.
doi:10.1016/j.neuroimage.2011.04.005
- Stefanics, G., Háden, G., Huotilainen, M., Balázs, L., Sziller, I., Beke, A., Fellman, V., & Winkler, I. (2007) Auditory temporal grouping in newborn infants. *Psychophysiol.* 44, 697–702.
- Stefanics, G., Háden, G., Sziller, I., Balázs, L., Beke, A., & Winkler, I. (2009). Newborn infants process pitch intervals. *Clin. Neurophysiol.* 120, 304-308.

- Stefanics, G., Kremláček, J., & Czigler, I. (2016). Mismatch Negativity and Neural Adaptation: Two Sides of the Same Coin. Response to: A Commentary on 'Visual mismatch negativity: a predictive coding view'. *Front. Hum. Neurosci.* 10:13. doi: 10.3389/fnhum.2016.00013
- Stefanics, G., Kremláček, J., & Czigler, I. I. (2014). Visual mismatch negativity: a predictive coding view. *Front. Hum. Neurosci.* 8:666. doi:10.3389/fnhum.2014.00666
- Stein, J. F. (2012). Visual Contributions to Reading Difficulties: The Magnocellular Theory. In J. Stein & Z. Kapoula (Eds.), *Visual Aspects of Dyslexia* (pp. 171–198). Oxford: Oxford University Press. doi:10.1093/acprof:oso/9780199589814.001.0001
- Stephan, K.E., Bach, D.R., Fletcher, P.C., Flint, J., Frank, M.J., Friston, K.J., Heinz, A., Huys, Q.J., Owen, M.J., Binder, E.B., Dayan, P., Johnstone, E.C., Meyer-Lindenberg, A., Montague, P.R., Schnyder, U., Wang, X.J., & Breakspear, M. (2016a). Charting the landscape of priority problems in psychiatry, part 1: classification and diagnosis. *Lancet Psychiatry.* 3, 77-83. doi: 10.1016/S2215-0366(15)00361-2.
- Stephan, K.E., Baldeweg, T., & Friston, K.J. (2006). Synaptic plasticity and dysconnection in schizophrenia. *Biol. Psychiatry,* 59, 929-939.
- Stephan, K.E., Binder, E.B., Breakspear, M., Dayan, P., Johnstone, E.C., Meyer-Lindenberg, A., Schnyder, U., Wang, X.J., Bach, D.R., Fletcher, P.C., Flint, J., Frank, M.J., Heinz, A., Huys, Q.J., Montague, P.R., Owen, M.J., & Friston, K.J. (2016b). Charting the landscape of priority problems in psychiatry, part 2: pathogenesis and aetiology. *Lancet Psychiatry.* 3, 84-90. doi: 10.1016/S2215-0366(15)00360-0.
- Stephan, K.E., Friston, K.J., & Frith, C.D. (2009). Dysconnection in schizophrenia: from abnormal synaptic plasticity to failures of self-monitoring. *Schizophr. Bull.* 35, 509-27.
- Stothart, G., Kazanina, N., Näätänen, R., Haworth, J., & Tales, A. (2014). Early Visual Evoked Potentials and Mismatch Negativity in Alzheimer's Disease and Mild Cognitive Impairment. *J. Alzheimers Dis.* 44, 397-408. doi: 10.3233/JAD-140930.
- Stothart, G., Tales, A., & Kazanina, N. (2013). Evoked potentials reveal age-related compensatory mechanisms in early visual processing. *Neurobiology of Aging,* 34, 1302–1308. doi:10.1016/j.neurobiolaging.2012.08.012
- Sulykos, I., & Czigler, I. (2011). One plus one is less than two: Visual features elicit non-additive mismatch-related brain activity. *Brain Res.* 1398, 64-71.
- Sulykos, I., & Czigler, I. (2014). Visual mismatch negativity is sensitive to illusory brightness changes. *Brain Res.* 1561, 48-59. doi: 10.1016/j.brainres.2014.03.008.

- Sulykos, I., & Czigler, I. (2011). One plus one is less than two: visual features elicit non-additive mismatch-related brain activity. *Brain Res.* 1398, 64-71. doi: 10.1016/j.brainres.2011.05.009.
- Summerfield, C., & de Lange, F.P. (2014). Expectation in perceptual decision making: neural and computational mechanisms. *Nat. Rev. Neurosci.* 15, 745-56. doi: 10.1038/nrn3838.
- Summerfield, C., Trittschuh, E.H., Monti, J.M., Mesulam, M.M., & Egner, T. (2008). Neural repetition suppression reflects fulfilled perceptual expectations. *Nat. Neurosci.* 11, 1004–1006.
- Susac, A., Ilmoniemi, R. J., Pihko, E., & Supek, S. (2003). Neurodynamic Studies on Emotional and Inverted Faces in an Oddball Paradigm. *Brain Topography*, 16, 265–268. doi:10.1023/B:BRAT.0000032863.39907.cb
- Susac, A., Ilmoniemi, R.J., Pihko, E., & Supek, S. (2004). Neurodynamic studies on emotional and inverted faces in an oddball paradigm. *Brain Topogr.* 16, 265-268.
- Susac, A., Ilmoniemi, R.J., Pihko, E., Ranken, D., & Supek, S. (2010). Early cortical responses are sensitive to changes in face stimuli. *Brain Res.* 1346, 155-164.
- Takei, Y., Kumano, S., Hattori, S., Uehara, T., Kawakubo, Y., Kasai, K., Fukuda, M., & Mikuni, M. (2009). Preattentive dysfunction in major depression: A magnetoencephalography study using auditory mismatch negativity. *Psychophysiol.* 46, 52–61. doi:10.1111/j.1469-8986.2008.00748.x
- Takei, Y., Kumano, S., Maki, Y., Hattori, S., Kawakubo, Y., Kasai, K., Fukuda, M., & Mikuni, M. (2010). Preattentive dysfunction in bipolar disorder: A MEG study using auditory mismatch negativity. *Prog Neuropsychopharmacol. Biol. Psychiatry.* 34, 903–912. doi:10.1016/j.pnpbp.2010.04.014
- Tales, A., & Butler, S. (2006). Visual mismatch negativity highlights abnormal preattentive visual processing in Alzheimer’s disease. *Neuroreport*, 17, 887–890. doi:10.1097/01.wnr.0000223383.42295.fa 00001756-200606260-00008 [pii]
- Tales, A., Haworth, J., Wilcock, G., Newton, P., & Butler, S. (2008). Visual mismatch negativity highlights abnormal pre-attentive visual processing in mild cognitive impairment and Alzheimer’s disease. *Neuropsychologia*, 46, 1224–1232.
- Tales, A., Troscianko, T., Wilcock, G. K., Newton, P., & Butler, S. R. (2002). Age-related changes in the preattentive detection of visual change. *Neuroreport*, 13, 969–972.
- Tanaka, M., Okubo, O., Fuchigami, T., & Harada, K. (2001). A study of mismatch negativity in newborns. *Pediatrics International*, 43, 281–286. doi:10.1046/j.1442-200x.2001.01395.x
- Tang, D., Xu, J., Chang, Y., Zheng, Y., Shi, N., Pang, X., & Zhang, B. (2013). Visual mismatch negativity in the detection of facial emotions in patients with panic disorder. *NeuroReport*, 24, 207–211. doi:10.1097/WNR.0b013e32835eb63a

- Thies, W., & Bieler, L. (2013). 2013 Alzheimer's disease facts and figures: Alzheimer's Association Report. *Alzheimers Dement.* 9, 208–45. doi:10.1016/j.jalz.2013.02.003
- Tiitinen, H., May, P., Reinikainen, K., & Näätänen, R. (1994). Attentive novelty detection in humans is governed by pre-attentive sensory memory. *Nature.* 372, 90–92.
- Todd, J., Harms, L., Schall, U., & Michie, P. T. (2013). Mismatch Negativity: Translating the Potential. *Front. Psychiatry.* 4:171. doi:10.3389/fpsy.2013.00171
- Todd, J., Michie, P. T., Schall, U., Karayanidis, F., Yabe, H., & Näätänen, R. (2008). Deviant Matters: Duration, Frequency, and Intensity Deviants Reveal Different Patterns of Mismatch Negativity Reduction in Early and Late Schizophrenia. *Biological Psychiatry,* 63, 58–64. doi:10.1016/j.biopsych.2007.02.016
- Todd, J., Michie, P. T., Schall, U., Ward, P. B., & Catts, S. V. (2012). Mismatch negativity (MMN) reduction in schizophrenia—Impaired prediction-error generation, estimation or salience? *International Journal of Psychophysiology,* 83, 222–231. doi:10.1016/j.ijpsycho.2011.10.003
- Tomio, N., Fuchigami, T., Fujita, Y., Okubo, O., & Mugishima, H. (2012). Developmental Changes of Visual Mismatch Negativity. *Neurophysiology,* 44, 138–143. doi:10.1007/s11062-012-9280-2
- Umbricht, D., & Krljes, S. (2005). Mismatch negativity in schizophrenia: a meta-analysis. *Schizophrenia Research,* 76, 1–23. doi:10.1016/j.schres.2004.12.002
- Umbricht, D., Schmid, L., Koller, R., Vollenweider, F. X., Hell, D., & Javitt, D. C. (2000). Ketamine-Induced Deficits in Auditory and Visual Context-Dependent Processing in Healthy Volunteers. *Archives of General Psychiatry,* 57, 1139–1147. doi:10.1001/archpsyc.57.12.1139
- Urban, A., Kremláček, J., Masopust, J., & Libiger, J. (2008). Visual mismatch negativity among patients with schizophrenia. *Schizophrenia Research,* 102, 320–328. doi:10.1016/j.schres.2008.03.014
- Vidyasagar, T. R., & Pammer, K. (2010). Dyslexia: a deficit in visuo-spatial attention, not in phonological processing. *Trends in Cognitive Sciences,* 14, 57–63. doi:10.1016/j.tics.2009.12.003
- Vogels, R. (2015). Sources of adaptation of inferior temporal cortical responses. *Cortex* (in press), <http://dx.doi.org/10.1016/j.cortex.2015.08.024>
- Wacongne, C., Changeux, J.P., & Dehaene, S. (2012). A neuronal model of predictive coding accounting for the mismatch negativity. *J. Neurosci.* 32, 3665–3678.
- Wang, J. J., Bi, H. Y., Gao, L. Q., & Wydell, T. N. (2010). The visual magnocellular pathway in Chinese-speaking children with developmental dyslexia. *Neuropsychologia,* 48(12), 3627–3633. doi:S0028-3932(10)00369-6 [pii] 10.1016/j.neuropsychologia.2010.08.015

Eperjesi, F., Ford, S., Larøi, F. (2014). Visual hallucinations in the psychosis spectrum and comparative information from neurodegenerative disorders and eye disease. *Schizophrenia Bulletin*, 40 Suppl 4, S233–S245. doi:10.1093/schbul/sbu036

- Widmann, A., Schröger, E., Tervaniemi, M., Pakarinen, S., & Kujala, T. (2012). Mapping symbols to sounds: electrophysiological correlates of the impaired reading process in dyslexia. *Frontiers in Psychology*, 3, 60
- Wig, G.S., Grafton, S.T., Demos, K.E., & Kelley, W.M. (2005). Reductions in neural activity underlie behavioral components of repetition priming. *Nat Neurosci.* 8, 1228-1233.
- Wiggs, C.L., & Martin, A. (1998). Properties and mechanisms of perceptual priming. *Current Opinion in Neurobiology*, 8, 227–233.
- Winkler, I. (2007). Interpreting the Mismatch Negativity. *Journal of Psychophysiology*, 21(3), 147–163.
- Winkler, I., & Czigler, I. (2012). Evidence from auditory and visual event-related potential (ERP) studies of deviance detection (MMN and vMMN) linking predictive coding theories and perceptual object representations. *Int. J. Psychophysiol.* 83, 132-143. doi:10.1016/j.ijpsycho.2011.10.001
- Winkler, I., Karmos, G., & Näätänen, R. (1996). Adaptive modelling of the unattended acoustic environment reflected in the mismatch negativity event-related potential. *Brain Res.* 742, 239–252.
- Wolfe, J.M. (1999). Inattentive amnesia. In Coltheart, V. (Ed.), *Fleeting memories* (pp.71-94). Cambridge, MA: MIT Press
- Wolkenstein, L., Schönenberg, M., Schirm, E., & Hautzinger, M. (2011). I can see what you feel, but i can't deal with it: Impaired theory of mind in depression. *Journal of Affective Disorders*, 132, 104–111. doi:10.1016/j.jad.2011.02.010
- Yehezkel, O., Sagi, D., Sterkin, A., Belkin, M., & Polat, U. (2010). Learning to adapt: Dynamics of readaptation to geometrical distortions. *Vision Research*, 50, 1550–1558.
- Yoon, JH., Sheremata, S.L., Rokem, A., & Silver, M.A. (2013). Windows to the soul: vision science as a tool for studying biological mechanisms of information processing deficits in schizophrenia. *Front Psychol.* 4:681. doi: 10.3389/fpsyg.2013.00681.
- Zhao, L., & Li, J. (2006). Visual mismatch negativity elicited by facial expressions under non-attentional condition. *Neurosci. Lett.* 401, 126-131.
- Zhao, L., & Li, J. (2006). Visual mismatch negativity elicited by facial expressions under non-attentional condition. *Neuroscience Letters*, 410, 126–31. doi:10.1016/j.neulet.2006.09.081

ACCEPTED MANUSCRIPT