



Swansea University
Prifysgol Abertawe



Cronfa - Swansea University Open Access Repository

This is an author produced version of a paper published in :

Cronfa URL for this paper:

<http://cronfa.swan.ac.uk/Record/cronfa30478>

Paper:

Seage, C. & Lee, M. (2016). Do disinhibited eaters pay increased attention to food cues?.

<http://dx.doi.org/10.1016/j.appet.2016.09.031>

This article is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Authors are personally responsible for adhering to publisher restrictions or conditions. When uploading content they are required to comply with their publisher agreement and the SHERPA RoMEO database to judge whether or not it is copyright safe to add this version of the paper to this repository.

<http://www.swansea.ac.uk/iss/researchsupport/cronfa-support/>

Accepted Manuscript

Do disinhibited eaters pay increased attention to food cues?

C.H. Seage, M. Lee

PII: S0195-6663(16)30481-0

DOI: [10.1016/j.appet.2016.09.031](https://doi.org/10.1016/j.appet.2016.09.031)

Reference: APPET 3170

To appear in: *Appetite*


Received Date: 27 May 2015

Revised Date: 6 September 2016

Accepted Date: 27 September 2016

Please cite this article as: Seage C.H. & Lee M., Do disinhibited eaters pay increased attention to food cues?, *Appetite* (2016), doi: 10.1016/j.appet.2016.09.031.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Abstract

The Three Factors Eating Questionnaire's measure of disinhibited eating is a robust predictor of long-term weight gain. This experiment explored if disinhibited eaters display attentional bias to food cues. Participants (N=45) completed a visual dot probe task which measured responses to food (energy dense and low energy foods) and neutral cues. Picture pairs were displayed either for a 100 ms or 2000 ms duration. All participants displayed attentional bias for energy dense food items. Indices of attentional bias were largest in disinhibited eaters. Attentional bias in disinhibited eaters appeared to be underpinned by facilitated attention.

Key Words: Attentional Bias, Food, Orientation, Visual Dot Probe, Disinhibition

Introduction

Drug cues acquire higher motivational value through the process of dopaminergic conditioning (Berridge & Robinson, 1997). This associative learning leads to the reward system becoming hypersensitive to drugs and their associated cues (Robinson & Berridge, 2001). A frequently used behavioural measure of neural sensitivity to drug cues is attentional bias. Attentional bias occurs when an individual is quicker at processing personally relevant information compared to neutral information (Macloed, Matthews & Tata, 1986). Attentional bias for drug cues has been consistently documented in smokers, frequent caffeine consumers, drug users and alcoholics (For a review see Field and Cox, 2008). It is thought that attentional bias serves a functional role in maintaining addictive behaviour. Selective attention to drug cues has been shown to underpin approach behaviour and craving (Cox, Klinger & Fadardi, 2016). It is also a robust predictor of relapse (Franken, 2003).

Overeating provides an interesting parallel to addictive behaviour. Much like habitual drug users, obese individuals commonly report experiencing craving and a preoccupation with food (Herman and Polivy, 2008; Jastreboff, Sinha, Lacadie, Small, Sherwin & Potenza, 2013). The influence that food relevant cues (e.g. sight, smell, taste) have on food intake has also been well documented (for review see Herman & Polivy, 2008). It is plausible that dopaminergic conditioning occurs in individuals who habitually overeat. Attempts to establish if attentional bias for food cues can be a useful predictor of obesity risk has had mixed success. However, there is a growing body of research that demonstrated that obese individuals allocate greater attentional resources to food stimuli compared to their lean counterparts. (Castellanos et al. 2009; Nijs, Franken & Muris, 2010, Yokum, & Stice, 2011; Braet & Crombez, 2003; Graham, Hoover, Ceballos & Komogrotsev, 2011; Kemps, Tiggemann & Hollitt, 2014; Long, Hinton & Gillespie, 1994; Nijs, Muris, Euser & Franken, 2010; Werthmann et al., 2011).

A recent review of this literature by Doolan, Breslin, Hanna & Gallagher (2015) proposes that attentional bias to food cues is influenced more by an individual's eating traits than body

54 explanation has been used to explain the paradoxical relationship that exists between body
55 weight and restrained eating patterns. Repeated attempts by restrained eaters to limit their
56 food intake to control body weight, seemingly increases the likelihood that they will become
57 obese (Herman & Polivy, 1980). A number of studies have demonstrated that restrained eaters
58 have high indices of attentional bias to food cues (Hollitt, et al. 2007; Tapper, Pothos, Fadardi
59 & Ziori, 2008). It can be proposed that attempts to restrict calorie intake made by restrained
60 eaters are thwarted by biased processing of food cues. Higher indices of food processing bias
61 have been linked to other eating patterns that are associated with obesity risk; these include
62 external eaters (Brignell, Griffiths, Bradley, & Mogg, 2009; Newman, O'Connor & Conner,
63 2008) and high chocolate cravers (Smeets, Roefs, & Jansen, 2009).

64
65 To date, there has been no published attempt to document attentional bias in individuals who
66 experience disinhibited eating. This oversight limits the existing literature as the Three
67 Factors Eating Questionnaire's measure of disinhibited eating (TFEQ_D, Stunkard &
68 Messick, 1985) is viewed as one of the most robust predictors of long-term weight gain
69 (Hays & Roberts, 2008). Conceptually the term disinhibition refers to a variety of eating
70 behaviours that can be characterised by a lack of self-regulation (e.g. binge eating, unhealthy
71 food choices, low awareness of satiety) (Lattimore & Malinowski, 2008). Research has
72 shown that individuals who score high on measures of trait disinhibition consistently have
73 higher body weights (Boschi et al 2001; Provencher et al. 2003), make unhealthy food
74 choices (Contento, Zybert, & Williams, 2005; Lahteenmaki & Tuorila, 1995), are more
75 impulsive (Yeomans, Leitch, & Mobini, 2008) and experience reduced success from weight
76 loss interventions (Bryant, Caudwell, Hopkins, King & Blundell 2012). This paper aims to
77 examine if the opportunistic eating pattern displayed by disinhibited eaters is indicative of
78 increased attentional bias to food cues.

79
80 The present research examined if individuals who have high levels of disinhibited eating (as
81 measured by the TFEQ, Stunkard & Messick, 1985) paid increased attention to food cues
82 during a visual dot probe task. Two visual stimuli were briefly presented side by side,
83 followed by a dot (probe) where one of the stimuli had been. Some trials involved a food
84 picture and a neutral picture, and others contained two neutral pictures. Participants had to
85 press a button on the side of the display to indicate where the probe had appeared. Response
86 time (RT) was used to calculate attentional bias. Faster RTs on trials where the probe
87 followed in the location of a food picture, compared with trials when it followed one of two
88 neutral stimuli was indicative of increased attention to food stimuli. To explore the impact of
89 motivational value on attentional bias the food pictures consisted of both energy dense and
90 low energy food items (Tapper, Pothos & Lawrence, 2010). It was predicted that attentional
91 bias would increase for all participants when responding to trials containing foods which are
92 energy dense (due to the cues higher motivational value). However, it is anticipated that this
93 effect will be exacerbated in disinhibited eaters who are typically more responsive to the
94 presence of hedonic food cues (Tapper et al. 2010).

95
96 During the visual dot probe task, picture pairs were displayed for either 100ms or 2000ms
97 exposures. A matched neutral design was used to allow the reaction time data to be analysed
98 in a way that provides both a traditional measure of attentional bias, but also establishes
99 whether bias reflects facilitated attention to food cues or delayed disengagement (Tapper et al
100 2010; Koster, Crombez, Verschuere & Houwer, 2004). If attentional bias for food cues is
101 driven by facilitated attention participants will make quicker responses when the probe
102 replaces a congruent stimulus (probe position replacing food item). Whereas delayed

104 (probe position replacing neutral items).

105

106 **Method**

107

108 The sample comprised of forty-five participants who were recruited from the undergraduate
109 population of the University of Swansea. The mean age of participants was 20.5 ± 1.8 years.
110 The sample's mean BMI was within the normal range ($23.6 \pm 4.8 \text{ kg/m}^2$). Disinhibition was
111 measured using the disinhibition subscale of the Three Factor Eating Questionnaire (Stunkard
112 and Melleck, 1985). This measure explores an individual's level of uncontrolled eating using
113 9 items. All potential participants were asked to complete the TFEQ_D; those whose scores
114 placed them in the bottom or top 40% of the sample were invited to complete the visual dot
115 probe task. Participants were grouped in terms of high and low disinhibited eating based on
116 their TFEQ_D scores. Recruitment adhered to the following selection criteria; all participants
117 were non-vegan or vegetarian, self-reported that had no history of disordered eating and were
118 not dieting.

119

120 Laboratory sessions were scheduled so that they occurred after meal times, all participants ate
121 their habitual breakfast or lunch prior to attendance. This was to ensure that any behavioural
122 differences in task performance were not caused by hunger. On arrival, participants were
123 required to rate their hunger measured using a general mood questionnaire (VAS 0-100)
124 which contained 10 items. Participants were asked to rate their mood (e.g. on a scale of 0-
125 100 how happy are you feeling?) Included in these ratings were questions on hunger and
126 thirst). Participants were then introduced to the visual dot probe task and were informed that
127 they would be required to attend and respond to stimuli in the form of pictures. The test
128 stimuli consisted of 64 pairs of colour pictures. Sixteen pairs were an energy dense food and
129 a household item; sixteen were a low energy food and a household item, and 32 were two
130 household items. All stimuli used in this task had been previously rated in a pilot study as
131 being representative of each of the two categories (Tapper et al. 2008) and none of the
132 household items selected altered the context of the food stimuli (e.g. related to food
133 preparation, cleaning). In addition 10 animal items were used to create practice trials.

134

135 Picture pairs were presented for 100 ms and 2000ms duration across two blocks of 258 trials
136 (128 critical trials, 128 matched neutral trials). Each block contained 4 presentations of each
137 of the experimental or matched neutral picture pairs (e.g. experimental stimulus shown on
138 left, followed by a probe on the left; experimental stimulus on left, followed by a probe on
139 the right; experimental stimulus shown on the right, followed by a probe on the right and
140 experimental stimulus shown on right followed by a probe on the left). These presentations
141 were randomised. The probe used in this task was a dot and was displayed until the
142 participant made a response. Participants responded to the probe by identifying which side of
143 the screen the probe had appeared. This was done by pressing one of two response buttons.
144 Reaction time (RT) was measured in Milliseconds (ms). At the end of the computer task,
145 participants were asked again to rate current mood and hunger. Finally, participant's height
146 (cm) and weight (kg) were recorded. An average laboratory session lasted 45 minutes.
147 All trials with incorrect responses were excluded from the data analysis. RT for correct
148 choices that were > 200 ms and < 2000 ms and < 2 SD longer than the participant's mean
149 RT was analysed. Attentional bias scores were calculated for each participant and picture
150 duration by subtracting the mean RT for probes replacing food items from the mean RT for
151 probes replacing neutral items. Thus positive values would reflect a bias favouring a food
152 stimulus relative to a neutral stimulus.

154 Data Analysis

155
156 Task Accuracy was compared across the two groups using an x 2 (Stimulus Duration) x 2
157 (Stimuli Set) X 2 (TFEQ_D) ANOVA. Attentional bias was compared across the two groups
158 using a 2 (Food Type) x 2 (Stimulus Duration) x 2 (TFEQ_D group) ANOVA was conducted.
159 Effect sizes for both ANOVA's were reported as Cohen's d (d). The significant interaction
160 between disinhibition group and food type was explored using four planned comparisons of
161 the mean attentional bias for energy dense and low energy foods (within and between each
162 disinhibition group). A significant interaction was also found between stimulus duration and
163 food type. Four planned comparisons were conducted, these compared stimulus duration
164 (energy dense 100ms vs. 2000ms; low energy 100ms vs. 2000ms) and food type (energy
165 dense 100ms vs. low energy 100ms; energy dense 2000ms vs. low energy, 2000ms).
166 Bonferroni's correction was used to find the true critical p value for these eight planned
167 comparisons. This critical p value was $p < 0.006$. The extent to which attentional bias for food
168 cues reflected increased facilitated attention or delayed disengagement was explored using an
169 approach set out by Koster et al (2004).

171 Results

172
173 The demographics of the two groups are shown in Table 1. As expected, the groups differed
174 significantly in terms of their TFEQ_D scores [$p < 0.01$] and although the high disinhibition
175 group had higher BMI this was not significantly higher [$p = 0.51$]. There were no significant
176 between group differences in baseline hunger [$p > 0.05$]. Rated hunger did not change
177 significantly in either group between the start (time point one) and end of the study (time
178 point two) [$p > 0.05$]

179
180 Accuracy was significantly improved for trials which displayed stimuli pairs for 2000ms
181 compared to 100ms (Mean 99.6% compared to 96.5%) [$F(1, 42) = 240.71$ $p < 0.01$]. However
182 the type of stimulus which the probe followed (food or household item) had no significant
183 impact on detection accuracy [$F(1, 42) = 0.51$ $p = 0.47$]. The groups did not differ in terms of
184 task accuracy [$F(1, 42) = 0.06$ $p = 0.80$].

185
186 A 2 (Food Type) x 2 (Stimulus Duration) x 2 (TFEQ_D group) ANOVA was conducted (For
187 F values, effect size and mean bias scores for each group refer to Table 2). Analysis revealed
188 that both groups displayed attentional bias for food cues on trials where picture pairs
189 contained energy dense food items. There was no evidence of attentional bias for low energy
190 foods. There was an interaction found between disinhibition group and food type. Planned
191 comparisons indicated that both groups had a significantly higher attentional bias for trials
192 where picture pairs contained an energy dense stimulus compared to a low energy stimulus
193 (Low TFEQ_D; $t(22) = 3.69$ $p < 0.001$; High TFEQ_D $t(21) = 8.11$ $p < 0.001$). Although mean
194 attentional bias for energy dense foods was highest in the high TFEQ_D group planned
195 comparisons indicated no significant between group differences in attentional bias scores
196 based on either food type (Energy Dense $t(43) = 0.55$ $p = 0.58$; Low Energy $t(43) = 1.11$
197 $p = 0.27$).

198
199 An interaction was also found between stimulus duration and food type. Planned contrasts
200 conducted across the two time durations indicate that there were no significant differences in
201 bias scores when trials contained energy dense picture pairs [$p > 0.05$]. At the 100ms duration,
202 attentional bias was significantly higher for energy dense foods compared to low energy

204 types across 2000 ms trials ($t(44) = 7.03$ $p < 0.001$).

205

206 The extent to which attentional bias scores reflected facilitated attention to food cues or
207 delayed disengagement from food cues was explored using an approach set out by Koster et
208 al (2004). RTs (ms) for congruent and incongruent trials were compared to mean RTs from
209 neutral trials to indicate whether FPB reflected orientation or disengagement. If attentional
210 bias reflected facilitated attention to food cues this shown in quicker responses on congruent
211 trials (compared to neutral and congruent matched neutral). Whereas difficulty disengaging
212 from food cues would result in slower responses on incongruent trials (compared to neutral
213 and matched neutral). Evidence of facilitated attention was found only for energy-dense foods
214 in the high TFEQ_D group. Here participants were significantly faster at identifying probes
215 replacing congruent food items compared to neutral items [$t(21) = -2.289$ $p < 0.05$]. There was
216 no evidence of delayed disengagement in either group [$p > 0.05$].

217

218 **Discussion**

219

220 The present study is the first to examine if disinhibited eaters pay more attention to food cues.
221 The results suggested that trait disinhibition (as measured by the TFEQ_D subscale) is
222 associated with increased attentional bias for energy dense food cues. Although both groups
223 were significantly quicker at identifying probes replacing energy dense food cues compared
224 to neutral cues); mean attentional bias was highest in disinhibited eaters. The mean difference
225 in attentional bias scores between the high and low disinhibition group was 12.7 ms. Though
226 this difference is small it does support the prediction that disinhibited eaters opportunistic
227 eating pattern is associated with heightened attention to food cues. The visual dot probe data
228 documented attentional bias only on trials where the picture pairs contained energy dense
229 foods. This finding is consistent with previous research that also identified attention bias only
230 for palatable food items (Hepworth et al. 2010; Tapper et al. 2010). Disparity in task
231 performance on energy dense and low-energy trials was largest for the high disinhibition
232 group. This group typically displayed attentional bias for energy dense foods and directed
233 attention away from low-energy foods. This pattern of avoiding low energy foods and while
234 having biased processing of high energy foods is most commonly documented in patients
235 with disordered eating (Shafran, Lee, Cooper, Palmer & Fairburn (2007)).

236

237 From a methodological standpoint the findings from this study may be a consequence of the
238 type of stimuli chosen to represent 'low energy foods'. Many of these items were foods
239 which would not typically be consumed immediately or by themselves (i.e. shredded wheat
240 biscuit, plain rice). The energy dense stimuli set contained foods which were more
241 representative of foods that can be eaten "at that moment" (i.e. burgers, chips, crisps and
242 sweets). This is a limitation of classifying food into energy dense and low-energy groups, as
243 it is likely that the energy-dense foods are those which are easily obtainable and can be
244 consumed then and there. These foods may also be viewed as 'forbidden' by individuals who
245 are aware that they have difficulty regulating their eating behaviour These are all features that
246 are likely to have high salience for individuals whose appetite control is disinhibited by the
247 availability of palatable foods. In light of these comments, this interaction suggests that
248 opportunistic eaters allocate more attentional resources to cues that signal the availability of
249 'forbidden' or 'hedonic' foods.

250

251 In this study the visual dot probe task measured two components of attentional bias,
252 facilitated attention and delayed disengagement from cues. Evidence of facilitated attention

254 evidence of delayed disengagement. As facilitated attention is likely to act as a reminder of
255 the presence of food in the environment, this together with the elevated biases displayed by
256 the high TFEQ_D group suggests albeit tentatively that individuals with this eating trait are
257 more responsive to food cues. This data adds further support to the prediction that overeating
258 is driven by an individual's sensitivity to food cues. It can be inferred that the opportunistic
259 eating patterns of individuals who with high TFEQ_D scores places them at increased risk of
260 long-term weight gain. It is important to acknowledge that the BMI range in this sample was
261 restricted due to the sample size. There was also limited variation in the mean age of
262 participants; the majority of participants were in their early twenties and it is likely that if the
263 high TFEQ_D group exhibit a phenotype associated with weight gain, this may not be
264 expressed as obesity until later life. With this in mind it would be valuable to replicate this
265 experiment using an older sample with the inclusion of a follow up at 12 months; this would
266 allow us to ascertain if the higher biases seen in the disinhibited eaters are indeed reflected in
267 long-term weight gain.

268
269 To summarise this study is the first to illustrate that disinhibited eaters have a higher
270 attentional enhanced attention to food cues in the environment may underpin overeating. This
271 work further substantiates the proposition that paying enhanced attention to food cues in the
272 environment may underpin overeating. This data suggests that disinhibited eaters have an
273 increased risk of developing obesity, as disinhibition is associated with opportunistic eating
274 patterns but also increased attentional bias to food cues. This interaction needs to be
275 considered when developing successful interventions for weight management. There remains
276 scope to explore if attentional retraining can lead to a reduction in responsivity to food cues
277 in this non-clinical population.

278 279 **References**

- 280 Berridge, K. C., & Robinson, T. E. (1998). What is the role of dopamine in reward: Hedonic impact, reward
281 learning, or incentive salience? Elsevier B.V. doi:10.1016/S0165-0173(98)00019-8
- 282 Boschi, V., Margiotta, D. I. N and Falconi, P. D. C (2001) The Three-Factor Eating Questionnaire in the
283 Evaluation of eating behaviour in subjects seeking participation in a dietotherapy programme. *Ann Nutr Metab*
284 45(1) 72-77
- 285 Braet, C., & Crombez, G. (2003). Cognitive interference due to food cues in childhood obesity. *Journal of*
286 *Clinical Child & Adolescent Psychology*, 32(1), 32-39. doi:10.1207/S15374424JCCP3201_04
- 287 Brignell, C., Griffiths, T., Bradley, B. P., & Mogg, K. (2009). Attentional and approach biases for pictorial food
288 cues. influence of external eating. *Appetite*, 52(2), 299-306. doi:10.1016/j.appet.2008.10.007
- 289 Bryant, E. J., Caudwell, P., Hopkins, M. E., King, N. A., & Blundell, J. E. (2012). Psycho-markers of weight
290 loss. the roles of TFEQ disinhibition and restraint in exercise-induced weight management. *Appetite*, 58(1), 234-
291 241. doi:10.1016/j.appet.2011.09.006
- 292 Calitri, R., Pothos, E. M., Tapper, K., Brunstrom, J. M., & Rogers, P. J. (2010). Cognitive biases to healthy and
293 unhealthy food words predict change in BMI. *Obesity*, 18(12), 2282-2287. doi:10.1038/oby.2010.78
- 294 Castellanos, E. H., Charboneau, E., Dietrich, M. S., Park, S., Bradley, B. P., Mogg, K., & Cowan, R. L. (2009).
295 Obese adults have visual attention bias for food cue images: Evidence for altered reward system function.
296 *International Journal of Obesity*, 33(9), 1063-1073. doi:10.1038/ijo.2009.138
- 297 Contento, I. R., Zybert, P., & Williams, S. S. (2005). Relationship of cognitive restraint of eating and
298 disinhibition to the quality of food choices of latina women and their young children. *Preventive Medicine*,
299 40(3), 326-336. doi:10.1016/j.ypmed.2004.06.008
- 300 Cox, W. M., Klinger, E., & Fadardi, J. S. (2015). The motivational basis of cognitive determinants of addictive
301 behaviors. *Addictive Behaviors*, 44, 16.
- 302 Doolan, K. J., Breslin, G., Hanna, D., & Gallagher, A. M. (2015). Attentional bias to food-related visual cues: Is
303 there a role in obesity? *The Proceedings of the Nutrition Society*, 74(1), 37-45.
304 doi:10.1017/S002966511400144X
- 305 Field, M., & Cox, W. M. (2008). Attentional bias in addictive behaviors: A review of its development, causes,
306 and consequences. *Drug and Alcohol Dependence*, 97(1), 1-20. doi:10.1016/j.drugalcdep.2008.03.030

308 approaches. *Prog Neuropsychopharmacol Biol Psychiatry*, 27(1) 563-579

309 Graham, R., Hoover, A., Ceballos, N. A., & Komogortsev, O. (2011). Body mass index moderates gaze orienting

310 biases and pupil diameter to high and low calorie food images. *Appetite*, 56(3), 577-586.

311 doi:10.1016/j.appet.2011.01.029

312 Hays, N. P., & Roberts, S. B. (2008). Aspects of eating behaviors "disinhibition" and "restraint" are related to

313 weight gain and BMI in women. *Obesity*, 16(1), 52-58. doi:10.1038/oby.2007.12

314 Hepworth, R., Mogg, K., Brignell, C., & Bradley, B. P. (2010). Negative mood increases selective attention to

315 food cues and subjective appetite. *Appetite*, 54(1), 134-142. doi:10.1016/j.appet.2009.09.019

316 Herman, C. P., & Polivy, J. (2008). External cues in the control of food intake in humans: The sensory-

317 normative distinction. *Physiology & Behavior*, 94(5), 722-728. doi:10.1016/j.physbeh.2008.04.014

318 Hollitt, S., Kemps, E., Tiggemann, M., Smeets, E., & Mills, J. S. (2010). Components of attentional bias for foo

319 Robinson, T. E., & Berridge, K. C. (2001). Incentive-sensitization and addiction. *Addiction*, 96(1), 103-115.d

320 cues among restrained eaters. *Appetite*, 54(2), 309-313. doi:10.1016/j.appet.2009.12.005

321 Jastreboff, A. M., Sinha, R., Lacadie, C., Small, D. M., Sherwin, R. S., & Potenza, M. N. (2013). Neural

322 correlates of stress- and food cue-induced food craving in obesity: Association with insulin levels. *Diabetes*

323 *Care*, 36(2), 394-402. doi:10.2337/dc12-1112

324 Kemps, E., Tiggemann, M., & Hollitt, S. (2014). Biased attentional processing of food cues and modification in

325 obese individuals. *Health Psychology : Official Journal of the Division of Health Psychology, American*

326 *Psychological Association*, 33(11), 1391-1401. doi:10.1037/hea0000069

327 Koster, E. H. W., Crombez, G., Verschuere, B., & De Houwer, J. (2006). Attention to threat in anxiety-prone

328 individuals: Mechanisms underlying attentional bias. *Cognitive Therapy and Research*, 30(5), 635-643.

329 doi:10.1007/s10608-006-9042-9

330 Lähteenmäki, L., & Tuorila, H. (1995). Three-factor eating questionnaire and the use and liking of sweet and fat

331 among dieters. *Physiology & Behavior*, 57(1), 81-88. doi:10.1016/0031-9384(94)00210-V

332 Lattimore, P., Fisher, N., & Malinowski, P. (2011). A cross-sectional investigation of trait disinhibition and its

333 association with mindfulness and impulsivity. *Appetite*, 56(2), 241-248. doi:10.1016/j.appet.2010.12.007

334 Long, G Hinton, C. H & Gillespie, N. K (1994) Selective processing of food and body size words : Application

335 of the stroop test with obese restrained eaters, anorexics and normal. *International Journal of Eating Disorders*,

336 15(3)279-283

337 MacLeod, C., Mathews, A., & Tata, P. (1986). Attentional bias in emotional disorders. *Journal of Abnormal*

338 *Psychology*, 95(1), 15-20. doi:10.1037/0021-843X.95.1.15

339 Newman, E., O'Connor, D. B., & Conner, M. (2008). Attentional biases for food stimuli in external eaters:

340 Possible mechanism for stress-induced eating? *Appetite*, 51(2), 339-342. doi:10.1016/j.appet.2008.03.007

341 Nijs, I. M. T., Franken, I. H. A., & Muris, P. (2010). Food-related stroop interference in obese and normal-

342 weight individuals: Behavioral and electrophysiological indices. *Eating Behaviors*, 11(4), 258-265.

343 doi:10.1016/j.eatbeh.2010.07.002

344 Nijs, I. M. T., Franken, I. H. A., & Muris, P. (2010). Food-related stroop interference in obese and normal-

345 weight individuals: Behavioral and electrophysiological indices. *Eating Behaviors*, 11(4), 258-265.

346 doi:10.1016/j.eatbeh.2010.07.002

347 Provencher, V., Drapeau, V., Tremblay, A., Despre, J. and Lemieux (2003) Eating behaviors and indexes of body

348 composition in men and women from the québec family study. (2003). *Obesity Research*, 11(6), 783-792.

349 doi:10.1038/oby.2003.109

350 Shafran, R., Lee, M., Cooper, Z., Palmer, R. L., & Fairburn, C. G. (2007). Attentional bias in eating disorders.

351 *International Journal of Eating Disorders*, 40(4), 369-380. doi:10.1002/eat.20375

352 Smeets, E., Roefs, A., & Jansen, A. (2009). Experimentally induced chocolate craving leads to an attentional

353 bias in increased distraction but not in speeded detection. *Appetite*, 53(3), 370-375.

354 doi:10.1016/j.appet.2009.07.020

355 Stunkard, A. J and Messick, S(1985) The three factor eating questionnaire to measure dietary restraint,

356 disinhibition and hunger. *Journal of Psychosomatic Research*. 29(1) 71-83

357 Tapper, K., Pothos, E. M., & Lawrence, A. D. (2010). Feast your eyes: Hunger and trait reward drive predict

358 attentional bias for food cues. *Emotion (Washington, D.C.)*, 10(6), 949-954. doi:10.1037/a0020305

359 Tapper, K., Pothos, E. M., Fadardi, J. S., & Ziori, E. (2008). Restraint, disinhibition and food-related processing

360 bias. *Appetite*, 51(2), 335-338. doi:10.1016/j.appet.2008.03.006

361 Veenstra, E. M., de Jong, P. J., Koster, E. H. W., & Roefs, A. (2010). Attentional avoidance of high-fat food in

362 unsuccessful dieters. *Journal of Behavior Therapy and Experimental Psychiatry*, 41(3), 282-288.

363 doi:10.1016/j.jbtep.2010.02.006

364 Werthmann, J., Jansen, A., & Roefs, A. (2015). Worry or craving? A selective review of evidence for food-

365 related attention biases in obese individuals, eating-disorder patients, restrained eaters and healthy samples. *The*

366 *Proceedings of the Nutrition Society*, 74(2), 99-114. doi:10.1017/S0029665114001451

368 restraint factor from the three factor eating questionnaire. *Appetite*, 50(2), 469-476.
 369 doi:10.1016/j.appet.2007.10.002
 370 Yokum, S., Ng, J., & Stice, E. (2011). Attentional bias to food images associated with elevated weight and future
 371 weight gain: An fMRI study. *Obesity*, 19(9), 1775-1783. doi:10.1038/oby.2011.168
 372

Tables

Table 1: Demographics of the TFEQ_D Groups (Mean±SD)

	Low TFEQ_D (n=23)	High TFEQ_D (n=224)	<i>t</i>
Age	20.6(2.3)	20.3(0.9)	0.71
BMI	22.2(4.5)	25.0(4.7)	2.01
TFEQ_D	4.2(1.4)	9.4(2.5)	-8.55**
TFEQ_R	3.87(3.6)	6.1(3.8)	2.00
Hunger Time 1	52.6(1.09)	55.5(1.1)	0.87
Hunger Time 2	55.0(1.1)	53.9(1.2)	0.39

* $p < 0.05$ ** $p < 0.01$

Table 2: F value and effect size (Cohen's *d*)

	F	<i>p</i>	Effect size (<i>d</i>)
Food Type	70.71	0.00**	0.78
Stimulus Duration	1.63	0.21	
TFEQ_D	0.11	0.73	
TFEQ_D*Food Type	10.89	0.002**	0.44
Stimulus Duration* Food Type	7.13	0.01**	0.38

* $p < 0.05$ ** $p < 0.01$

Table 3: Mean±SD Bias Scores (ms) based on stimuli exposure and food type

Group	Stimulus Duration	Energy Dense	Low Energy	<i>t</i>
Low TFEQ_D	100ms	17.88±11.9	9.01±12.1	1.12
	2000ms	8.206±15.7	-20.08±15.9	4.59**
High TFEQ_D	100ms	19.80±8.7	-10.99±10.45	4.78**
	2000ms	20.95±15.7	-33.42±16.3	5.80**

* $p < 0.05$ ** $p < 0.01$

- Food cues command greater attentional resources than neutral items on the visual dot probe task.
- High disinhibition (as measured by the Three Factor Eating Questionnaire disinhibition subscale [TFEQ_D]) is predictive of increased attentional bias to high calorie stimuli.
- There is limited evidence for the existence of attentional bias for low calorie food items.
- Stimuli duration has limited impact on attentional bias to food cues.

ACCEPTED MANUSCRIPT