



Swansea University
Prifysgol Abertawe



Cronfa - Swansea University Open Access Repository

This is an author produced version of a paper published in :
PLOS ONE

Cronfa URL for this paper:

<http://cronfa.swan.ac.uk/Record/cronfa33140>

Paper:

Muema, D., Macharia, G., Olusola, B., Hassan, A., Fegan, G., Berkley, J., Urban, B. & Nduati, E. (2017). Proportions of circulating follicular helper T cells are reduced and correlate with memory B cells in HIV-infected children. *PLOS ONE*, 12(4), e0175570

<http://dx.doi.org/10.1371/journal.pone.0175570>

This article is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Authors are personally responsible for adhering to publisher restrictions or conditions. When uploading content they are required to comply with their publisher agreement and the SHERPA RoMEO database to judge whether or not it is copyright safe to add this version of the paper to this repository.

<http://www.swansea.ac.uk/iss/researchsupport/cronfa-support/>

RESEARCH ARTICLE

Proportions of circulating follicular helper T cells are reduced and correlate with memory B cells in HIV-infected children

Daniel M. Muema^{1#a*}, Gladys N. Macharia^{1#b}, Babatunde A. Olusola², Amin S. Hassan¹, Greg W. Fegan^{1,3#c}, James A. Berkley^{1,3}, Britta C. Urban^{1,4}, Eunice W. Nduati¹


1 Kenya Medical Research Institute-Wellcome Trust Research Programme, Centre for Geographic Medicine Research-Coast, Kilifi, Kenya, **2** Department of Virology, University of Ibadan, Ibadan, Nigeria, **3** Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom, **4** Department of Parasitology, Liverpool School of Tropical Medicine, Liverpool, United Kingdom

#a Current address: Africa Health Research Institute, Durban, South Africa.

#b Current address: Faculty of Medicine, Imperial College London, London, United Kingdom.

#c Current address: Swansea University Medical School, Swansea, United Kingdom.

* Daniel.muema@ahri.org



OPEN ACCESS

Citation: Muema DM, Macharia GN, Olusola BA, Hassan AS, Fegan GW, Berkley JA, et al. (2017) Proportions of circulating follicular helper T cells are reduced and correlate with memory B cells in HIV-infected children. PLoS ONE 12(4): e0175570. <https://doi.org/10.1371/journal.pone.0175570>

Editor: Derya Unutmaz, Jackson Laboratory, UNITED STATES

Received: September 12, 2016

Accepted: March 28, 2017

Published: April 26, 2017

Copyright: © 2017 Muema et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by Wellcome Trust UK [grant numbers 095068 (EWN), 089351 (ASH), 083579 (JAB) and 079082 (BCU)], <https://wellcome.ac.uk/>. BAO was supported by HIV Research Trust, <http://www.hivresearchtrust.org.uk/>.

Competing interests: The authors have declared that no competing interests exist.

Abstract

Introduction

HIV causes defects in memory B cells in children, but the mechanisms of those defects have not been fully elucidated. One possible mechanism is the lack of T-cell help to B cells during immune reactions. However, few studies have assessed the effect of HIV on follicular helper T cells (T_{FH} cells) in children.

Methods

In this study, follicular-homing CD4 T cells and memory B cells were assessed in HIV-infected children and compared with children from the community. CXCR5 and CD45RO were used as markers of follicular-homing T cells and memory T cells, respectively. Memory T_{FH} cells were identified as CD3⁺CD8⁻CD4⁺CXCR5⁺CD45RO⁺PD1⁺. Central memory T cells were identified based on CCR7 expression. Relationship between the proportions of follicular-homing CD4 T cells and memory B cells were determined in multivariable regression models.

Results

Highly viremic HIV-infected children had lower proportions of memory T_{FH} cells when compared with community control children. In multivariable analyses, high proportions of memory T_{FH} cells were associated with increased percentages of resting memory B cells after adjusting for other covariates.

Conclusion

The impact of HIV on follicular helper T cells could influence the accumulation of memory B cells in HIV-infected children.

Introduction

Even though depletion of CD4 T cells is the hallmark of HIV-induced immune dysfunction, the virus causes many other immunological abnormalities within the CD4 T-cell compartment. CD4 T cells from HIV patients are qualitatively defective, displaying features of aberrant immune activation as depicted by high levels of markers of activation [1]. Paradoxically, they also have impaired responsiveness to stimuli, an observation that has been attributed to the lymphocyte exhaustion that is characterized by up-regulation of inhibitory molecules [2, 3]. HIV is also associated with skewing of the subset-distribution of CD4 T cells. Viremic patients have fewer IL-2 producing central memory CD4 T cells [4]. In addition, active HIV viremia is associated with increased frequencies of follicular helper T cells (T_{FH} cells) in lymphoid tissues, suggesting increased T_{FH} activity [5].

HIV patients also make poor antibody and memory B-cell responses to routine vaccines and common infections [6–14]. The poor memory B-cell responses leave the patients, especially children, prone to repeated infections despite previous exposures and/or immunizations. Considering that one of the major functions of CD4 T cells is to provide help to B cells, the HIV-induced B-cell defects could be due to either depletion of CD4 T cells or HIV-induced qualitative defects in the CD4 T cells. Investigating the effect of HIV on T_{FH} cells, the subset of CD4 T cells that provides help to B cells in germinal centres, is necessary to comprehensively understand the mechanisms by which HIV impairs B-cell responses. Indeed, T_{FH} cells from the lymphoid tissues of HIV patients have been shown to be poor at helping the patients' B cells *in vitro*, an effect that has been attributed to increased PD1-PDL1 interaction [15].

Unfortunately, access to lymphoid tissues, the anatomical location of T_{FH} cells, entails performing invasive procedures and is logistically complicated. Attempts have therefore been made to identify counterparts of T_{FH} cells in peripheral circulation. Morita et al identified circulating T_{FH} on the basis of their CXCR5 expression, the marker for follicular homing, and showed that Th2 and Th17 skewing within this subset was associated with active disease in juvenile dermatomyositis [16]. Similarly, Pallikkuth et al used CXCR5 to identify circulating T_{FH} cells and associated their expansion with the magnitude of antibody response against the 2009 H1N1/09 vaccine in HIV patients [17]. Locci et al and Cohen et al described them as CXCR5+CXCR3-PD1+ and CXCR5+PD1+, respectively, and observed an association with eventual development of HIV cross-reactive antibodies [18, 19]. Boswell et al reported that the best B-cell helper capabilities were in the CXCR5^{high}CCR6^{high}PD1^{high} subset of CD4 T cells, though their frequencies did not correlate with development of cross-reactive neutralizing antibodies [20]. More recently, Schultz et al suggested that IL-21 secretion was the best marker for circulating memory T_{FH} cells [21].

In this study on HIV-infected children, the proportions of circulating T_{FH} cells and other follicular-homing CD4 T cells, and their relationship with memory B cells, were assessed. Considering that most previous studies in HIV used CXCR5 and PD1 to identify circulating T_{FH} cells, the same markers were used here.

Materials and methods

Study population

HIV-infected children aged 18 months to 10 years were recruited from the Comprehensive Care and Research Clinic at Kilifi County Hospital in 2012. The children were treated in accordance to the WHO guidelines at the time; those younger than 24 months were put on HAART regardless of their immunological and clinical profile, those between 25 months and 59 months were put on HAART if their CD4 percentages were below 25% or if they were in

WHO clinical stages 3 or 4 whereas children above 60 months of age were put on HAART if their CD4 percentages were below 20% or if they were in WHO clinical stages 3 or 4 [22]. Some children were recruited when it was their first time in the clinic (i.e. newly enrolled). The other children were recruited if they had received co-trimoxazole prophylaxis alone or in combination with HAART for at least six months. Viral loads were determined at the International Centre for Reproductive Health in Mombasa using a RT-qPCR test from Agence Nationale de Recherches sur le SIDA with a detection limit of 300 RNA copies/ml [23]. Community control children were recruited from the same community. They did not have fever, malaria parasitemia or any other active infection at the time of collecting the blood. Due to ethical constraints, the community control children were not directly assessed for HIV infection. However, the HIV prevalence among women in the coastal region was 6.1%, and, due to Prevention of Mother To Child Transmission (PMTCT) programs, the HIV prevalence was less than 1% among children in the community [24]. Written informed consent was obtained from the parents or guardians of all children. The study protocol was reviewed and approved by Kenya National Ethical Review Committee of Kenya Medical Research Institute (SSC numbers 1633 and 1131).

T-cell and B-cell subset determination in PBMCs by flow cytometry

A total of five milliliters of blood were obtained from each child. Plasma and the cellular fractions were separated by centrifugation. Peripheral blood mononuclear cells (PBMCs) were isolated from the cellular fraction by gradient centrifugation and stored in liquid nitrogen at 5 million PBMCs/ml in new born bovine serum containing 10% DMSO. On the day of the assay, PBMCs were thawed and washed in RPMI media containing 10% new born bovine serum, 100 units/ml penicillin, 0.1 mg/ml Streptomycin, 2 mM L-glutamine and 10 mM Hepes buffer.

The following antibodies were used: anti-IgD PE (IADB6), anti-CD19 ECD (J3-119), anti-CD3 ECD (UCHT1) and anti-CD27 PE-Cy5 (1A4CD27) from Beckman Coulter; anti-CD38-PE-Cy7 (HIT2) from eBiosciences; anti-CD8 FITC (DK25) from DAKO; anti-CXCR5 Brilliant Violet 421 (RF8B2), anti-CD4 PE-Cy7 (SK3), anti-CD21 APC (B-ly4) and anti-CD27 Brilliant Violet 421(M-T271) from BD Biosciences; anti-CCR7 APC (G043H7), anti-CD45RO APC-Cy7 (UCHL1) and anti-CD10 PE-Cy5 (HI10a) from Biolegend.

PBMCs were stained with various cocktails of the surface markers. At least 80,000 events were acquired on a nine-colour Cyan ADP flow cytometer (Beckman Coulter). Analyses and determination of different subsets were done in FlowJo software version 10 (TreeStar inc, Flowjo Africa Scheme).

Total follicular-homing CD4 T cells were identified based on the expression of CXCR5 i.e. CD3⁺CD8⁻CD4⁺CXCR5⁺. Since circulating follicular helper cells are expected to be antigen-experienced, the proportions of memory follicular-homing CD4 T cells were also determined based on the additional expression of CD45RO, the marker of memory in T cells i.e. CD3⁺CD8⁻CD4⁺CXCR5⁺CD45RO⁺. Since some studies have shown that the B-cell helper capabilities are enriched in the PD1⁺CXCR5⁺ memory CD4 T cells, the proportions of CD3⁺CD8⁻CD4⁺CXCR5⁺CD45RO⁺PD1⁺ were also determined, here referred as memory follicular helper CD4 T cells (memory T_{FH} cells). Within the memory compartments, central memory CD4 T cells were determined as CCR7⁺ cells, whereas effector memory CD4 T cells were determined as CCR7⁻ cells.

Total resting memory B cells were identified as CD19⁺CD27⁺CD21⁺CD10⁻CD38^{-/+} as described in previous studies [13, 25]. Further analyses were done to identify IgD⁺ (unswitched) and IgD⁻ (switched) resting memory B cells.

Statistical analyses

Active HIV viremia has been previously associated with B-cell defects [9, 13, 14, 26]. Therefore, the HIV-infected children were stratified into high viremia and low viremia groups based on a cut off of 5000 copies/ml, similar to our previous study and the WHO definition of treatment failure as at the time of enrollment of the children [13, 22].

Comparisons between different groups of children were done using the Mann-Whitney test (Wilcoxon rank sum test). Correlations were done using Spearman's rank-order correlation. Multivariable quantile regression analyses were done to assess independent relationships between various parameters. P values were considered significant if they were less than 0.05. STATA version 13 (STATA corporation) was used for all analyses.

Results

Characteristics of the study population

The population characteristics of the participants are shown in Table 1. A total of 52 children were recruited into the study of whom 36 were HIV-infected and 16 were community control children. Of the HIV-infected children, 15 had high viremia. Median viral loads were 4.6 (IQR, 4.3–5.0) log₁₀RNA copies/ml in the high viremia group and 1.9 (IQR, 0.0–3.1) log₁₀RNA copies/ml in the low viremia group (P<0.01). Nine of the high viremia children and 20 of the low viremia children were on HAART (P<0.01). CD4 T percentages data were available for 28 of the 36 HIV-infected children. Full blood counts data were available for 32 of the 36 HIV-infected children and all 16 community controls. The high viremia group had fewer monocytes when compared with the community controls (median 0.40 [IQR, 0.29–0.53] x 10³ cells/μL versus median 0.49 [IQR, 0.41–0.60] x 10³ cells/μL, P = 0.04). Interestingly, the low viremia group had fewer eosinophils when compared with both high viremia group (median 0.19 [IQR, 0.11–0.34] x 10³ cells/μL versus median 0.53 [IQR, 0.15–1.02] x 10³ cells/μL, P = 0.01) and community controls (median 0.19 [IQR, 0.11–0.34] x 10³ cells/μL versus median 0.49 [IQR, 0.30–0.68] x 10³ cells/μL, P<0.01). The high viremia group had lower hemoglobin levels

Table 1. Characteristics of the study population.

| | High viremia (HV) | Low viremia (LV) | Community controls (CC) | Normal range | P value | | |
|-------------------------------------|-------------------|------------------|-------------------------|--------------|-----------------|-------------|-----------------|
| | | | | | HV vs LV | HV vs CC | LV vs CC |
| N | 15 | 21 | 16 | N/A | | | |
| Age (years) | 5.9 (3.8–6.9) | 5.3 (2.6–8.0) | 5.8 (5.0–6.6) | N/A | 0.96 | 0.75 | 0.69 |
| % CD4+ T cells | 22.0 (16.7–34.4) | 30.0 (20.7–36.2) | N/D | | 0.31 | N/A | N/A |
| HIV log ₁₀ RNA copies/ml | 4.6 (4.3–5.0) | 1.9 (0.0–3.1) | N/A | N/A | <0.01 | N/A | N/A |
| % on HAART (n) | 60% (9) | 95% (20) | N/A | N/A | <0.01 | N/A | N/A |
| Lymphocytes (x 10 ³ /μL) | 3.8 (2.3–4.2) | 3.6 (3.1–4.5) | 3.4 (3.1–3.9) | 1.7–7.6 | 0.65 | 0.63 | 0.43 |
| Monocytes (x 10 ³ /μL) | 0.40 (0.29–0.53) | 0.52 (0.34–0.66) | 0.49 (0.41–0.60) | 0.3–1.5 | 0.22 | 0.04 | 0.83 |
| Neutrophils (x 10 ³ /μL) | 2.0 (1.3–2.5) | 2.6 (1.6–3.9) | 2.3 (2.1–3.6) | 1.2–5.5 | 0.23 | 0.13 | 0.79 |
| Eosinophils (x 10 ³ /μL) | 0.53 (0.15–1.02) | 0.19 (0.11–0.34) | 0.49 (0.30–0.68) | 0.1–1.3 | 0.01 | 0.93 | <0.01 |
| Platelets (x 10 ³ /μL) | 349 (293–449) | 365 (292–459) | 370 (278–444) | 159–564 | 0.98 | 0.93 | 0.76 |
| Hemoglobin (g/dL) | 10 (9–11) | 11 (10–11) | 11 (11–12) | 8.2–12.7 | 0.15 | 0.02 | 0.11 |

Values shown for age, % CD4+ T cells, HIV log₁₀ RNA copies/ml, lymphocytes, monocytes, neutrophils, eosinophils, platelets and hemoglobin are medians and interquartile ranges. Values under the normal range column refer to predetermined hematology reference values of normal children in Kilifi county. % CD4+ T cells refer to frequencies of CD4 T cells as percentage of total lymphocytes. Statistical tests used: Wilcoxon rank sum test was used for age, % CD4+ T cells, HIV log₁₀ RNA copies/ml, lymphocytes, monocytes, neutrophils, eosinophils, platelets and hemoglobin. Chi-squared test was used for % on HAART. N/D—Not determined. N/A—Not applicable.

<https://doi.org/10.1371/journal.pone.0175570.t001>

when compared with community controls (median 10 [IQR, 9–11] g/dL versus median 11 [IQR, 11–12] g/dL, $P = 0.02$).

HIV-infected children have reduced proportions of circulating follicular-homing CD4 T cells

Median proportions of total follicular-homing CD4 T cells ($CD3^+CD8^-CD4^+CXCR5^+$), memory follicular-homing CD4 T cells ($CD3^+CD8^-CD4^+CXCR5^+CD45RO^+$) and memory T_{FH} cells ($CD3^+CD8^-CD4^+CXCR5^+CD45RO^+PD1^+$) were lower in both HIV-infected groups when compared with community controls, except for memory T_{FH} cells in the low viremia group (Fig 1A–1C). The high viremia group also had significantly lower frequencies of central memory follicular-homing CD4 T cells and central memory T_{FH} cells when compared with the low viremia and community control groups (Fig 1D and 1F). In addition, the two HIV-infected groups had lower frequencies of effector memory follicular-homing CD4 T cells when compared with the community controls group (Fig 1E). In the memory T_{FH} compartment, the high viremia group, but not the low viremia group, had lower frequencies of effector memory T_{FH} cells when compared with the community controls group (Fig 1G).

Among all HIV-infected children, there was a direct correlation between total CD4 percentages and total follicular-homing CD4 T cells. A similar but non-significant trend was observed between total CD4 percentages and memory follicular-homing CD4 T cells ($P = 0.06$) but not memory T_{FH} cells (Fig 2A–2C). We also observed a trend towards direct correlation between total CD4 percentages and effector memory follicular-homing CD4 T cells ($P = 0.09$), but no relationship with other memory follicular-homing subsets (Fig 2D–2G).

Low proportions of memory T_{FH} cells are independent predictors of reduced proportions of resting memory B cells in HIV-infected children

To assess the relationship between memory follicular-homing CD4 T cells and memory B cells, the proportions of resting memory B cells were determined in accordance to previous




Fig 1. HIV infection is associated with low proportions of circulating follicular helper T cells (T_{FH} cells). Proportions of (A) total follicular-homing CD4 T cells ($CXCR5^+CD4^+$), (B) memory follicular-homing CD4 T cells ($CXCR5^+CD45RO^+CD4^+$), (C) memory T_{FH} cells ($CXCR5^+CD45RO^+PD1^+CD4^+$), (D) central memory follicular-homing CD4 T cells ($CXCR5^+CD45RO^+CCR7^+CD4^+$), (E) effector memory follicular-homing CD4 T cells ($CXCR5^+CD45RO^+CCR7^+CD4^+$), (F) central memory T_{FH} cells ($CXCR5^+CD45RO^+PD1^+CCR7^+CD4^+$) and (G) effector memory T_{FH} cells ($CXCR5^+CD45RO^+PD1^+CCR7^+CD4^+$) as percentages of total CD4 T cells. Each symbol represents a child. High viremia, ≥ 5000 RNA copies/ml; Low viremia, < 5000 RNA copies/ml. Horizontal lines and error bars represent median, 25th and 75th percentiles. Red symbols represent HIV-infected children who were not on HAART. Blue symbols represent HAART-treated children. Statistical test: Wilcoxon's rank sum test.

<https://doi.org/10.1371/journal.pone.0175570.g001>




Fig 2. In the HIV-infected cohort, proportions of total follicular-homing CD4 T cells are directly correlated with CD4 percentages. Correlation analyses between CD4 T cells (as percentage of total lymphocytes) and: (A) total follicular-homing CD4 T cells (CXCR5⁺CD4⁺), (B) memory follicular-homing CD4 T cells (CXCR5⁺CD45RO⁺CD4⁺), (C) memory T_{FH} cells (CXCR5⁺CD45RO⁺PD1⁺CD4⁺), (D) central memory follicular-homing CD4 T cells (CXCR5⁺CD45RO⁺CCR7⁺CD4⁺), (E) effector memory follicular-homing CD4 T cells (CXCR5⁺CD45RO⁺CCR7⁺CD4⁺), (F) central memory T_{FH} cells (CXCR5⁺CD45RO⁺PD1⁺CCR7⁺CD4⁺) and (G) effector memory T_{FH} cells (CXCR5⁺CD45RO⁺PD1⁺CCR7⁺CD4⁺). Circles represent highly viremic children and squares represent lowly viremic children. Red symbols represent HAART-naïve children and blue symbols represent HAART-treated children. Statistical test: Spearman's rank-order correlation.

<https://doi.org/10.1371/journal.pone.0175570.g002>

studies [13, 14, 25]. Similar to the previous reports, when compared with community controls, both high viremia and low viremia groups had lower proportions of total resting memory B cells (P<0.01 in both cases), IgD⁺ (Unswitched) resting memory B cells (P<0.01 and P = 0.01, respectively) and IgD⁻ (switched) resting memory B cells (P<0.01 and P = 0.01, respectively) (Fig 3A–3C) [9, 11, 13, 14, 27].

Some reports have suggested that CXCR5+PD1+ memory CD4 T cells are the main predictors of B-cell function with regard to the development of broadly neutralizing antibodies in HIV patients [18, 19]. Therefore, the association between memory T_{FH} cells (CD3⁺CD8⁻CD4⁺CXCR5⁺CD45RO⁺PD1⁺) and resting memory B cells was assessed in multivariable quantile regression analyses. When the independent predictions of resting memory B-cell subsets by HIV infection, age, memory T_{FH} cells, high viremia, interaction of age with high viremia and HAART were assessed, lower proportions of memory T_{FH} cells were predictors of lower proportions of total and IgD⁻ resting memory B cells. Being older was a predictor of higher proportions of IgD⁻ resting memory B cells (Table 2). The covariates HIV infection, age, high viremia, interaction of age with high viremia and HAART had been identified in a previous larger study [13].




Fig 3. HIV-infected children have reduced proportions of resting memory B cells. Proportions of (A) total resting memory B cells, (B) IgD⁺ resting memory B cells and (C) IgD⁻ resting memory B cells as percentages of total B cells. Each symbol represents a child. High viremia, ≥5000 RNA copies/ml; Low viremia, <5000 RNA copies/ml. Horizontal lines and error bars represent median, 25th and 75th percentiles. Red symbols represent HIV-infected children who were not on HAART. Blue symbols represent HAART-treated children. Statistical test: Wilcoxon's rank sum test.

<https://doi.org/10.1371/journal.pone.0175570.g003>

Table 2. Estimated change (beta coefficients) in frequencies of memory B-cell subsets with variations in age, HIV status, proportions of memory T_{FH} cells, viremia and HAART in multivariable regression.

| | Beta Coefficient (Standard Error) | P value |
|------------------------------|--------------------------------------|-----------------|
| Total resting memory | | |
| Age (Years) | 0.3 (0.3) | 0.30 |
| HIV infection | -3.9 (2.2) | 0.08 |
| Memory T _{FH} cells | 3.2 (1.1) | <0.01 |
| High viremia | -1.1 (3.0) | 0.72 |
| Age*high viremia | 0.0 (0.5) | 0.96 |
| HAART | 0.6 (1.9) | 0.76 |
| IgD+ resting memory | | |
| Age (Years) | -0.1 (0.1) | 0.46 |
| HIV infection | -1.4 (1.1) | 0.22 |
| Memory T _{FH} cells | 0.9 (0.5) | 0.13 |
| High viremia | -1.9 (1.6) | 0.23 |
| Age*high viremia | 0.3 (0.3) | 0.27 |
| HAART | 0.4 (1.0) | 0.68 |
| IgD- resting memory | | |
| Age (Years) | 0.5 (0.2) | 0.02 |
| HIV infection | -2.0 (1.5) | 0.20 |
| Memory T _{FH} cells | 2.0 (0.7) | 0.01 |
| High viremia | 0.8 (2.1) | 0.71 |
| Age*high viremia | -0.3 (0.3) | 0.39 |
| HAART | 0.2 (1.3) | 0.89 |

Each memory B-cell subset was independently predicted in a quantile regression model incorporating the reported variables. Age*high viremia is an interaction term for age and level of viremia. P values <0.05 were considered significant. Significant results are in bold text.

<https://doi.org/10.1371/journal.pone.0175570.t002>

In additional analyses on the predictive ability of other subsets within the follicular-homing CD4 T cells compartment, lower proportions of total follicular-homing CD4 T cells were predictors for lower proportions of total and IgD⁺ resting memory B cells (P<0.01 and P = 0.02, respectively). Similarly, low proportions of memory follicular-homing CD4 T cells were predictors for lower proportions of total and IgD⁺ resting memory B cells (P<0.01 and P = 0.02, respectively). Low proportions of central memory follicular-homing CD4 T cells were predictors for low proportions of total and IgD⁻ resting memory B cells (P<0.01 and P = 0.03, respectively) whereas low proportions of effector memory follicular-homing CD4 T cells were predictors for low proportions of total and IgD⁺ resting memory B cells (P = 0.02 and P<0.01, respectively). Low proportions of central memory T_{FH} cells were predictors for low proportions of total and IgD⁻ resting memory B cells (P = 0.02 in both cases) whereas low proportions of effector memory T_{FH} cells were predictors for low proportions of total and IgD⁺ resting memory B cells (P<0.01 in both cases) (Data not shown).

Discussion

Since antibodies have proven to play a critical role in vaccine mediated immunity against many infections, it would be important to understand the factors that determine the magnitude and quality of B-cell responses. Defects in the B-cell compartment leave HIV patients exposed to many opportunistic infections that would otherwise be controlled by a normal

antibody response [6, 7, 10, 12]. This problem is particularly significant in HIV-infected children, whose immature immune system develops in a background of HIV infection, leading to faster disease progression and probably more severe defects [28, 29]. Understanding the mechanisms by which HIV causes B-cell defects could inform interventions to improve antibody responses in HIV-infected children.

T_{FH} cells are the distinct CD4 T cell subset that provides help to B cells inside B-cell follicles in lymphoid tissues [30]. The recent identification of memory circulating counterparts of T_{FH} cells in blood has enabled studies that evaluate how HIV-induced defects on the T_{FH} compartment affect B-cell responses [17–19, 27].

In the current study, we show that the proportions of circulating follicular-homing CD4 T cells are lower in HIV-infected children when compared to community control children, similar to a previous report [27]. HIV preferentially replicates in T_{FH} cells in lymphoid tissues, probably leading to their preferential destruction, a phenomenon that could spill over into the circulating memory T_{FH} cells compartment [31]. Furthermore, B cells play a major role in reciprocally maintaining T_{FH} cells via co-stimulatory signals, and the detrimental effect of HIV on the B-cell compartment could indirectly impact the T_{FH} compartment [15]. Increased numbers of T_{FH} cells have been observed in lymphoid tissues of HIV patients, suggesting that depletion in peripheral blood could alternatively be due to, at least in part, increased trafficking into lymphoid organs [5, 32].

Notably, low proportions of memory T_{FH} cells in HIV-infected children were associated with reduced proportions of total and IgD⁺ (class-switched) resting memory B cells in multi-variable analyses after adjusting for other covariates. Previous reports have shown a similar link between frequencies of circulating memory T_{FH} cells and the quality of B-cell responses [17–19, 27]. T_{FH} cells in lymphoid tissues are important for the formation of germinal centers, the histological sites where long-lived plasma cells and memory B cells are generated during immune reactions. They also provide important contact and soluble signals that drive B-cell affinity maturation and class switch recombination, processes that determine the quality of the B-cell response [30]. The observed paucity of circulating memory T_{FH} cells implies that HIV-infected children have impaired T_{FH} activity resulting into the observed impaired generation of memory B cells. The reduced B-cell helper activity could be exacerbated by other qualitative defects in addition to these quantitative defects. For instance, memory T_{FH} cells from HIV-infected children could have altered differentiation patterns into less functional effector phenotypes. In this study, viremic HIV-infected children had fewer proportions of both central and effector memory T_{FH} cells, suggesting that the paucity of central memory T_{FH} cells was not due to differentiation into effector memory T_{FH} cells. However, T_{FH} cells from HIV-infected children have previously been shown to be defective at secreting IL-4, a cytokine that constitutes T-cell help to B cells and is required for maintenance of germinal centers, thus influencing the development of memory B cells [27, 30]. It is not known if the expression of surface molecules that mediate T-cell to B-cell interactions is altered in various subsets of follicular-homing CD4 T cells in HIV-infected children. Additional studies are required to assess such possibilities. Of note, the observed relationship between memory T_{FH} cells and memory B cells does not rule out the possibility of direct inhibition of B-cell responses by virion factors as suggested by some studies [33, 34].

Notably, the low viremia children also had reduced proportions of some of the subsets within follicular-homing CD4 T-cell compartment when compared with community controls. This observation implies that controlling viremia does not fully restore the memory T_{FH} compartment and that the HAART-treated children would need to rebuild the memory T_{FH} compartment through re-exposures to the various antigens. As such, re-immunization programmes could be beneficial to HAART-treated HIV-infected children.

Due to sample limitation, our study could not assess antigen-specific responses. It would be interesting to determine if similar results will be obtained with regard to antigen-specific memory T_{FH} cells and antigen specific B cells in HIV-infected children, and whether such responses predict the quality of antibody responses upon revaccination. Nevertheless, the data suggest that designing interventions that modulate T_{FH} cells could lead to better B-cell responses in HIV-infected children.

Supporting information

S1 Dataset.

(DTA)

S1 Fig. Representative flow-cytometry plots showing the gating strategy for identifying various subsets of follicular-homing CD4 T cells.

(PDF)

S2 Fig. Representative flow-cytometry plots showing the gating strategy for identifying resting memory B cells.

(PDF)

Acknowledgments

The authors are grateful to all the study participants, their guardians and the staff members of the Comprehensive Care and Research Clinic in Kilifi County Hospital. They are also grateful to Francis Ndungu and Jenniffer Maroa for the discussions. License for the FlowJo software was donated by Treestar Corp (Flowjo Africa scheme).

This paper is published with the permission of the director of Kenya Medical Research Institute.

Author Contributions

Conceptualization: DMM ASH JAB BCU EWN.

Formal analysis: DMM GWF EWN.

Funding acquisition: BCU EWN.

Investigation: DMM GNM ASH BAO.

Project administration: ASH.

Supervision: JAB BCU EWN.

Writing – original draft: DMM.

Writing – review & editing: DMM ASH JAB BCU EWN GNM BAO GWF.

References

1. Hazenberg MD, Otto SA, van Benthem BH, Roos MT, Coutinho RA, Lange JM, et al. Persistent immune activation in HIV-1 infection is associated with progression to AIDS. *Aids*. 2003; 17(13):1881–8. <https://doi.org/10.1097/01.aids.0000076311.76477.6e> PMID: 12960820
2. Day CL, Kaufmann DE, Kiepiela P, Brown JA, Moodley ES, Reddy S, et al. PD-1 expression on HIV-specific T cells is associated with T-cell exhaustion and disease progression. *Nature*. 2006; 443(7109):350–4. <https://doi.org/10.1038/nature05115> PMID: 16921384
3. Kaufmann DE, Kavanagh DG, Pereyra F, Zaunders JJ, Mackey EW, Miura T, et al. Upregulation of CTLA-4 by HIV-specific CD4+ T cells correlates with disease progression and defines a reversible

- immune dysfunction. *Nature immunology*. 2007; 8(11):1246–54. <https://doi.org/10.1038/ni1515> PMID: 17906628
4. Younes SA, Yassine-Diab B, Dumont AR, Boulassel MR, Grossman Z, Routy JP, et al. HIV-1 viremia prevents the establishment of interleukin 2-producing HIV-specific memory CD4+ T cells endowed with proliferative capacity. *The Journal of experimental medicine*. 2003; 198(12):1909–22. PubMed Central PMCID: PMC2194146. <https://doi.org/10.1084/jem.20031598> PMID: 14676302
 5. Lindqvist M, van Lunzen J, Soghoian DZ, Kuhl BD, Ranasinghe S, Kranias G, et al. Expansion of HIV-specific T follicular helper cells in chronic HIV infection. *The Journal of clinical investigation*. 2012; 122(9):3271–80. Epub 2012/08/28. PubMed Central PMCID: PMC3428098. <https://doi.org/10.1172/JCI64314> PMID: 22922259
 6. Bekker V, Scherpbier H, Pajkrt D, Jurriaans S, Zaaijer H, Kuijpers TW. Persistent humoral immune defect in highly active antiretroviral therapy-treated children with HIV-1 infection: loss of specific antibodies against attenuated vaccine strains and natural viral infection. *Pediatrics*. 2006; 118(2):e315–22. Epub 2006/07/19. <https://doi.org/10.1542/peds.2005-2616> PMID: 16847077
 7. Farquhar C, Wamalwa D, Selig S, John-Stewart G, Mabuka J, Majiwa M, et al. Immune responses to measles and tetanus vaccines among Kenyan human immunodeficiency virus type 1 (HIV-1)-infected children pre- and post-highly active antiretroviral therapy and revaccination. *The Pediatric infectious disease journal*. 2009; 28(4):295–9. Epub 2009/03/05. PubMed Central PMCID: PMC2779204. <https://doi.org/10.1097/INF.0b013e3181903ed3> PMID: 19258919
 8. Jacobsen MC, Thiebaut R, Fisher C, Sefe D, Clapson M, Klein N, et al. Pediatric human immunodeficiency virus infection and circulating IgD+ memory B cells. *The Journal of infectious diseases*. 2008; 198(4):481–5. Epub 2008/06/28. <https://doi.org/10.1086/590215> PMID: 18582200
 9. Moir S, Buckner CM, Ho J, Wang W, Chen J, Waldner AJ, et al. B cells in early and chronic HIV infection: evidence for preservation of immune function associated with early initiation of antiretroviral therapy. *Blood*. 2010; 116(25):5571–9. Epub 2010/09/15. PubMed Central PMCID: PMC3031405. <https://doi.org/10.1182/blood-2010-05-285528> PMID: 20837780
 10. Titanji K, De Milito A, Cagigi A, Thorstensson R, Grutzmeier S, Atlas A, et al. Loss of memory B cells impairs maintenance of long-term serologic memory during HIV-1 infection. *Blood*. 2006; 108(5):1580–7. Epub 2006/04/29. <https://doi.org/10.1182/blood-2005-11-013383> PMID: 16645169
 11. Iwajomo OH, Finn A, Moons P, Nkhata R, Sepako E, Ogunniyi AD, et al. Deteriorating pneumococcal-specific B-cell memory in minimally symptomatic African children with HIV infection. *The Journal of infectious diseases*. 2011; 204(4):534–43. Epub 2011/07/28. <https://doi.org/10.1093/infdis/jir316> PMID: 21791655
 12. Fernandes SJ, Shessarenko N, Souto FJ. Effects of vertical HIV infection on the persistence of anti-HBs after a schedule of three doses of recombinant hepatitis B vaccine. *Vaccine*. 2008; 26(8):1032–7. <https://doi.org/10.1016/j.vaccine.2007.12.020> PMID: 18242796
 13. Muema DM, Macharia GN, Hassan AS, Mwangi SM, Fegan GW, Berkley JA, et al. Control of Viremia Enables Acquisition of Resting Memory B Cells with Age and Normalization of Activated B Cell Phenotypes in HIV-Infected Children. *J Immunol*. 2015; 195(3):1082–91. <https://doi.org/10.4049/jimmunol.1500491> PMID: 26116511
 14. Pensiero S, Galli L, Nozza S, Ruffin N, Castagna A, Tambussi G, et al. B-cell subset alterations and correlated factors in HIV-1 infection. *Aids*. 2013; 27(8):1209–17. <https://doi.org/10.1097/QAD.0b013e32835edc47> PMID: 23343911
 15. Cubas RA, Mudd JC, Savoye AL, Perreau M, van Grevenynghe J, Metcalf T, et al. Inadequate T follicular cell help impairs B cell immunity during HIV infection. *Nature medicine*. 2013; 19(4):494–9. Epub 2013/03/12. <https://doi.org/10.1038/nm.3109> PMID: 23475201
 16. Morita R, Schmitt N, Bentebibel SE, Ranganathan R, Bourdery L, Zurawski G, et al. Human blood CXCR5(+)CD4(+) T cells are counterparts of T follicular cells and contain specific subsets that differentially support antibody secretion. *Immunity*. 2011; 34(1):108–21. PubMed Central PMCID: PMC3046815. <https://doi.org/10.1016/j.immuni.2010.12.012> PMID: 21215658
 17. Pallikkuth S, Parmigiani A, Silva SY, George VK, Fischl M, Pahwa R, et al. Impaired peripheral blood T-follicular helper cell function in HIV-infected nonresponders to the 2009 H1N1/09 vaccine. *Blood*. 2012; 120(5):985–93. PubMed Central PMCID: PMC3412336. <https://doi.org/10.1182/blood-2011-12-396648> PMID: 22692510
 18. Locci M, Havenar-Daughton C, Landais E, Wu J, Kroenke MA, Arlehamn CL, et al. Human circulating PD-(+)/CXCR3(-)/CXCR5(+) memory Tfh cells are highly functional and correlate with broadly neutralizing HIV antibody responses. *Immunity*. 2013; 39(4):758–69. Epub 2013/09/17. <https://doi.org/10.1016/j.immuni.2013.08.031> PMID: 24035365
 19. Cohen K, Altfeld M, Alter G, Stamatatos L. Early preservation of CXCR5+ PD-1+ helper T cells and B cell activation predict the breadth of neutralizing antibody responses in chronic HIV-1 infection. *Journal*

- of virology. 2014; 88(22):13310–21. PubMed Central PMCID: PMC4249103. <https://doi.org/10.1128/JVI.02186-14> PMID: 25210168
20. Boswell KL, Paris R, Boritz E, Ambrozak D, Yamamoto T, Darko S, et al. Loss of circulating CD4 T cells with B cell helper function during chronic HIV infection. *PLoS pathogens*. 2014; 10(1):e1003853. Epub 2014/02/06. PubMed Central PMCID: PMC3911819. <https://doi.org/10.1371/journal.ppat.1003853> PMID: 24497824
 21. Schultz BT, Teigler JE, Pissani F, Oster AF, Kranias G, Alter G, et al. Circulating HIV-Specific Interleukin-21(+)CD4(+) T Cells Represent Peripheral Tfh Cells with Antigen-Dependent Helper Functions. *Immunity*. 2016; 44(1):167–78. <https://doi.org/10.1016/j.immuni.2015.12.011> PMID: 26795249
 22. WHO. Antiretroviral therapy for HIV infection in infants and children: towards universal access. 2010; Accessed 10th September 2016:<http://www.who.int/hiv/pub/paediatric/infants2010/en/>.
 23. Rouet F, Chaix ML, Nerrienet E, Ngo-Giang-Huong N, Plantier JC, Burgard M, et al. Impact of HIV-1 genetic diversity on plasma HIV-1 RNA Quantification: usefulness of the Agence Nationale de Recherches sur le SIDA second-generation long terminal repeat-based real-time reverse transcriptase polymerase chain reaction test. *Journal of acquired immune deficiency syndromes*. 2007; 45(4):380–8. Epub 2007/05/01. <https://doi.org/10.1097/QAI.0b013e3180640cf5> PMID: 17468666
 24. National AIDS and STI Control Programme (NASCOP), Kenya. Kenya AIDS Indicator Survey 2012: Final Report. June 2014. Accessed 10th September 2016:<http://nacc.or.ke/wp-content/uploads/2015/10/KAIS-2.pdf>.
 25. Moir S, Fauci AS. B cells in HIV infection and disease. *Nature reviews Immunology*. 2009; 9(4):235–45. Epub 2009/03/26. PubMed Central PMCID: PMC2779527. <https://doi.org/10.1038/nri2524> PMID: 19319142
 26. Rethi B, Sammiceli S, Amu S, Pensieroso S, Hejdeman B, Schepis D, et al. Concerted effect of lymphopenia, viraemia and T-cell activation on Fas expression of peripheral B cells in HIV-1-infected patients. *Aids*. 2013; 27(2):155–62. Epub 2012/12/15. <https://doi.org/10.1097/QAD.0b013e32835b8c5e> PMID: 23238551
 27. Bekele Y, Amu S, Bobosha K, Lantto R, Nilsson A, Endale B, et al. Impaired Phenotype and Function of T Follicular Helper Cells in HIV-1-Infected Children Receiving ART. *Medicine*. 2015; 94(27):e1125. <https://doi.org/10.1097/MD.0000000000001125> PMID: 26166114
 28. Spira R, Lepage P, Msellati P, Van De Perre P, Leroy V, Simonon A, et al. Natural history of human immunodeficiency virus type 1 infection in children: a five-year prospective study in Rwanda. Mother-to-Child HIV-1 Transmission Study Group. *Pediatrics*. 1999; 104(5):e56. PMID: 10545582
 29. Newell ML, Coovadia H, Cortina-Borja M, Rollins N, Gaillard P, Dabis F. Mortality of infected and uninfected infants born to HIV-infected mothers in Africa: a pooled analysis. *Lancet*. 2004; 364(9441):1236–43. Epub 2004/10/07. [https://doi.org/10.1016/S0140-6736\(04\)17140-7](https://doi.org/10.1016/S0140-6736(04)17140-7) PMID: 15464184
 30. Crotty S. Follicular helper CD4 T cells (TFH). *Annual review of immunology*. 2011; 29:621–63. <https://doi.org/10.1146/annurev-immunol-031210-101400> PMID: 21314428
 31. Perreau M, Savoye AL, De Crignis E, Corpataux JM, Cubas R, Haddad EK, et al. Follicular helper T cells serve as the major CD4 T cell compartment for HIV-1 infection, replication, and production. *The Journal of experimental medicine*. 2013; 210(1):143–56. PubMed Central PMCID: PMC3549706. <https://doi.org/10.1084/jem.20121932> PMID: 23254284
 32. Petrovas C, Yamamoto T, Gerner MY, Boswell KL, Wloka K, Smith EC, et al. CD4 T follicular helper cell dynamics during SIV infection. *The Journal of clinical investigation*. 2012; 122(9):3281–94. PubMed Central PMCID: PMC3428091. <https://doi.org/10.1172/JCI63039> PMID: 22922258
 33. Jelacic K, Cimbri R, Nawaz F, Huang da W, Zheng X, Yang J, et al. The HIV-1 envelope protein gp120 impairs B cell proliferation by inducing TGF-beta1 production and FcRL4 expression. *Nature immunology*. 2013; 14(12):1256–65. PubMed Central PMCID: PMC3870659. <https://doi.org/10.1038/ni.2746> PMID: 24162774
 34. Qiao X, He B, Chiu A, Knowles DM, Chadburn A, Cerutti A. Human immunodeficiency virus 1 Nef suppresses CD40-dependent immunoglobulin class switching in bystander B cells. *Nature immunology*. 2006; 7(3):302–10. Epub 2006/01/24. <https://doi.org/10.1038/ni1302> PMID: 16429138