



Swansea University
Prifysgol Abertawe



Cronfa - Swansea University Open Access Repository

This is an author produced version of a paper published in :
Proceedings of the Royal Society B: Biological Sciences

Cronfa URL for this paper:
<http://cronfa.swan.ac.uk/Record/cronfa33667>

Paper:

Cornioley, T., Jenouvrier, S., Börger, L., Weimerskirch, H. & Ozgul, A. (2017). Fathers matter: male body mass affects life-history traits in a size-dimorphic seabird. *Proceedings of the Royal Society B: Biological Sciences*, 284 (1854), 20170397
<http://dx.doi.org/10.1098/rspb.2017.0397>

This article is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Authors are personally responsible for adhering to publisher restrictions or conditions. When uploading content they are required to comply with their publisher agreement and the SHERPA RoMEO database to judge whether or not it is copyright safe to add this version of the paper to this repository.

<http://www.swansea.ac.uk/iss/researchsupport/cronfa-support/>

Author Queries

Journal: Proceedings of the Royal Society B

Manuscript: RSPB20170397

As the publishing schedule is strict, please note that this might be the only stage at which you are able to thoroughly review your paper.

Please pay special attention to author names, affiliations and contact details, and figures, tables and their captions.

The corresponding author must provide an ORCID ID if they haven't done so already. If you or your co-authors have an ORCID ID please supply this with your corrections. More information about ORCID can be found at <http://orcid.org/>.

No changes can be made after publication.

SQ1 Please confirm that this paper is intended to be Open Access. The charge for Open Access should be paid before publication. If you have not yet received an email requesting payment please let us know when returning your corrections.

SQ2 Your supplementary material will be published online alongside your article and on rs.figshare.com exactly as the file(s) are provided. Therefore, please could you either confirm that your supplementary material is correct, or – if you have any changes to make to these files – email these along with your proof corrections to the journal inbox. Your ESM files are listed here for your convenience:

SI1Mass_stand_res.2.0docx.pdf

SI2Adult_sur.2.0docx.pdf

Q1 Please provide the volume number for ref. [58].

Q2 Reference [58] is provided in the list but not cited in the text. Please supply citation details or delete the reference from the reference list.



Cite this article: Cornioley T, Jenouvrier S, Börger L, Weimerskirch H, Ozgul A. 2017

Fathers matter: male body mass affects life-history traits in a size dimorphic seabird.

Proc. R. Soc. B 20170397.

<http://dx.doi.org/10.1098/rspb.2017.0397>

Received: 24 February 2017

Accepted: 28 March 2017

Subject Category:

Ecology

Subject Areas:

ecology

Keywords:

wandering albatross, bi-parental care, sexual dimorphism, survival, reproduction

Author for correspondence:

Tina Cornioley

e-mail: tina.cornioley@ieu.uzh.ch

Electronic supplementary material is available online at rs.figshare.com.

Fathers matter: male body mass affects life-history traits in a size dimorphic seabird

Tina Cornioley¹, Stéphanie Jenouvrier^{2,3}, Luca Börger⁴, Henri Weimerskirch³ and Arpat Ozgul¹

¹Department of Evolutionary Biology and Environmental Studies, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland

²Woods Hole Oceanographic Institution, Mailstop 50, Woods Hole, MA 02543, USA

³Centre d'Etudes Biologiques de Chizé, Station d'Ecologie de Chizé-La Rochelle, CNRS UMR 7372, 79360 Villiers-en-Bois, France

⁴Department of Biosciences, College of Science, Swansea University, Singleton Park, Swansea, Wales SA2 8PP, UK

TC, 0000-0002-9882-6235

One of the predicted consequences of climate change is a shift in body mass distributions within animal populations. Yet, body mass, an important component of the physiological state of an organism, can affect key life-history traits and consequently population dynamics. Over the past decades, the wandering albatross—a pelagic seabird providing bi-parental care with marked sexual size dimorphism—has exhibited an increase in average body mass and breeding success in parallel to experiencing increasing wind speeds. To assess the impact of these changes, we examined how body mass affects five key life-history traits at the individual level: adult survival, breeding probability, breeding success, chick mass and juvenile survival. We found that male mass impacted all traits examined except breeding probability, whereas female mass affected none. Adult male survival increased with increasing mass. Increasing adult male mass increased breeding success and mass of sons but not of daughters. Juvenile male survival increased with their chick mass. These results suggest that a higher investment in sons by fathers can increase their inclusive fitness, which is not the case for daughters. Our study highlights sex-specific differences in the effect of body mass on the life history of a monogamous species with bi-parental care.

1. Introduction

Body mass is an important component of an organism's body condition because it reflects the intrinsic amount of energy reserve available to survive and breed [1]. Yet body mass of numerous species have shifted with climate change [2–4]. Therefore, investigating how body mass shapes life-history traits is critical to ultimately understand the impact of environmental changes on individual fitness and population dynamics [5–7].

Body mass and other indices of body condition positively impact adult survival in several species of mammals [5,8–10] and in some long-lived birds such as geese and seabirds [11–13], although sometimes the relationship is only apparent in extreme climatic conditions [14]. This is because long-lived species are expected to allocate their resources primarily to survival rather than reproduction [15]. For example, in poor environmental conditions, yellow-nosed albatrosses (*Thalassarche chlororhynchos*) were only slightly lighter, experienced no change in survival but substantially decreased their provisioning rate to their offspring [16]. Similarly, southern fulmars (*Fulmarus glacialisoides*) exposed

64 to poor environmental conditions had lower breeding suc-
65 cess, while survival of adults did not vary significantly [17].
66 This allocation strategy may cause body mass to impact
67 reproductive performances but not survival. Thus, a relation-
68 ship between body mass and survival is not trivial in
69 long-lived species.

70 In species with sexual size dimorphism, life-history traits
71 can be expected to respond differently to male and female
72 body mass. The larger sex may suffer costs associated with
73 growth and maintenance of a large size [18–20]. Thus, its
74 life-history traits can be expected to be more sensitive to
75 body mass than the smaller sex which has lower energetic
76 requirements. Furthermore, sex-specific relationships can
77 arise from segregation in diet choice and space utilization
78 determined by size ([20,21], but see [22]). For example,
79 sexual dimorphism could reduce food competition between
80 sexes in predatory birds as males and females hunt prey
81 related to their respective size [23].

82 Arguably, the life-history traits most expected to respond
83 differently to male and female body mass are those linked to
84 reproductive performance, even in species providing bi-par-
85 ental care. This is perhaps straightforward in species where
86 each parent performs a role suited to its size. In predatory
87 birds, where the female is typically the larger sex, large size
88 confers higher reproductive performance to females, whereas
89 it is disadvantageous for males [23–25].

90 Less obvious are cases of species performing similar par-
91 ental role, parental contribution and/or parental investment.
92 Parental role here refers to the tasks performed by each
93 parent during reproduction. It is more or less equally parti-
94 tioned depending on whether the tasks are performed by
95 one or both parents. Parental investment refers to the cost
96 to the parent, i.e. the amount of energy spent by a parent
97 during reproduction. Parental contribution refers to the
98 benefit to the offspring from each parent, i.e. the amount of
99 energy it received from each parent. In the wandering
100 albatross (*Diomedea exulans*), partners incubate and feed
101 their single chick in turns. Thus, parental roles are roughly
102 equally partitioned. However, when rearing chicks,
103 fathers contributes in absolute terms more energy to the
104 chick [26], resulting in unequal parental contribution. There-
105 fore, reproductive performance may respond predominantly
106 to the body condition of the parent contributing the most
107 to reproduction, even in species with bi-parental care.

108 Here, we examined the effects of adult and chick mass on
109 a suite of life-history traits in the wandering albatross, a long-
110 lived species providing bi-parental care with sexual size
111 dimorphism. Given that this species has increased in mass
112 over the past decades in parallel to an increase in wind
113 speed [27], it is valuable to investigate the effect of mass on
114 individual life-history traits to understand the consequences
115 of environmental changes. Using data from a long-term indi-
116 vidual-based study, we studied five life-history traits: adult
117 survival, breeding probability, breeding success, chick mass
118 and juvenile survival.

119 We predicted that adult mass should increase breed-
120 ing probability, breeding success and chick mass, but not
121 adult survival and that chick mass should increase juvenile
122 survival, with possible sex-specific differences. Despite
123 equal role partitioning, given the larger contribution of
124 males during reproduction [28], a more pronounced effect
125 of adult male body mass on reproductive performance
126 was expected.

2. Material and methods

(a) Data collection

The wandering albatross population of Possession Island (Crozet Archipelago), ranging from 250 to 500 breeding pairs, was monitored annually since 1966. The colony is visited multiple times during the breeding season to assess status of individuals [29]. They were classified as chicks (from hatching to departure from the colony), immature (have never bred), non-breeder (have breeding experience but are not breeding during the year of observation), pre-breeder (have breeding experience and are at the colony at the beginning of the breeding season), incubating or chick rearing. Individuals were weighed in 18 different years between 1988 and 2013 in varying status and in some cases, tarsus length was also measured. For a complete description of the field methods, see [30].

(i) Standardization of mass

We chose to focus on body mass rather than another index of body condition because (i) measures of structural size, such as tarsus length, were available for only a subset of the individuals with known mass, and (ii) for birds, body mass is a reliable index of body condition [1]. This was also the case in the wandering albatross; a common body condition index, the residuals of a generalized additive model of mass as a function of tarsus length, age and sex of incubating individuals, was strongly correlated to body mass (Pearson correlation coefficient = 0.94 for males and 0.88 for females). The same test for chicks showed the same pattern (Pearson correlation coefficient = 0.94 for males and 0.95 for females). The very strong correlation between the two metrics suggests that they contain the same information.

Mass is a plastic trait that can vary across time, status and age [30]. Thus, mass measurements collected at different point in time are not directly comparable. Prior to performing life-history trait analyses, we built two mass standardization functions, one for adults and another one for juveniles. These enabled us to standardize mass at a reference status relevant for each life-history trait. The standardization functions are presented in detail in the electronic supplementary material, S1.

(b) Life-history traits and mass

(i) Adult and juvenile survival

Mass measurements were available for 662 adults with breeding experience aged between 6 and 30 years, i.e. before the onset of senescence in survival [30,31]; excluding the minor set of senescent individuals and focusing on the majority of the breeding population led to more robust models. Mass measurements were standardized within the year they were taken to the non-breeding mass using the adult mass standardization function as mortality is assumed to occur during the non-breeding period.

The survival analyses for both adults (model set S) and juveniles (model set SJ) were performed in a capture–mark–recapture framework in the program E-Surge to account for imperfect detection [32]. Mass was considered to affect only adult survival from the year of measurement to the next. As the wandering albatross is a facultative biennial breeder, successive capture events are not independent, thus we implemented an Immediate Trap Effect on Capture model [33] (more details in the electronic supplementary material, S2).

For the juvenile survival analysis, we distinguished juveniles—individuals aged 1–2 years old never visiting the colony, from the other immatures—individuals at least 3 years old that have not yet bred but can potentially visit the colony. Individuals were grouped into three age classes (1–2, 3–8 and greater than 8), following [34]. We examined the effect of chick mass on survival for age 1–2—referred to as

juvenile survival—and allowed age-class-specific and sex-specific observation probabilities.

For each survival analysis, we compared 10 models: (1, 2) without mass and (3–10) eight with mass. The effect of mass was either (3, 4) distinct for each sex, (5, 6) the same for both sexes or applied only to (7, 8) males or only to (9, 10) females. Models were constructed with (1, 3, 5, 7, 9) or without (2, 4, 6, 8, 10) a different intercept for each sex on the transition of interest. All continuous variables were standardized. The other transitions were sex- and mass-independent. Models were ranked based on their AICc. Only the model with the lowest AICc and the models within $2\Delta\text{AICc}$ that contained no more variables than the previous one were retained following the principle of parsimony [35]. The same model selection procedure has been applied to the other analyses.

(ii) Breeding probability

Here, breeding probability is conditioned upon the presence on the island as only present individuals can be weighed. This distinction is important because most individuals skipping breeding do not visit the island [36]. Mass measurements were available for 356 breeders and 55 non-breeders with breeding experience and were standardized within the year they were taken to the non-breeding mass because non-breeders do not have a breeding mass the years they do not reproduce.

Three breeding probability analyses were performed. The first examined the effect of mass (model set PB). We fitted six generalized linear mixed effect (GLME) models with a binomial distribution and a logit link function including year as a random effect with the package lme4 (v. 1.1–7, [37]). Only one randomly selected entry per individual was kept as the full model did not converge with individual ID as a random effect due to the low number of repeated individuals. To account for the effect of age, we built on the model retained by Pardo *et al.* [31]; consequently, our simplest model included a linear age effect in interaction with sex. We compared this model with a series of age-and-mass models. These models included mass with or without an interaction between mass and sex, or with or without an interaction effect between age and mass, or with or without a three-way interaction between mass, age and sex. Comparison of all possible combination of variables was performed with the ‘dredge’ command from the MuMin package (v. 1.13.4, [38]) on models fitted with maximum likelihood. The estimates reported are from the selected models fitted with restricted maximum likelihood.

Because sons are more demanding to rear [26], we additionally tested for a potential impact of fledging a son versus a daughter on male breeding probability the year immediately after the reproductive event (model set PB_{y+1}) and the following year (model set PB_{y+2}). We fitted GLME models with a binomial distribution and a logit function with ID as a random effect. In addition to sex of the previous chick, we considered age of father as a potential explanatory variable.

(iii) Breeding success

In 116 cases, mass measurements of both partners of known age and breeding success were available (model set BS). Often, the mass of only one partner was known. Thus, we also examined exclusively the effect of female mass (274 cases; model set BSF) and exclusively of male mass (283 cases; model set BSM). Mass was standardized to the first day of incubation.

We fitted GLME models with a binomial distribution and a logit link function with year and ID as random effects. The minimum model included linear and quadratic effects of age in interaction with sex [31]. We compared the minimum model with age-and-mass models. These models included a combination of an interaction between mass and sex, an interaction between mass and age or a three-way interaction among mass, age and sex.

(iv) Chick mass

Mass measurements of at least one of the parents and of the chick were taken the year of hatching for 89 sexed chicks. Because mass of both parents was not always available, the same statistical approach was performed on three datasets: with both parents (model set MaP, 40 cases), with only the mother (model set MaF, 62 cases) and with only the father (model set MaM, 67 cases). Linear mixed effect models with a Gaussian distribution and year as a random effect were fitted with the lme4 package (v. 1.1–7, [37]). The models compared included a combination of sex of the chick, mass of one or both parents and the interaction between the variables.

3. Results

Male adult survival was positively influenced by mass, whereas this was not the case for females (model S1 in tables 1 and 2 and figure 1a). The set of models investigating a potential effect of mass on breeding probability found that breeding probability was mass-independent in both sexes as mass was never retained during model selection (model PB1 in table 1). Breeding probability was also independent of the sex of the previous chick, both 1 and 2 years after the last breeding event (models $\text{PB1}_{(y+1)}$ and $\text{PB1}_{(y+2)}$ in table 1). A positive effect of mass on breeding success was detected in the model fitted to the dataset including only male mass. Conversely, no effect of mass was detected in the models fitted on the datasets including mass of both partners or only female mass (models BS1, BSF1 and BSM1 in tables 1 and 2 and figure 1c).

We did not detect any effect of parent mass on chick mass in models fitted to datasets including both parents or only the mother (models MaP,1 MaF1 and MaF2 in tables 1 and 2), whereas father’s mass was retained in the top model when using the dataset including only father’s mass (model MaM1 in table 2). Father’s mass was found to positively impact chick mass in sons, but not in daughters (model MaM1 in table 2 and figure 1d). Chick mass had a positive effect on the annual male juvenile survival but not for females (model SJ1 in table 2 and figure 1b).

4. Discussion

This study provides a comprehensive overview of the effect of body mass on life-history traits of a sexually dimorphic long-lived species providing bi-parental care. We found a clear difference between sexes: male mass enhanced performance in four life-history traits, namely adult survival, breeding success, chick mass and juvenile survival, whereas female mass impacted none. Sex-specific relationships between mass and life-history traits were expected, given the different ecology of the two sexes. Indeed, males and females forage in different oceanic sectors [40], follow a different relationship between wind, mass and foraging statistics such as flight speed and maximum distance from the colony [41], and contribute differently to reproduction [26,28].

(a) Effect of mass on survival

We found that body mass of adult males enhances survival in one of the longest-lived bird, irrespective of environmental conditions. This is not trivial as long-lived species are expected to allocate resources to survival before reproduction [15], which can potentially buffer an influence of mass on

Table 1. Retained models for adult survival (model set S), breeding probability (model sets PB, $PB_{(y+1)}$ and $PB_{(y+2)}$), breeding success (model sets BS, BSF and BSM), chick mass (model sets Ma, MaF and MaM) and juvenile survival (SJ). Model set PB tested for the effect of mass on breeding probability, model set $PB_{(y+1)}$ for the effect of previous-year offspring sex, model set $PB_{(y+2)}$ for the effect of sex of offspring two year previous. Model sets BS, BSF and BSM are performed on different datasets: including both partners, only females and only males, respectively. The same applied to model sets testing for an effect of adult mass on chick mass; including both partners (Ma), only females (MaF) and only males (MaM). If not specified, mass refers to adult mass. AICc values are estimated from models fitted by ML. Within each model set, only the model with the lowest AICc ($\Delta AICc = 0$) and the models with fewer parameters within $2\Delta AICc$ from the model with $\Delta AICc = 0$ are reported. The $\Delta AICc$ is the difference in AICc from the model with the lowest AICc within the model set. For the survival analyses, d.f. refers to the number of estimable parameters, it adds up to more than the number of explanatory variables listed because it also includes the parameters for the probability of recapture (not listed because they were identical in all models). For the other models, d.f. refers to the degree of freedom for linear mixed effect models based on the inner–outer rules [39].

model set	model	resp. var	expl. var	d.f.	AICc	$\Delta AICc$
adult survival						
S	S1	<i>Sur.</i> ~	Sex + Mass (M)	8	8940.68	0
	S2	<i>Sur.</i> ~	Mass (M)	7	8941.65	0.97
breeding probability						
PB	PB1	<i>Prob. Bre</i> ~	Sex + Age + Sex : Age	5	260.87	0
$PB_{(y+1)}$	$PB1_{(y+1)}$	<i>Prob. Bre</i> $_{(y+1)}$ ~	Age	3	502.8	0
$PB_{(y+2)}$	$PB1_{(y+2)}$	<i>Prob. Bre</i> $_{(y+2)}$ ~	Age	3	2244.5	0
breeding success						
BS	BS1	<i>Bre. Suc.</i> ~	Age (F) + Age ² (F) + Age (M) + Age ² (M)	7	122.83	0
BSF	BSF1	<i>Bre. Suc.</i> ~	Age + Age ²	5	321.59	0
BSM	BSM1	<i>Bre. Suc.</i> ~	Age + Age ² + Mass	6	294.38	0
	BSM2	<i>Bre. Suc.</i> ~	Age + Age ²	5	294.76	0.38
chick mass						
Ma	Ma1	Chick mass ~	Sex chick	4	652.56	0
MaF	MaF1	Chick mass ~	Sex chick + Age	5	997.51	0
	MaF2	Chick mass ~	Sex chick	4	997.83	0.33
MaM	MaM1	Chick mass ~	Sex chick + Mass + Mass : Sex chick	6	1074.59	0
	MaM2	Chick mass ~	Sex chick	4	1075.01	0.42
Juvenile survival						
SJ	SJ1	Chick <i>sur.</i> ~	Age(1–2) + Age(3–8) + Age(>8) + Mass Chicks(M)	8	2981.00	0

survival even in extremely poor environmental conditions. For instance, there is no such relationship in the blue petrel (*Halobaena caerulea*) [42] and the lack of variation in survival in the southern fulmar, even in poor environmental conditions, suggest likewise [17]. Yet the case of the wandering albatross is not unique. Shorter-lived seabirds, the little auk (*Alle alle*), in extreme environmental conditions [14], and the black-legged kittiwake (*Rissa tridactyla*) [11,43], also had enhanced survival when in good body condition.

Unlike male survival, female adult survival of wandering albatrosses was not impacted by body mass. Such difference may reflect the higher energetic requirements of the largest sex to maintain its large size [19,20] or may be related to sex-specific energetically demanding activities. For example, in the greater white-fronted goose (*Anser albifrons*), only female survival varies with body mass, which is explained by the higher energetic requirement of this sex, the only one that incubates [13]. Similarly, male wandering albatrosses experience male-specific energetically demanding periods in their life cycle, in particular during the pre-breeding period when they are guarding the nest for one month without feeding, and at the end of the breeding season when they provision for the chick for longer than females [44].

The contrasted relationships between mass and survival among sexes may have important consequences for the population dynamics. Indeed, they can affect population growth rate directly and indirectly through the population structure, operational sex ratio (OSR) and mating process [45]. For example, male emperor penguin survival is determined by environmental conditions while female survival is not [46], which impacts OSR. The sensitivity of the population growth rate to female survival is negative [45] as the indirect effects through OSR can sometimes overwhelm the direct effects. For the wandering albatross, the consequences on population growth rate of the contrasted adult survivals remain an open question.

(b) Mass-independent breeding probability

As in the blue petrel [42], breeding probability of adult wandering albatrosses was not influenced by body mass. This is intriguing because body mass determines the first breeding attempt: among individuals of the same age, immature birds tend to be lighter than breeders [30]. It is possible that the absence of an effect of body mass on breeding probability is an artefact. Breeding probability examined here is

Table 2. Estimates on the logit scale of the coefficients of the fixed effects included in the models in table 1 fitted by REML. *1st trans.* refers to the yearly survival from the first year after mass measurement and *subs. trans.* to all the subsequent yearly survival.

model	resp. var	expl. var	estimate	s.e.
S1	<i>Sur.</i> ~	Intercept for 1st <i>trans.</i> (F)	3.106	0.072
		Intercept for 1st <i>trans.</i> (M)	2.759	0.321
		<i>Subs. trans.</i>	1.791	0.434
		Mass (M)	1.078	0.583
S2	<i>Sur.</i> ~	Intercept for 1st <i>trans.</i> (F)	3.105	0.072
		Intercept for 1st <i>trans.</i> (M)	3.681	0.666
		<i>Subs. trans.</i>	1.913	0.421
PB1	<i>Prob. Bre.</i> ~	Intercept (F)	2.700	0.694
		Sex (M)	-0.147	0.364
		Age	-0.133	0.298
		Sex (M):Age	0.151	0.398
		<i>Marg. R²: 0.02, Cond. R²: 0.60, Rand eff.: Year</i>		
BS1	<i>Bre. Suc.</i> ~	Intercept	2.338	0.603
		Age (M)	0.529	0.498
		Age ² (M)	-0.871	0.324
		Age (F)	-0.420	0.582
		Age ² (F)	0.076	0.250
		<i>Marg. R²: 0.25, Cond. R²: 0.34, Rand eff.: Year + ID</i>		
PB1 _(y+1)	<i>Prob. Bre.</i> _(y+1) ~	Intercept	-8.15	0.68
		Age	0.47	0.19
		<i>Marg. R²: 0.00, Cond. R²: 0.00, Rand eff.: ID</i>		
PB1 _(y+2)	<i>Prob. Bre.</i> _(y+2) ~	Intercept	0.86	0.05
		Age	-0.08	0.05
		<i>Marg. R²: 0.00, Cond. R²: 0.03, Rand eff.: ID</i>		
SF1	<i>Bre. Suc.</i> (F)~	Intercept	1.253	0.233
		Age	0.172	0.196
		Age ²	-0.129	0.109
		<i>Marg. R²: 0.01, Cond. R²: 0.06, Rand eff.: Year + ID</i>		
BSM1	<i>Bre. Suc.</i> (M)~	Intercept	1.645	0.247
		Age	-0.182	0.216
		Age ²	-0.312	0.129
		Mass	0.307	0.193
		<i>Marg. R²: 0.13, Cond. R²: 0.16, Rand eff.: Year + ID</i>		
BSM2	<i>Bre. Suc.</i> (M)~	Intercept	1.740	0.266
		Age	-0.044	0.199
		Age ²	-0.378	0.126
		<i>Marg. R²: 0.12, Cond. R²: 0.17, Rand eff.: Year + ID</i>		
MaP1	Mass chick ~	Intercept (F)	10997	157.7
		Sex chick (M)	1324.2	249.3
		<i>Marg. R²: 0.42, Cond. R²: 0.42, Rand eff.: Year</i>		
MaF1	Mass chick ~	Intercept (F)	11368.11	230.25
		Sex chick (M)	1246.83	178.66
		Age	142.64	87.25
		<i>Marg. R²: 0.40, Cond. R²: 0.55, Rand eff.: Year</i>		

(Continued.)

Table 2. (Continued.)

model	resp. var	expl. var	estimate	s.e.
MaF2	Mass chick ~	Intercept (F)	11352.34	228.6
		Sex chick (M)	1236.66	181.12
		<i>Marg. R²: 0.38, Cond. R²: 0.52, Rand eff.: Year</i>		
MaM1	Mass chick ~	Intercept (F)	10958.32	172.28
		Sex chick (M)	1512.05	167.86
		Mass	-42.47	100.7
		Mass:Sex chick	367.5	177.19
<i>Marg. R²: 0.54, Cond. R²: 0.59, Rand eff.: Year</i>				
MaM2	Mass chick ~	Intercept (F)	10970.98	174.71
		Sex chick (M)	1505.86	171.79
		<i>Marg. R²: 0.51, Cond. R²: 0.57, Rand eff.: Year</i>		
SJ1	Juv sur. ~	Intercept for age (1–2)	0.741	0.122
		Intercept for age (3–8)	7.791	40.82
		Intercept for age (>8)	2.737	0.417
		Chick mass (M)	0.259	0.130

conditional upon return to the colony. The non-breeders sampled may have returned because they were in good enough condition to breed but some failed to breed for reasons independent of their own condition, for instance, because their partners were absent. Birds in poor condition may simply have not returned to the colony and remain at sea. We cannot test for this assumption, as individuals that are absent from the colony cannot be weighed.

(c) Effect of mass on reproductive performance

The reproductive performance varied with the body mass of the father, the parent contributing the most to reproduction [28], but not with that of the mother. The dependence on exclusively one parent can be linked to how parents resolve conflict over offspring care. In a bi-parental system with perfect information, parents should incompletely compensate for a reduction in their partner contribution when increasing contribution yields decelerating benefits to the offspring at a non-decelerating cost to the parent [47]. That is, when the costs to the parent are high and the benefit to the offspring low, the parent has little incentive to increase investment. When from the same level of cost, the contribution to reproductive effort (i.e. the benefit to the offspring) differs between the parents, the parent with the lowest contribution for the same cost can compensate less than the other parent. This is possibly the case in the wandering albatross.

The absence of a relationship between wandering albatross female mass and either breeding success or chick mass probably does not reflect a lack of investment of females. Rather, it indicates that no matter the condition of the female, her absolute contribution to reproduction cannot compensate for a reduced contribution of the male. Females likely bear costs proportional to that of males as suggested by the paralleled mass variation of males and females during chick rearing [28]. Furthermore, in proportion to their mass, females provide larger meals—even though

they visit the nest less often [26]. Yet, for the same mass, the wing loading of females, which have smaller wing surface, changes more than that of a male, suggesting that the energetic cost of females per meal is larger as higher wing loading increases the cost of taking-off and landing [48]. Indeed, in the wandering albatross population of South Georgia, there is evidence that female cost of chick rearing is higher than that of males [49]. Similarly, in the northern giant petrel, a seabird with a sexual size dimorphism even more pronounced than the wandering albatross, females foraging costs are much higher than that of males [22]. It thus is probable that females, the smaller sex, do not invest less, but that their contribution has less impact.

Unequal contribution to reproduction is not uncommon in bi-parental systems. Other seabirds show higher food provisioning from one parent, usually the largest sex [50,51], but also in the absence of sexual size dimorphism [52,53]. In species like the wandering albatross where there is an asymmetry in contribution between parents, the reproductive performance can be expected to depend more on the condition of parent contributing the most.

(d) Fathers invest more in sons

Numerous previous studies which experimentally manipulated body conditions of the parents (or one parent) and/or chick need, reported parents' (or one parent's) ability to adjust food provisioning to both chick and parents conditions (e.g. [45,47]). However, here we found evidence that this is the case in an unmanipulated system. By examining jointly the effect of adult body mass on chick body mass and the effect of chick body mass on juvenile survival, we could detect indirect benefit to the adult of investing more in a chick to improve its post-fledging survival.

The strategy of wandering albatross fathers is consistent with the current theory stating that long-lived seabirds should adjust their reproductive performance to both their




Figure 1. Effect of mass on life-history traits. Predicted annual male and female survival for (a) adult based on model S1, for (b) juveniles based on model SJ1. The annual breeding success of males as predicted by the BSM1 is plotted in (c) as function of age and mass. The colour gradient represents the effect of mass, confidence interval in this case have been omitted to improve readability. Peak chick mass of offspring in relation to the mass of father is depicted in (d). The dots are the observed values and lines the predicted values. The grey ribbons are the 95% CIs (except for c) and have been calculated with the delta method for (a,b).

own body conditions and the need of their offspring [54]. We found evidence of higher investment of fathers in sons, but not in daughters with increasing father mass. This trend corroborates well with the higher cost of rearing sons observed in many species [55–57]. In the wandering albatross, both parents provide more food to sons than to daughters, sons have faster growth rates, reach higher fledgling mass [26] and parents adjust meal size to male chick needs [28]. The higher energetic need of males is often explained by the higher vulnerability of male juveniles to food shortage compared with females [18], a hypothesis likely applicable to the wandering albatross as male juveniles have a lower survival ([34], this study).

Our results suggest that, in wandering albatrosses, investing further in sons to respond to their higher energy requirement is an efficient strategy for fathers to increase their inclusive fitness because it increased sons' juvenile survival, and it came at no cost to future breeding performance. Indeed, heavy male chicks had a higher juvenile survival than lighter ones, whereas mass had no effect on female juvenile survival. Furthermore, the breeding probability of fathers that produced sons 1 or 2 years later was not lower than that of fathers that produced daughters. This suggests that producing a son has no carry-over effect on male breeding performance in the short term. Not only do fathers adjust investment in their son to their son's conditions [28], but also to their own. Fathers seem to share with their sons only the resources that they can spare within a given year.

5. Conclusion

Our results showed that among long-lived species, a relationship between body mass and survival can be present independently of environmental conditions. The survival of only one of the two sexes can be affected by body mass but not the other, probably due to sex-specific energy requirements [19]. Furthermore, we showed that life-history traits related to reproduction of species with sexual size dimorphism with biparental care can vary exclusively with body condition of the sex contributing the most to reproduction. Theory on resolution of parental conflict over care provides a framework to explain the emergence of such a pattern [54]: when for the same cost to the parents, the benefits derived by the offspring produced from the care of each parents are unequal, then the parents contributing the less may not be able to compensate.

When some life-history traits are mass-dependent, variation in mass distribution is expected to have consequences at population level. Given that body mass has been reported to have changed over the past years with wind speed in both sexes [27], understanding how a climate-driven change in this trait will affect population dynamics will prove to be crucial for the conservation of this species.

Ethics. The Ethics Committee of IPEV and Comité de l'Environnement Polaire approved the field procedures.

Data accessibility. Data will be made available on Dryad.

Authors' contributions. All contributed to conception and design of the study. H.W. collected field data. T.C. and S.J. analysed the data.

T.C. wrote the manuscript in collaboration with all. All authors gave final approval for publication.

Competing interests. We declare we have no competing interests.

Funding. This study is supported by the Swiss National Science Foundation project grant #31003A_146445 and the ERC Starting Grant no. 337785 to A.O. and is a contribution to the Program EAR-LYLIFE funded by an ERC Advanced Grant under the European Community's Seven Framework Program FP7/2007-2013 (ERC-2012-ADG_20120314 to H.W.). The long-term demographic study at

Crozet was supported by the French Polar Institute IPEV (programme no. 109 to H.W.). S.J. acknowledges support from NSF project no.1246407.

Acknowledgements. We thank the field workers since 1966 at Possession Island and Dominique Besson and Karine Delord for invaluable help with data management. We are grateful to Koen van Benthem and Marine Desprez for their help with the analysis and stimulating discussions. We are grateful to two anonymous reviewers for their insightful comments on an earlier version of the manuscript.

References

- Labocha MK, Hayes JP. 2012 Morphometric indices of body condition in birds: a review. *J. Ornithol.* **153**, 1–22. (doi:10.1007/s10336-011-0706-1)
- Millien V, Kathleen Lyons S, Olson L, Smith FA, Wilson AB, Yom-Tov Y. 2006 Ecotypic variation in the context of global climate change: revisiting the rules. *Ecol. Lett.* **9**, 853–869. (doi:10.1111/j.1461-0248.2006.00928.x)
- Yom-Tov Y, Geffen E. 2006 Geographic variation in body size: the effects of ambient temperature and precipitation. *Oecologia* **148**, 213–218. (doi:10.1007/s00442-006-0364-9)
- Gardner JL, Peters A, Kearney MR, Joseph L, Heinsohn R. 2011 Declining body size: a third universal response to warming? *Trends Ecol. Evol.* **26**, 285–291. (doi:10.1016/j.tree.2011.03.005)
- Ozgul A, Childs DZ, Oli MK, Armitage KB, Blumstein DT, Olson LE, Tuljapurkar S, Coulson T. 2010 Coupled dynamics of body mass and population growth in response to environmental change. *Nature* **466**, 482–485. (doi:10.1038/nature09210)
- Plard F, Gaillard J-MM, Coulson T, Delorme D, Warnant C, Michallet J, Tuljapurka S, Krishnakumar S, Bonenfant C. 2015 Quantifying the influence of measured and unmeasured individual differences on demography. *J. Anim. Ecol.* **84**, 1434–1445. (doi:10.1111/1365-2656.12393)
- Vindenes Y, Langangen Ø. 2015 Individual heterogeneity in life histories and eco-evolutionary dynamics. *Ecol. Lett.* **18**, 417–432. (doi:10.1111/ele.12421)
- Bérubé CH, Festa-Bianchet M, Jorgenson JT. 1999 Individual differences, longevity, and reproductive senescence in bighorn ewes. *Ecology* **80**, 2555–2565. (doi:10.1890/0012-9658(1999)080[2555:IDLARS]2.0.CO;2)
- Clutton-Brock TH, Stevenson IR, Marrow P, MacColl AD, Houston AI, McNamara JM. 1996 Population fluctuations, reproductive costs and life-history tactics in female Soay sheep. *J. Anim. Ecol.* **65**, 675–689. (doi:10.2307/5667)
- Ozgul A, Bateman AW, English S, Coulson T, Clutton-Brock TH. 2014 Linking body mass and group dynamics in an obligate cooperative breeder. *J. Anim. Ecol.* **83**, 1357–1366. (doi:10.1111/1365-2656.12239)
- Oro D, Furness RW. 2002 Influences of food availability and predation on survival of kittiwakes. *Ecology* **83**, 2516–2528. (doi:10.1890/0012-9658(2002)083[2516:IOFAAP]2.0.CO;2)
- Haramis GM, Nichols JD, Pollock KH, Hines JE. 1986 The relationship between body mass and survival of wintering canvasback. *Auk* **103**, 506–514.
- Schmutz JA, Ely CR. 1999 Survival of greater white-fronted geese: effects of year, season, sex and body condition. *J. Wildl. Manage.* **63**, 1239–1249. (doi:10.2307/3802841)
- Harding AMA *et al.* 2011 Adverse foraging conditions may impact body mass and survival of a high Arctic seabird. *Oecologia* **167**, 49–59. (doi:10.1007/s00442-011-1971-7)
- Williams GC. 1966 Natural selection, the costs of reproduction, and a refinement of Lack's principle. *Am. Nat.* **100**, 687–690. (doi:10.1086/282461)
- Weimerskirch H, Zimmermann L, Prince PA. 2001 Influence of environmental variability on breeding effort in a long-lived seabird, the yellow-nosed albatross. *Behav. Ecol.* **12**, 22–30. (doi:10.1093/oxfordjournals.beheco.a000374)
- Jenouvrier S, Péron C, Weimerskirch H. 2015 Extreme climate events and individual heterogeneity shape lifehistory traits and population dynamics. *Ecol. Monogr.* **85**, 605–623. (doi:10.1890/14-1834.1)
- Clutton-Brock TH, Albon SD, Guinness FE. 1985 Parental investment and sex differences in juvenile mortality in birds and mammals. *Nature* **313**, 131–133. (doi:10.1038/313131a0)
- Isaac JL. 2005 Potential causes and life-history consequences of sexual size dimorphism in mammals. *Mamm. Rev.* **35**, 101–115. (doi:10.1111/j.1365-2907.2005.00045.x)
- Clutton-Brock TH, Guinness FE, Albon SD. 1982 *Red deer, behavior and ecology of two sexes*. Chicago, IL: The University of Chicago Press.
- Conradt L, Clutton-Brock T, Guinness F. 2000 Sex differences in weather sensitivity can cause habitat segregation: red deer as an example. *Anim. Behav.* **59**, 1049–1060. (doi:10.1006/anbe.2000.1409)
- González-Solís J, Croxall JP, Wood AG. 2000 Sexual dimorphism and sexual segregation in foraging strategies of northern giant petrels, *Macronectes halli*, during incubation. *Oikos* **90**, 390–398. (doi:10.1034/j.1600-0706.2000.900220.x)
- Andersson AM, Norberg RÅ. 1981 Evolution of reversed sexual size dimorphism and role partitioning among predatory birds, with a size scaling of flight performance. *Biol. J. Linn. Soc.* **15**, 105–130. (doi:10.1111/j.1095-8312.1981.tb00752.x)
- McDonald PG, Olsen PD, Cockburn A. 2005 Selection on body size in a raptor with pronounced reversed sexual size dimorphism: are bigger females better? *Behav. Ecol.* **16**, 48–56. (doi:10.1093/beheco/arh118)
- Sunde P, Bolstad MS, Moller JD. 2003 Reversed sexual dimorphism in tawny owls, *Strix aluco*, correlates with duty division in breeding effort. *Oikos* **101**, 265–278. (doi:10.1034/j.1600-0706.2003.12203.x)
- Weimerskirch H, Barbraud C, Lys P. 2000 Sex differences in parental investment and chick growth in wandering albatrosses: fitness consequences. *Ecology* **81**, 309–318. (doi:10.1890/0012-9658(2000)081[0309:SDIPIA]2.0.CO;2)
- Weimerskirch H, Louzao M, de Grissac S, Delord K. 2012 Changes in wind pattern alter albatross distribution and life-history traits. *Science* **335**, 211–214. (doi:10.1126/science.1212070)
- Weimerskirch H, Lys P. 2000 Seasonal changes in the provisioning behaviour and mass of male and female wandering albatrosses in relation to the growth of their chick. *Polar Biol.* **23**, 733–744. (doi:10.1007/s003000000144)
- Weimerskirch H, Brothers N, Jouventin P. 1997 Population dynamics of wandering albatross *Diomedea exulans* and Amsterdam albatross *D. amsterdamensis* in the Indian Ocean and their relationships with long-line fisheries: conservation implications. *Biol. Conserv.* **79**, 257–270. (doi:10.1016/S0006-3207(96)00084-5)
- Weimerskirch H. 1992 Reproductive effort in long-lived birds: age-specific patterns of condition, reproduction and survival in the wandering albatross. *Oikos* **64**, 464–473. (doi:10.2307/3545162)
- Pardo D, Barbraud C, Weimerskirch H. 2014 What shall I do now? State-dependent variations of life-history traits with aging in wandering albatrosses. *Ecol. Evol.* **4**, 474–487. (doi:10.1002/ece3.882)
- Choquet R, Rouan L, Pradel R. 2009 Program E-Surge: a software application for fitting multievent models. In *Modeling demographic processes in marked populations* (eds DL Thomson, EG Cooch, MJ Conroy), pp. 845–865. Boston, MA: Springer US.
- Pradel R, Sanz-Aguilar A. 2012 Modeling trap-awareness and related phenomena in capture-recapture studies. *PLoS ONE* **7**, 1–4. (doi:10.1371/annotation/e240f425-0375-4c32-b0a7-85fa586d0f40)

- 505 34. Fay R, Weimerskirch H, Delord K, Barbraud C. 2015
506 Population density and climate shape early-life
507 survival and recruitment in a long-lived pelagic
508 seabird. *J. Anim. Ecol.* **84**, 1423–1433. (doi:10.
509 1111/1365-2656.12390)
- 510 35. Burnham KP, Anderson DR. 2002 *Model selection*
511 *and multimodel inference: a practical information-*
512 *theoretic approach*. New York, NY: Springer.
- 513 36. Barbraud C, Weimerskirch H. 2012 Estimating
514 survival and reproduction in a quasi-biennially
515 breeding seabird with uncertain and unobservable
516 states. *J. Ornithol.* **152**, 605–615. (doi:10.1007/
517 s10336-011-0686-1)
- 518 37. Bates D, Mächler M, Bolker BM, Walker SC. 2015
519 Fitting linear mixed-effects models using lme4.
520 *J. Stat. Softw.* **67**, 1–48. (doi:10.18637/jss.v067.i01)
- 521 38. Barton K. 2015 MuMIn: multi-model inference.
522 R package version 1.13.4. Version 1, 18.
- 523 39. Pinheiro J, Bates D, DebRoy S, Sarkar D, orpR Core
524 Team. 2016 nlme: Linear and nonlinear mixed
525 effects models.
- 526 40. Weimerskirch H, Chérel Y, Delord K, Jaeger A, Patrick
527 SC, Riotte-Lambert L. 2014 Lifetime foraging
528 patterns of the wandering albatross: life on the
529 move! *J. Exp. Mar. Biol. Ecol.* **450**, 68–78. (doi:10.
530 1016/j.jembe.2013.10.021)
- 531 41. Cornioley T, Börger L, Ozgul A, Weimerskirch H.
532 2016 Impact of changing wind conditions on
533 foraging and incubation success in male and female
534 wandering albatrosses. *J. Anim. Ecol.* **85**,
535 1318–1327. (doi:10.1111/1365-2656.12552)
- 536 42. Barbraud C, Weimerskirch H. 2005 Environmental
537 conditions and breeding experience affect costs of
538 reproduction in Blue Petrels. *Ecology* **86**, 682–692.
539 (doi:10.1890/04-0075)
- 540 43. Kitaysky AS, Piatt JF, Hatch SA, Kitaiskaia EV, Benowitz-
541 Fredericks ZM, Shultz MT, Wingfield JC. 2010 Food
542 availability and population processes: severity of
543 nutritional stress during reproduction predicts survival
544 of long-lived seabirds. *Funct. Ecol.* **24**, 625–637.
545 (doi:10.1111/j.1365-2435.2009.01679.x)
- 546 44. Tickell WLN. 1968 The biology of the great
547 albatrosses, *Diomedea exulans* and *Diomedea*
548 *epomophora*. *Antarct. Res. Ser.* **12**, 1–55.
- 549 45. Jenouvrier S, Caswell H, Barbraud C, Weimerskirch
550 H. 2010 Mating behavior, population growth,
551 and the operational sex ratio: a periodic two-sex
552 model approach. *Am. Nat.* **175**, 739–752. (doi:10.
553 1086/652436)
- 554 46. Jenouvrier S, Barbraud C, Weimerskirch H. 2005
555 Long-term contrasted responses to climate of two
556 antarctic seabird species. *Ecology* **86**, 2889–2903.
557 (doi:10.1890/05-0514)
- 558 47. Johnstone RA, Hinde CA. 2006 Negotiation over
559 offspring care—how should parents respond to
560 each other's efforts? *Behav. Ecol.* **17**, 818–827.
561 (doi:10.1093/beheco/arl009)
- 562 48. Shaffer S, Weimerskirch H, Costa D. 2001 Functional
563 significance of sexual dimorphism in wandering
564 albatrosses, *Diomedea exulans*. *Funct. Ecol.* **15**,
565 203–210. (doi:10.1046/j.1365-2435.2001.00514.x)
- 566 49. Berrow SD, Croxall JP. 2001 Provisioning rate and
567 attendance patterns of wandering albatrosses at
568 Bird Island, South Georgia. *Condor* **103**, 230.
569 (doi:10.1650/0010-5422(2001)103[0230:PRAAPO]
570 2.0.CO;2)
- 571 50. Huin N, Prince PA, Briggs DR. 2000 Chick
572 provisioning rates and growth in black-browed
573 Albatross *Diomedea melanophris* and grey-headed
574 albatross *D. chrysostoma* at Bird Island, South
575 Georgia. *Ibis (Lond. 1859)* **142**, 550–565. (doi:10.
576 1111/j.1474-919X.2000.tb04456.x)
- 577 51. Weimerskirch H, Le Corre M, Gadenne H, Pinaud D,
578 Kato A, Ropert-Coudert Y, Bost C-A. 2009
579 Relationship between reversed sexual dimorphism,
580 breeding investment and foraging ecology in a
581 pelagic seabird, the masked booby. *Oecologia* **161**,
582 637–649. (doi:10.1007/s00442-009-1397-7)
- 583 52. Gray CM, Hamer KC. 2001 Food-provisioning
584 behaviour of male and female Manx shearwaters,
585 *Puffinus puffinus*. *Anim. Behav.* **62**, 117–121.
586 (doi:10.1006/anbe.2001.1717)
- 587 53. Peck DR, Congdon BC. 2006 Sex-specific chick
588 provisioning and diving behaviour in the wedge-
589 tailed shearwater *Puffinus pacificus*. *J. Avian Biol.*
590 **37**, 245–251. (doi:10.1111/j.2006.0908-8857.
591 03558.x)
- 592 54. Erikstad KE, Fauchald P, Tveraa T, Steen H. 1998 On
593 the cost of reproduction in long-lived birds: the
594 influence of environmental variability. *Ecology* **79**,
595 1781–1788. (doi:10.1890/0012-9658(1998)079
596 [1781:OTCORI]2.0.CO;2)
- 597 55. Berube CH, Festa-Bianchet M, Jorgenson JT. 1996
598 Reproductive costs of sons and daughters in Rocky
599 Mountain bighorn sheep. *Behav. Ecol.* **7**, 60–68.
600 (doi:10.1093/beheco/7.1.60)
- 601 56. Bercovitch FB, Widdig A, Nürnberg P. 2000 Maternal
602 investment in rhesus macaques (*Macaca mulatta*):
603 reproductive costs and consequences of raising sons.
604 *Behav. Ecol. Sociobiol.* **48**, 1–11. (doi:10.1007/
605 s002650000204)
- 606 57. Harding AMA, Kitaysky AS, Hall ME, Welcker J,
607 Karnovsky NJ, Talbot SL, Hamer KC, Grémillet D.
608 2009 Flexibility in the parental effort of an Arctic-
609 breeding seabird. *Funct. Ecol.* **23**, 348–358. (doi:10.
610 1111/j.1365-2435.2008.01488.x)
- 611 58. Baylis AMM, Orben R, Costa DP, Arnould JPY,
612 Staniland IJ. 2016 Sexual segregation in habitat use
613 is smaller than expected in a highly dimorphic
614 marine predator, the southern sea lion. *Mar. Ecol.*
615 *Prog. Ser.* 201–211. (doi:10.3354/meps11759)