



Swansea University
Prifysgol Abertawe



Cronfa - Swansea University Open Access Repository

This is an author produced version of a paper published in:
Journal of Mathematical Physics

Cronfa URL for this paper:
<http://cronfa.swan.ac.uk/Record/cronfa34436>

Paper:

Boejko, M., Lytvynov, E. & Wysoczanski, J. (2017). Fock representations of Q-deformed commutation relations.
Journal of Mathematical Physics, 58(7), 073501
<http://dx.doi.org/10.1063/1.4991671>

This item is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Copies of full text items may be used or reproduced in any format or medium, without prior permission for personal research or study, educational or non-commercial purposes only. The copyright for any work remains with the original author unless otherwise specified. The full-text must not be sold in any format or medium without the formal permission of the copyright holder.

Permission for multiple reproductions should be obtained from the original author.

Authors are personally responsible for adhering to copyright and publisher restrictions when uploading content to the repository.

<http://www.swansea.ac.uk/iss/researchsupport/cronfa-support/>

Fock representations of Q-deformed commutation relations

Marek Bożejko, Eugene Lytvynov, and Janusz Wysoczański

Citation: *Journal of Mathematical Physics* **58**, 073501 (2017); doi: 10.1063/1.4991671

View online: <http://dx.doi.org/10.1063/1.4991671>

View Table of Contents: <http://aip.scitation.org/toc/jmp/58/7>

Published by the *American Institute of Physics*

Articles you may be interested in

[KP hierarchy for the cyclic quiver](#)

Journal of Mathematical Physics **58**, 071702 (2017); 10.1063/1.4991031

[Superintegrable classical Zernike system](#)

Journal of Mathematical Physics **58**, 072901 (2017); 10.1063/1.4990793

[Analytic properties of the electromagnetic Green's function](#)

Journal of Mathematical Physics **58**, 071501 (2017); 10.1063/1.4993199

[Quantum superintegrable Zernike system](#)


Journal of Mathematical Physics **58**, 072101 (2017); 10.1063/1.4990794

[Mode stability on the real axis](#)

Journal of Mathematical Physics **58**, 072501 (2017); 10.1063/1.4991656

[Particle-like structure of Lie algebras](#)

Journal of Mathematical Physics **58**, 071703 (2017); 10.1063/1.4991657



Fock representations of Q -deformed commutation relations

Marek Bożejko,^{1,a)} Eugene Lytvynov,^{2,b)} and Janusz Wysoczański^{3,c)}

¹*Institute of Mathematics Polish Academy of Science, Kopernika 18, 51-617 Wrocław, Poland*

²*Department of Mathematics, Swansea University, Singleton Park, Swansea SA2 8PP, United Kingdom*

³*Instytut Matematyczny, Uniwersytet Wrocławski, pl. Grunwaldzki 2/4, 50-384 Wrocław, Poland*

(Received 17 March 2017; accepted 21 June 2017; published online 12 July 2017)

We consider Fock representations of the Q -deformed commutation relations $\partial_s \partial_t^\dagger = Q(s, t) \partial_t^\dagger \partial_s + \delta(s, t)$ for $s, t \in T$. Here $T := \mathbb{R}^d$ (or more generally T is a locally compact Polish space), the function $Q: T^2 \rightarrow \mathbb{C}$ satisfies $|Q(s, t)| \leq 1$ and $Q(s, t) = \overline{Q(t, s)}$, and $\int_{T^2} h(s)g(t)\delta(s, t)\sigma(ds)\sigma(dt) := \int_T h(t)g(t)\sigma(dt)$, σ being a fixed reference measure on T . In the case, where $|Q(s, t)| \equiv 1$, the Q -deformed commutation relations describe a generalized statistics studied by Liguori and Mintchev. These generalized statistics contain anyon statistics as a special case (with $T = \mathbb{R}^2$ and a special choice of the function Q). The related Q -deformed Fock space $\mathcal{F}(\mathcal{H})$ over $\mathcal{H} := L^2(T \rightarrow \mathbb{C}, \sigma)$ is constructed. An explicit form of the orthogonal projection of $\mathcal{H}^{\otimes n}$ onto the n -particle space $\mathcal{F}_n(\mathcal{H})$ is derived. A scalar product in $\mathcal{F}_n(\mathcal{H})$ is given by an operator $\mathcal{P}_n \geq 0$ in $\mathcal{H}^{\otimes n}$ which is strictly positive on $\mathcal{F}_n(\mathcal{H})$. We realize the smeared operators ∂_t^\dagger and ∂_t as creation and annihilation operators in $\mathcal{F}(\mathcal{H})$, respectively. Additional Q -commutation relations are obtained between the creation operators and between the annihilation operators. They are of the form $\partial_s^\dagger \partial_t^\dagger = Q(t, s) \partial_t^\dagger \partial_s^\dagger$, $\partial_s \partial_t = Q(t, s) \partial_t \partial_s$, valid for those $s, t \in T$ for which $|Q(s, t)| = 1$. *Published by AIP Publishing.* [<http://dx.doi.org/10.1063/1.4991671>]

I. INTRODUCTION

The aim of the paper is to construct Fock representations of the Q -commutation relations

$$\partial_s \partial_t^\dagger = Q(s, t) \partial_t^\dagger \partial_s + \delta(s, t), \quad s, t \in T. \quad (1)$$

Here $T = \mathbb{R}^d$, or more generally, T is a locally compact Polish space, the function $Q: T^2 \rightarrow \mathbb{C}$ is Hermitian, i.e., $Q(s, t) = \overline{Q(t, s)}$, and satisfies $|Q(s, t)| \leq 1$, ∂_t and ∂_t^\dagger are operator-valued distributions, adjoint of each other, and

$$\int_{T^2} \delta(s, t) f(s, t) \sigma(ds) \sigma(dt) := \int_T f(t, t) \sigma(dt),$$

where σ is a fixed Radon measure on X (typically $\sigma(dt) = dt$ being the Lebesgue measure if $T = \mathbb{R}^d$). We will call (1) the Q -deformed commutation relations, or just Q -CR.

For a function Q satisfying $|Q(s, t)| \equiv 1$, a Fock representation of the Q -CR was constructed by Liguori and Mintchev.²⁴ In that case, creation operators ∂_t^\dagger and annihilation operators ∂_t satisfy the

^{a)}E-mail: bozejko@math.uni.wroc.pl

^{b)}E-mail: e.lytvynov@swansea.ac.uk

^{c)}Author to whom correspondence should be addressed: jwys@math.uni.wroc.pl and janusz.wysoczanski@math.uni.wroc.pl.
 Tel.: +48 71 3757095.

additional commutation relations

$$\partial_s^\dagger \partial_t^\dagger = Q(t, s) \partial_t^\dagger \partial_s^\dagger, \quad \partial_s \partial_t = Q(t, s) \partial_t \partial_s. \quad (2)$$

The term *Fock representation* means that for each annihilation operator, one has $\partial_t \Omega = 0$, where Ω is the vacuum vector.

In the present study, relations (2) will hold for those $s, t \in T$ which satisfy $|Q(s, t)| = 1$. Note that under the assumption that the function Q is Hermitian, the commutation relations (2) are consistent if and only if $|Q(s, t)| = 1$.

For the first time, an interpolation between the canonical (bosonic) commutation relations (CCR) and the canonical (fermionic) anticommutation relations (CAR) was rigorously constructed in Ref. 7. Let \mathcal{H} be a separable Hilbert space and let $q \in (-1, 1)$. On a q -deformed Fock space over \mathcal{H} , Bożejko and Speicher⁷ constructed q -creation operators $a^+(f)$ [in fact $a^+(f)$ were *free* creation operators], and q -annihilation operators $a^-(f) := (a^+(f))^*$, for $f \in \mathcal{H}$, which satisfy the q -commutation relations

$$a^-(f)a^+(g) = qa^+(g)a^-(f) + (f, g)_{\mathcal{H}}, \quad f, g \in \mathcal{H}. \quad (3)$$

The limits $q = 1$ and $q = -1$ correspond to the boson and fermion statistics, respectively, thus giving the CCR and CAR. The case $q = 0$ corresponds to the creation and annihilation operators acting in the full Fock space; these operators are particularly important for models of free probability, see, e.g., Refs. 30 and 5. Aspects of noncommutative probability related to the general q -commutation relations (3) were discussed, e.g., in Refs. 7, 4, and 1.

By using probabilistic methods, Speicher³⁶ proved existence of a representation of the (discrete) q_{ij} -commutation relations of the form

$$\partial_i \partial_j^\dagger = q_{ij} \partial_j^\dagger \partial_i + \delta_{ij}, \quad (4)$$

with $-1 \leq q_{ij} = q_{ji} \leq 1$, $i, j \in \mathbb{N}$, and $(\partial_i^\dagger)^* = \partial_i$. Bożejko and Speicher⁸ constructed a Fock representation of the following commutation relations between creation operators ∂_j^\dagger and annihilation operators ∂_i , with $i, j \in \mathbb{N}$,

$$\partial_i \partial_j^\dagger = \sum_{k,l} q_{jl}^{ik} \partial_k^\dagger \partial_l + \delta_{ij}. \quad (5)$$

They showed that if the operator Ψ given by the matrix $(q_{jl}^{ik})_{i,j,k,l}$ is self-adjoint, satisfies the braid relations, and has norm < 1 , then there exists a Fock representation of the commutation relations (5). As a consequence, they obtained a Fock representation of the q_{ij} -commutation relations (4) even for complex q_{ij} with $\overline{q_{ij}} = q_{ji}$ and $\sup_{i,j} |q_{ij}| = \|\Psi\| < 1$. By taking the weak limit of corresponding operator algebras, Bożejko and Speicher⁸ also derived existence of a representation of the q_{ij} -commutation relations (4) with $\sup_{i,j} |q_{ij}| = \|\Psi\| = 1$. Also Jørgensen, Schmitt, and Werner^{18,19} considered representations of the commutation relations (5).

In the case, where $\|\Psi\| = 1$, Jørgensen, Proskurin, and Samoilenko²⁰ found, for $n \geq 2$, the kernel of the nonnegative operator which determines the scalar product in the n -particle space of the Fock space corresponding to the commutation relations (5). The papers Refs. 8 and 20 taken together give then a Fock representation of the commutation relations (4) with $\sup_{i,j} |q_{ij}| = 1$.

Properties of the algebras generated by such operators were studied by many authors. In the context of C^* -algebras, let us mention the works by Dykema and Nica¹¹ and Kennedy and Nica²¹ (who studied relations of the C^* -algebras generated by the q -commutation relations with the Cuntz algebra), Jørgensen, Schmitt, and Werner^{18,19} (who studied the Wick order generated C^* -algebras), and Proskurin and Samoilenko³² (who studied general Wick $*$ -algebras). There are also a number of studies of the q -commutation relations in the context of von Neumann algebras, in particular, by Lust-Piquard²⁵ (who studied properties of the Riesz transform), Królak,²² Nou,³¹ Śniady,³⁵ and Ricard³³ (who studied factoriality problems), Shlyakhtenko³⁴ (who studied Voiculescu's free entropy for families of q -Gaussian operators), and Bożejko² (who studied positivity of the symmetrization operators constructed through a self-adjoint Yang–Baxter operator $\Psi \geq -1$). Also Dabrowski,¹⁰ Guionnet and Shlyakhtenko,¹⁷ and Nelson and Zeng^{28,29} proved that q -factors or, more generally, q_{ij} -factors are isomorphic to the free group factors ($q = 0$) for small values of q or q_{ij} , respectively. Another possible

generalization of the commutation relations (3) related to the group of signed permutations can be found in Ref. 3.

All the above mentioned investigations are of *discrete* type so that the set T is at most countable. As we have already mentioned above, in the *continuous* setting, a Fock representation of Q -CR (1) and (2), called a *generalized statistics*, was constructed by Liguori and Mintchev,²⁴ see also Refs. 14–16, 13, and 6. A rigorous meaning of these commutation relations is given by smearing them with functions from $\mathcal{H} := L^2(T \rightarrow \mathbb{C}, \sigma)$. More precisely, defining for $f \in \mathcal{H}$ operators $a^+(f) := \int_T \sigma(dt) f(t) \partial_t^\dagger$ and $a^-(f) := \int_T \sigma(dt) \overline{f(t)} \partial_t$, we get the commutation relations

$$\begin{aligned} a^-(f)a^+(g) &= \int_{T^2} \sigma(ds) \sigma(dt) \overline{f(s)} g(t) Q(s, t) \partial_t^\dagger \partial_s + \int_T \overline{f(t)} g(t) \sigma(dt), \\ a^+(f)a^+(g) &= \int_{T^2} \sigma(ds) \sigma(dt) f(s) g(t) Q(t, s) \partial_t^\dagger \partial_s^\dagger, \\ a^-(f)a^-(g) &= \int_{T^2} \sigma(ds) \sigma(dt) \overline{f(s)} \overline{g(t)} Q(t, s) \partial_t \partial_s, \end{aligned}$$

where $f, g \in \mathcal{H}$. (Of course, the operator-valued integrals in these relations should be given a rigorous meaning.)

From the physical point of view, the most important case of a generalized statistics is the anyon statistics, where $T = \mathbb{R}^2$ and the function $Q(s, t)$ is determined by a complex parameter q with $|q| = 1$, namely,

$$Q(s, t) = \begin{cases} q, & \text{if } s^1 < t^1, \\ \bar{q}, & \text{if } s^1 > t^1. \end{cases} \tag{6}$$

Here, $s = (s^1, s^2), t = (t^1, t^2) \in \mathbb{R}^2$. Note that the value of the function Q on the set $\{(s, t) \in T^2 \mid s^1 = t^1\}$ does not matter for the Fock representation of the Q -CR. For an explanation as to why such commutation relations describe an anyon statistic, we refer the reader to Liguori and Mintchev’s paper²⁴ and to Goldin and Sharp’s paper.¹⁶

Goldin and Majid¹³ proved the following anyonic exclusion principle, which generalizes Pauli’s exclusion principle for fermions: If $q^m = 1$ and $q \neq 1$, then the creation operators $a^+(f)$ in the Fock representation of the anyon commutation relations satisfies $a^+(f)^m = 0$, or equivalently, the Q -symmetrization of the function $f^{\otimes m}$ is equal to zero.

In Ref. 26, non-Fock representations of the anyon commutation relations have been constructed, whose vacuum states are gauge-invariant quasi-free. Note that for those representations, the (real) value of the function $Q(s, t)$ for $s = t$ must be specified.

Let us mention that anyon systems have also been considered in the discrete setting, i.e., when $T \subset \mathbb{N}$, see, e.g., Refs. 12, 23, and 13. It should be, however, mentioned that when discussing the anyons in the discrete setting, Goldin and Majid¹³ dropped the assumption that the annihilation operator is adjoint of the creation operator and proved an anyonic exclusion principle for their model.

In this paper, we study the continuous case with a function Q satisfying $|Q(s, t)| \leq 1$. This natural choice of Q contains generalized statistics and the relations (3) as special cases. We would also like to draw the reader’s attention to the study by Merberg,²⁷ where the case $Q : T^2 \rightarrow (-1, 1)$ was considered and factoriality of the related von Neumann algebras generated by the Q -Gaussian operators was discussed.

In Sec. II, we present a construction of the Fock representation of Q -CR (1). To this end, we construct a certain Q -deformed Fock space over $\mathcal{H} = L^2(T \rightarrow \mathbb{C}, \sigma)$, denoted by $\mathcal{F}(\mathcal{H})$. We describe the n -particle subspaces, $\mathcal{F}_n(\mathcal{H})$, of $\mathcal{F}(\mathcal{H})$. As a set, each $\mathcal{F}_n(\mathcal{H})$ is a subset of $\mathcal{H}^{\otimes n} = L^2(T^n \rightarrow \mathbb{C}, \sigma^{\otimes n})$ and consists of all functions $f^{(n)} \in \mathcal{H}^{\otimes n}$ that are Q -quasisymmetric, meaning that, almost every, for each $k \in \{1, \dots, n - 1\}$,

$$f^{(n)}(t_1, \dots, t_n) = Q(t_k, t_{k+1}) f(t_1, \dots, t_{k+1}, t_k, \dots, t_n), \tag{7}$$

provided $|Q(t_k, t_{k+1})| = 1$. We derive an explicit formula for the orthogonal projection of $\mathcal{H}^{\otimes n}$ onto $\mathcal{F}_n(\mathcal{H})$. A scalar product in $\mathcal{F}_n(\mathcal{H})$ is given by an operator $\mathcal{P}_n \geq 0$ in $\mathcal{H}^{\otimes n}$ which is strictly positive on

$\mathcal{F}_n(\mathcal{H})$. We then realize $a^+(f)$ and $a^-(f)$ ($f \in \mathcal{H}$) as creation and annihilation operators acting in the Q -deformed Fock space $\mathcal{F}(\mathcal{H})$. These operators satisfy Q -CR (1). Additionally, due to the Q -symmetry (7) in each $\mathcal{F}_n(\mathcal{H})$, we get the following commutation relations between the creation operators and between the annihilation operators:

$$\begin{aligned} \partial_s^\dagger \partial_t^\dagger &= Q(t, s) \partial_t^\dagger \partial_s^\dagger, & \text{if } |Q(s, t)| = 1, \\ \partial_s \partial_t &= Q(t, s) \partial_t \partial_s, & \text{if } |Q(s, t)| = 1. \end{aligned} \tag{8}$$

We note that by choosing T to be a discrete set and σ to be the counting measure on T , one can apply our results in a discrete setting. In fact, the explicit description of the n -particle space $\mathcal{F}_n(\mathcal{H})$, explicit formula for the orthogonal projection of $\mathcal{H}^{\otimes n}$ onto $\mathcal{F}_n(\mathcal{H})$, and the additional commutation relations (8) appear to be new results even in the discrete setting.

We finish Sec. II with a proposition that shows that discrete anyons of fermion type satisfy the anyon exclusion principle, compare with Ref. 13.

In Sec. III, we prove the results formulated in Sec. II.

II. CONSTRUCTION OF THE FOCK REPRESENTATION OF Q -CR

In this section, we will construct a Fock representation of the commutation relation (1), and we will note that the additional commutation relations (8) then also hold.

A. Operator \mathcal{P}_n

Let T be a locally compact Polish space, let $\mathcal{B}(T)$ denote the Borel σ -algebra on T , and let σ be a Radon measure on $(T, \mathcal{B}(T))$. Let $E \in \mathcal{B}(T^2)$ be a symmetric subset of T : if $(s, t) \in E$ then $(t, s) \in E$. We assume that $\sigma^{\otimes 2}(E) = 0$. Denote $T^{(2)} := T^2 \setminus E$, which is also a symmetric set. We fix a complex-valued measurable function

$$Q : T^{(2)} \rightarrow \{z \in \mathbb{C} : |z| \leq 1\},$$

which is Hermitian: for all $(s, t) \in T^{(2)}$, we have $Q(s, t) = \overline{Q(t, s)}$. This function is defined $\sigma^{\otimes 2}$ -almost everywhere on T^2 .

Remark 1. The case where $|Q(s, t)| = 1$ for all $(s, t) \in T^{(2)}$ corresponds to a generalized statistics studied by Liguori and Mintchev.²⁴ The special case where $T = \mathbb{R}^2$, $\sigma(dt) = dt$ is the Lebesgue measure on T , $E = \{(s, t) \in T^2 \mid s^1 = t^1\}$, and the function Q is defined by formula (6) with $q \in \mathbb{C}$, $|q| = 1$, corresponds to anyon statistics, see Refs. 24, 16, and 13. The choice $Q(s, t) = q$ for all $(s, t) \in T^{(2)} = T^2$ with $q \in (-1, 1)$ corresponds to the q -commutations (3), see Ref. 7.

Let us consider an operator Ψ which transforms a measurable function $f^{(2)} : T^{(2)} \rightarrow \mathbb{C}$ into the function

$$(\Psi f^{(2)})(s, t) := Q(s, t) f^{(2)}(t, s), \quad (s, t) \in T^{(2)}. \tag{9}$$

Analogously to $T^{(2)}$, we define, for $n \geq 3$,

$$T^{(n)} := \{(t_1, \dots, t_n) \in T^n : (t_i, t_j) \notin E \text{ for all } 1 \leq i < j \leq n\}.$$

It is clear that $\sigma^{\otimes n}(T^n \setminus T^{(n)}) = 0$. The operator Ψ can be extended to a transformation of functions $f^{(n)} : T^{(n)} \rightarrow \mathbb{C}$ by setting, for $k \in \{1, \dots, n - 1\}$,

$$(\Psi_k f^{(n)})(t_1, \dots, t_n) := Q(t_k, t_{k+1}) f^{(n)}(t_1, \dots, t_{k-1}, t_{k+1}, t_k, t_{k+2}, \dots, t_n). \tag{10}$$

Let $\mathcal{H} := L^2(T \rightarrow \mathbb{C}, \sigma)$ be the complex L^2 -space over T . We agree that the scalar product $(\cdot, \cdot)_{\mathcal{H}}$ is antilinear in the first dot and linear in the second. For $n \geq 2$, the n th tensor power of \mathcal{H} , denoted by $\mathcal{H}^{\otimes n}$, can be identified with the complex L^2 -space $L^2(T^{(n)} \rightarrow \mathbb{C}, \sigma^{\otimes n})$. Each Ψ_k is a contraction in $\mathcal{H}^{\otimes n}$. The following trivial lemma shows that the operators Ψ_k are self-adjoint and satisfy the braid relations.

Lemma 2. The operators Ψ_k satisfy

$$\begin{aligned} \Psi_k^* &= \Psi_k, \\ \Psi_k \Psi_l &= \Psi_l \Psi_k \quad \text{if } |k - l| \geq 2, \\ \Psi_k \Psi_{k+1} \Psi_k &= \Psi_{k+1} \Psi_k \Psi_{k+1}. \end{aligned} \tag{11}$$

Let S_n denote the symmetric group on $\{1, \dots, n\}$. We represent a permutation $\pi \in S_n$ as an arbitrary product of adjacent transpositions,

$$\pi = \pi_{j_1} \dots \pi_{j_m}, \tag{12}$$

where $\pi_j := (j, j + 1) \in S_n$ for $1 \leq j \leq n - 1$. A permutation $\pi \in S_n$ can be represented (not in a unique way, in general) as a reduced product of a minimal number of adjacent transpositions, i.e., in form (12) with a minimal m . This number m is then called the length of π , denoted by $|\pi|$. It is well known that $|\pi|$ is equal to the number of inversions of π , i.e., the number of $1 \leq i < j \leq n$ such that $\pi(i) > \pi(j)$.

The mapping $\pi_k \mapsto \Psi_{\pi_k} := \Psi_k$ can be multiplicatively extended to S_n by setting

$$S_n \ni \pi \mapsto \Psi_\pi := \Psi_{j_1} \dots \Psi_{j_m}. \tag{13}$$

Although representation (12) of $\pi \in S_n$ in a reduced form is not unique, formulas (11) yield that extension (13) is well defined, i.e., it does not depend on the representation. (This fact also follows from the proof of Proposition 3 below.)

We will use the notations $\mathbf{t}^{(n)} := (t_1, \dots, t_n) \in T^{(n)}$ and $\mathbf{t}_\pi^{(n)} := (t_{\pi(1)}, \dots, t_{\pi(n)})$ for $\pi \in S_n$.

Proposition 3. For each $\pi \in S_n$ and $f^{(n)} \in \mathcal{H}^{\otimes n}$, we have

$$(\Psi_\pi f^{(n)})(\mathbf{t}^{(n)}) = Q_{\pi^{-1}}(\mathbf{t}^{(n)}) f^{(n)}(\mathbf{t}_\pi^{(n)}), \tag{14}$$

where

$$Q_\pi(\mathbf{t}^{(n)}) := \prod_{\substack{1 \leq i < j \leq n \\ \pi(i) > \pi(j)}} Q(t_i, t_j), \quad \mathbf{t}^{(n)} \in T^{(n)}. \tag{15}$$

For $n \geq 2$, we define an operator \mathcal{P}_n on $\mathcal{H}^{\otimes n}$ by

$$\mathcal{P}_n := \frac{1}{n!} \sum_{\pi \in S_n} \Psi_\pi. \tag{16}$$

The operator \mathcal{P}_n is a self-adjoint contraction in $\mathcal{H}^{\otimes n}$, so are the operators Ψ_k .

The following result is a special case of Theorem 1.1 in Ref. 8.

Theorem 4 (Ref. 8). For each $n \geq 2$, we have $\mathcal{P}_n \geq 0$.

For any $f^{(n)}, g^{(n)} \in \mathcal{H}^{\otimes n}$, we define

$$(f^{(n)}, g^{(n)})_{\mathcal{F}_n(\mathcal{H})} := (\mathcal{P}_n f^{(n)}, g^{(n)})_{\mathcal{H}^{\otimes n}}. \tag{17}$$

We consider the factor space

$$\mathcal{F}_n(\mathcal{H}) := \mathcal{H}^{\otimes n} / \{f^{(n)} \in \mathcal{H}^{\otimes n} : (f^{(n)}, f^{(n)})_{\mathcal{F}_n(\mathcal{H})} = 0\}$$

and define a scalar product on $\mathcal{F}_n(\mathcal{H})$ by (17).

Below, for a bounded linear operator L in a Hilbert space \mathfrak{H} , we denote by $\text{Ker}(L)$ and $\text{Ran}(L)$ the kernel of L and the range of L , respectively. Recall that $\text{Ker}(L)$ is a closed linear subspace of \mathfrak{H} and, if L is self-adjoint,

$$\mathfrak{H} = \text{Ker}(L) \oplus \overline{\text{Ran}(L)},$$

where $\overline{\text{Ran}(L)}$ denotes the closure of the linear subspace $\text{Ran}(L)$. The following lemma only uses the fact that $\mathcal{P}_n \geq 0$.

Lemma 5. (i) We have

$$\{f^{(n)} \in \mathcal{H}^{\otimes n} : (f^{(n)}, f^{(n)})_{\mathcal{F}_n(\mathcal{H})} = 0\} = \text{Ker}(\mathcal{P}_n).$$

(ii) For each $f^{(n)} \in \overline{\text{Ran}(\mathcal{P}_n)}$, $f^{(n)} \neq 0$,

$$(f^{(n)}, f^{(n)})_{\mathcal{F}_n(\mathcal{H})} > 0.$$

By Lemma 5, we can identify $\mathcal{F}_n(\mathcal{H})$ with the set $\overline{\text{Ran}(\mathcal{P}_n)}$ equipped with scalar product (17). The result below follows from Theorem 2 and Remark 4 in Ref. 20.

Theorem 6 (Ref. 20). *We have*

$$\text{Ker}(\mathcal{P}_n) = \overline{\sum_{k=1}^{n-1} \text{Ker}(\mathbf{1} + \Psi_k)}, \tag{18}$$

i.e., the kernel of \mathcal{P}_n is equal to the closure of the linear span of the subspaces $\text{Ker}(\mathbf{1} + \Psi_k)$, $k = 1, \dots, n - 1$.

We will now give an explicit description of the space $\mathcal{F}_n(\mathcal{H}) = \overline{\text{Ran}(\mathcal{P}_n)}$. We denote

$$\Theta := \{(s, t) \in T^{(2)} : |Q(s, t)| = 1\}, \quad \Theta' := T^{(2)} \setminus \Theta = \{(s, t) \in T^{(2)} : |Q(s, t)| < 1\}. \tag{19}$$

Theorem 7. *The space $\mathcal{F}_n(\mathcal{H}) = \overline{\text{Ran}(\mathcal{P}_n)}$ is equal (as a set) to the subspace of $\mathcal{H}^{\otimes n}$ consisting of all $f^{(n)} \in \mathcal{H}^{\otimes n}$ that are Q -quasisymmetric, i.e., formula (7) holds for each $k \in \{1, \dots, n - 1\}$ and for $\sigma^{\otimes n}$ -a.a. $(t_1, \dots, t_n) \in T^{(n)}$ such that $|Q(t_k, t_{k+1})| = 1$, i.e., for $\sigma^{\otimes n}$ -a.a. $(t_1, \dots, t_n) \in T_k^{(n)}$, where*

$$T_k^{(n)} := \{(t_1, \dots, t_n) \in T^{(n)} : (t_k, t_{k+1}) \in \Theta\}. \tag{20}$$

B. Orthogonal projection onto $\overline{\text{Ran}(\mathcal{P}_n)}$

We will now describe the orthogonal projection \mathbb{P}_n of $\mathcal{H}^{\otimes n}$ onto $\overline{\text{Ran}(\mathcal{P}_n)} = \mathcal{F}_n(\mathcal{H})$. For this purpose, we define a function

$$R(s, t) := \begin{cases} Q(s, t), & \text{if } (s, t) \in \Theta, \\ 0, & \text{if } (s, t) \in \Theta'. \end{cases}$$

Observe that

$$|R(s, t)| = \begin{cases} 1, & \text{if } (s, t) \in \Theta, \\ 0, & \text{if } (s, t) \in \Theta', \end{cases}$$

and that the function R is Hermitian. Hence, for each $\pi \in S_n$, similarly to the operator $\Psi_\pi : \mathcal{H}^{\otimes n} \rightarrow \mathcal{H}^{\otimes n}$ defined in Subsection II A for the function $Q(s, t)$, we may define an operator $\Phi_\pi : \mathcal{H}^{\otimes n} \rightarrow \mathcal{H}^{\otimes n}$ for the function $R(s, t)$. By Proposition 3, we get

$$(\Phi_\pi f^{(n)})(\mathbf{t}^{(n)}) = R_{\pi^{-1}}(\mathbf{t}^{(n)}) f^{(n)}(\mathbf{t}_\pi^{(n)}), \tag{21}$$

where

$$R_\pi(\mathbf{t}^{(n)}) := \prod_{\substack{1 \leq i < j \leq n \\ \pi(i) > \pi(j)}} R(t_i, t_j), \quad \mathbf{t}^{(n)} \in T^{(n)}. \tag{22}$$

Let $\pi \in S_n$ and let $\mathbf{t}^{(n)} \in T^{(n)}$ be such that for some $1 \leq i < j \leq n$, we have $\pi(i) > \pi(j)$ and $(t_i, t_j) \in \Theta'$. Then, it follows from (22) that $R_\pi(\mathbf{t}^{(n)}) = 0$. Otherwise, i.e., if such i and j do not exist, we get $|R_\pi(\mathbf{t}^{(n)})| = 1$.

Given $\mathbf{t}^{(n)} \in T^{(n)}$, we define a splitting

$$S_n = S_n^1(\mathbf{t}^{(n)}) \sqcup S_n^0(\mathbf{t}^{(n)})$$

of the set S_n into two disjoint subsets:

$$\begin{aligned} S_n^1(\mathbf{t}^{(n)}) &:= \{\pi \in S_n : |R_{\pi^{-1}}(\mathbf{t}^{(n)})| = 1\}, \\ S_n^0(\mathbf{t}^{(n)}) &:= \{\pi \in S_n : |R_{\pi^{-1}}(\mathbf{t}^{(n)})| = 0\}. \end{aligned} \tag{23}$$

Let $c_n(\mathbf{t}^{(n)}) := |S_n^1(\mathbf{t}^{(n)})|$ denote the cardinality. We define an operator $\mathbb{P}_n : \mathcal{H}^{\otimes n} \rightarrow \mathcal{H}^{\otimes n}$ by setting, for each $f^{(n)} \in \mathcal{H}^{\otimes n}$,

$$\begin{aligned} (\mathbb{P}_n f^{(n)})(\mathbf{t}^{(n)}) &:= \frac{1}{c_n(\mathbf{t}^{(n)})} \sum_{\pi \in S_n^1(\mathbf{t}^{(n)})} (\Phi_\pi f^{(n)})(\mathbf{t}^{(n)}) \\ &= \frac{1}{c_n(\mathbf{t}^{(n)})} \sum_{\pi \in S_n^1(\mathbf{t}^{(n)})} R_{\pi^{-1}}(\mathbf{t}^{(n)}) f^{(n)}(\mathbf{t}_\pi^{(n)}). \end{aligned} \tag{24}$$

Theorem 8. For each $n \geq 2$, the operator \mathbb{P}_n is the orthogonal projection of $\mathcal{H}^{\otimes n}$ onto $\overline{\text{Ran}(\mathbb{P}_n)} = \mathcal{F}_n(\mathcal{H})$.

The corollary below is a straightforward consequence of Theorem 8.

Corollary 9. For each $n \geq 2$,

$$\mathbb{P}_n \mathcal{P}_n = \mathcal{P}_n \mathbb{P}_n = \mathcal{P}_n.$$

We will also need the following result about the operators \mathbb{P}_n , which follows from Theorem 8 and its proof.

Corollary 10. For each $n \geq 2$ and $k \in \{1, \dots, n - 1\}$, we have

$$\mathbb{P}_n = \mathbb{P}_n (\mathbb{P}_k \otimes \mathbb{P}_{n-k}). \tag{25}$$

Here we denote by $\mathbb{P}_1 := \mathbf{1}$ the identity operator in \mathcal{H} .

Remark 11. For $f^{(n)} \in \mathcal{F}_n(\mathcal{H})$ and $g^{(m)} \in \mathcal{F}_m(\mathcal{H})$, we may define a Q -quasisymmetric tensor product of $f^{(n)}$ and $g^{(m)}$ by

$$f^{(n)} \otimes g^{(m)} := \mathbb{P}_{n+m}(f^{(n)} \otimes g^{(m)}).$$

Then Corollary 10 implies that the Q -quasisymmetric tensor product \otimes is associative.

C. Creation and annihilation operators and their Q -commutation relations

Recall that we have defined complex Hilbert spaces $\mathcal{F}_n(\mathcal{H})$ for $n \geq 2$. Let also $\mathcal{F}_1(\mathcal{H}) := \mathcal{H}$ and $\mathcal{F}_0(\mathcal{H}) := \mathbb{C}$. We define a Q -deformed Fock space to be the Hilbert space

$$\mathcal{F}(\mathcal{H}) := \bigoplus_{n=0}^{\infty} \mathcal{F}_n(\mathcal{H}) n!.$$

Thus, every $f \in \mathcal{F}(\mathcal{H})$ is represented as $f = (f^{(n)})_{n=0}^{\infty}$, where $f^{(n)} \in \mathcal{F}_n(\mathcal{H})$, and the norm of f is given by

$$\|f\|_{\mathcal{F}(\mathcal{H})}^2 := \sum_{n=0}^{\infty} \|f^{(n)}\|_{\mathcal{F}_n(\mathcal{H})}^2 n!.$$

The vector $\Omega := (1, 0, 0, \dots)$ is called the vacuum.

Let $\mathcal{F}_{\text{fin}}(\mathcal{H}) \subset \mathcal{F}(\mathcal{H})$ be the subspace consisting of all finite sequences of the form $f = (f^{(0)}, f^{(1)}, \dots, f^{(k)}, 0, 0, \dots)$ for some $k \in \mathbb{N}$. The subspace $\mathcal{F}_{\text{fin}}(\mathcal{H})$ is evidently dense in $\mathcal{F}(\mathcal{H})$.

For each $h \in \mathcal{H}$, we define a creation operator $a^+(h) : \mathcal{F}_{\text{fin}}(\mathcal{H}) \rightarrow \mathcal{F}_{\text{fin}}(\mathcal{H})$ by setting

$$a^+(h)\Omega := h, \quad a^+(h)f^{(n)} := \mathbb{P}_{n+1}(h \otimes f^{(n)}), \quad f^{(n)} \in \mathcal{F}_n(\mathcal{H}), \quad n \in \mathbb{N}. \tag{26}$$

The domain of the adjoint operator of $a^+(h)$ in $\mathcal{F}(\mathcal{H})$ contains $\mathcal{F}_{\text{fin}}(\mathcal{H})$, and furthermore the annihilation operator $a^-(h) := (a^+(h))^* \upharpoonright \mathcal{F}_{\text{fin}}(\mathcal{H})$ also maps $\mathcal{F}_{\text{fin}}(\mathcal{H})$ into itself.

The following proposition gives an explicit form of the action of the annihilation operator.

Proposition 12. For each $h \in \mathcal{H}$, we have $a^-(h)\Omega = 0$, $a^-(h)g = (h, g)_{\mathcal{H}}$ for $g \in \mathcal{H}$, and

$$\begin{aligned} &(a^-(h)f^{(n)})(t_1, \dots, t_{n-1}) \\ &= \sum_{k=1}^n \mathbb{P}_{n-1} \left[\int_T \overline{h(s)} \left(\prod_{i=1}^{k-1} Q(s, t_i) \right) f^{(n)}(t_1, \dots, t_{k-1}, s, t_k, \dots, t_{n-1}) \sigma(ds) \right] \end{aligned} \tag{27}$$

for any $f^{(n)} \in \mathcal{F}_n(\mathcal{H})$, $n \geq 2$. In formula (27), the operator \mathbb{P}_{n-1} acts on the function of t_1, \dots, t_{n-1} variables. Furthermore, for any $g^{(n)} \in \mathcal{H}^{\otimes n}$, $n \geq 2$,

$$\begin{aligned} & (a^-(h)\mathbb{P}_n g^{(n)})(t_1, \dots, t_{n-1}) \\ &= \sum_{k=1}^n \mathbb{P}_{n-1} \left[\int_T \overline{h(s)} \left(\prod_{i=1}^{k-1} Q(s, t_i) \right) g^{(n)}(t_1, \dots, t_{k-1}, s, t_k, \dots, t_{n-1}) \sigma(ds) \right]. \end{aligned} \tag{28}$$

For $t \in T$, we now informally define creation and annihilation operators at point t , denoted by ∂_t^\dagger and ∂_t , respectively. A rigorous meaning to these operators is given through smearing them with functions $h \in \mathcal{H}$,

$$a^+(h) = \int_T \sigma(dt) h(t) \partial_t^\dagger, \quad a^-(h) = \int_T \sigma(dt) \overline{h(t)} \partial_t. \tag{29}$$

So we have the following informal equalities:

$$\begin{aligned} \partial_t^\dagger f^{(n)} &= \mathbb{P}_{n+1}(\delta_t \otimes f^{(n)}), \\ \partial_t f^{(n)}(t_1, \dots, t_{n-1}) &= \sum_{k=1}^n \mathbb{P}_{n-1} \left[\left(\prod_{i=1}^{k-1} Q(t, t_i) \right) f^{(n)}(t_1, \dots, t_{k-1}, t, t_k, \dots, t_{n-1}) \right], \end{aligned}$$

where δ_t denotes the delta function at t .

Using (26) and Corollary 10, we see that for any $g, h \in \mathcal{H}$ and $f^{(n)} \in \mathcal{F}_n(\mathcal{H})$,

$$a^+(g)a^+(h)f^{(n)} := \mathbb{P}_{n+2}(g \otimes h \otimes f^{(n)}). \tag{30}$$

In view of (29) and (30), for each $\varphi^{(2)} \in \mathcal{H}^{\otimes 2}$, we can naturally define an operator

$$\int_{T^2} \sigma(ds) \sigma(dt) \varphi^{(2)}(s, t) \partial_s^\dagger \partial_t^\dagger : \mathcal{F}_{\text{fin}}(\mathcal{H}) \rightarrow \mathcal{F}_{\text{fin}}(\mathcal{H})$$

by setting

$$\int_{T^2} \sigma(ds) \sigma(dt) \varphi^{(2)}(s, t) \partial_s^\dagger \partial_t^\dagger f^{(n)} := \mathbb{P}_{n+2}(\varphi^{(2)} \otimes f^{(n)}) \tag{31}$$

for $f^{(n)} \in \mathcal{F}_n(\mathcal{H})$. In particular, choosing $\varphi^{(2)} = g \otimes h$ with $g, h \in \mathcal{H}$, we get

$$\int_{T^2} \sigma(ds) \sigma(dt) g(s)h(t) \partial_s^\dagger \partial_t^\dagger = a^+(g)a^+(h).$$

Remark 13. Note that we also accept the natural formula

$$\int_{T^2} \sigma(ds) \sigma(dt) \varphi^{(2)}(s, t) \partial_t^\dagger \partial_s^\dagger = \int_{T^2} \sigma(ds) \sigma(dt) \varphi^{(2)}(t, s) \partial_s^\dagger \partial_t^\dagger. \tag{32}$$

Similarly, using also Proposition 12, we may define, for each $\varphi^{(2)} \in \mathcal{H}^{\otimes 2}$, linear operators

$$\begin{aligned} & \int_{T^2} \sigma(ds) \sigma(dt) \varphi^{(2)}(s, t) \partial_s \partial_t : \mathcal{F}_{\text{fin}}(\mathcal{H}) \rightarrow \mathcal{F}_{\text{fin}}(\mathcal{H}), \\ & \int_{T^2} \sigma(ds) \sigma(dt) \varphi^{(2)}(s, t) \partial_s^\dagger \partial_t : \mathcal{F}_{\text{fin}}(\mathcal{H}) \rightarrow \mathcal{F}_{\text{fin}}(\mathcal{H}). \end{aligned}$$

Note that

$$\begin{aligned} \left(\int_{T^2} \sigma(ds) \sigma(dt) \varphi^{(2)}(s, t) \partial_s^\dagger \partial_t^\dagger \right)^* &= \int_{T^2} \sigma(ds) \sigma(dt) \overline{\varphi^{(2)}(s, t)} \partial_t \partial_s \\ &= \int_{T^2} \sigma(ds) \sigma(dt) \overline{\varphi^{(2)}(t, s)} \partial_s \partial_t. \end{aligned} \tag{33}$$

Also, for any $g, h \in \mathcal{H}$, we denote

$$\int_{T^2} \sigma(ds) \sigma(dt) g(s)h(t) \partial_s \partial_t^\dagger := a^-(\overline{g})a^+(h).$$

We will now present the commutation relations for the creation and annihilation operators.

Theorem 14 (Q-CR). *The operators ∂_t^\dagger and ∂_t ($t \in T$) satisfy the (informal) commutations relations (1) and (8). Rigorously, this means the following: for any $g, h \in \mathcal{H}$,*

$$\int_{T^2} \sigma(ds) \sigma(dt) g(s)h(t) \partial_s \partial_t^\dagger = \int_T g(t)h(t) \sigma(dt) + \int_{T^2} \sigma(ds) \sigma(dt) g(s)h(t) Q(s, t) \partial_t^\dagger \partial_s, \quad (34)$$

and for any function $\varphi^{(2)} \in \mathcal{H}^{\otimes 2}$ that vanishes almost every in Θ' [see (19)],

$$\int_{T^2} \sigma(ds) \sigma(dt) \varphi^{(2)}(s, t) \partial_s^\dagger \partial_t^\dagger = \int_{T^2} \sigma(ds) \sigma(dt) \varphi^{(2)}(s, t) Q(t, s) \partial_t^\dagger \partial_s^\dagger, \quad (35)$$

$$\int_{T^2} \sigma(ds) \sigma(dt) \varphi^{(2)}(s, t) \partial_s \partial_t = \int_{T^2} \sigma(ds) \sigma(dt) \varphi^{(2)}(s, t) Q(t, s) \partial_t \partial_s. \quad (36)$$

We finish this section with several remarks.

Remark 15. We can naturally identify the diagonal $\Delta := \{(s, t) \in T^2 \mid s = t\}$ with T . Denote by $\tilde{\sigma}$, the measure σ on Δ . We may consider $\tilde{\sigma}$ as a measure on T^2 which is equal to zero outside of Δ . Denote

$$\mathfrak{G} := L^2(T^2 \rightarrow \mathbb{C}, \sigma^{\otimes 2}) \cap L^1(T^2 \rightarrow \mathbb{C}, \tilde{\sigma}).$$

In view of (34), for each $\varphi^{(2)} \in \mathfrak{G}$, we may define an operator

$$\int_{T^2} \sigma(ds) \sigma(dt) \varphi^{(2)}(s, t) \partial_s \partial_t^\dagger : \mathcal{F}_{\text{fin}}(\mathcal{H}) \rightarrow \mathcal{F}_{\text{fin}}(\mathcal{H}),$$

which satisfies

$$\int_{T^2} \sigma(ds) \sigma(dt) \varphi^{(2)}(s, t) \partial_s \partial_t^\dagger = \int_T \varphi^{(2)}(t, t) \sigma(dt) + \int_{T^2} \sigma(ds) \sigma(dt) \varphi^{(2)}(s, t) Q(s, t) \partial_t^\dagger \partial_s.$$

Remark 16. Denote $B(\varphi) := a^+(\varphi) + a^-(\varphi)$. The family of operators $(B(\varphi))_{\varphi \in \mathcal{H}}$ can be thought of as a *noncommutative Brownian motion* (or a *noncommutative Gaussian white noise*). Let \mathcal{P} denote the complex unital $*$ -algebra generated by $(B(\varphi))_{\varphi \in \mathcal{H}}$, i.e., the algebra of noncommutative polynomials in the variables $B(\varphi)$. We define a vacuum state on \mathcal{P} by $\tau(p) := (p\Omega, \Omega)_{\mathcal{F}(\mathcal{H})}$, $p \in \mathcal{P}$. By analogy with the proofs of Theorem 4.4 in Ref. 8 and Corollary 4.9 in Ref. 6, one can prove the following result: the state τ is tracial (i.e., it satisfies $\tau(p_1 p_2) = \tau(p_2 p_1)$ for all $p_1, p_2 \in \mathcal{P}$) if and only if the function Q is real-valued, i.e., $Q : T^{(2)} \rightarrow [-1, 1]$.

Remark 17. The results of this section hold, in particular, in the case where $\sigma^{\otimes 2}(\Theta') = 0$, i.e., when $|Q(s, t)| < 1$ for $\sigma^{\otimes 2}$ -a.a. $(s, t) \in T^2$. Then, for each $n \geq 2$, the equality $\mathcal{F}_n(\mathcal{H}) = \mathcal{H}^{\otimes n}$ holds (in the sense of sets). Evidently, there are no commutation relations (35) and (36) in this case. Note also that if $|Q(s, t)| \leq r < 1$ for some number $0 < r < 1$, then the creation operators $a^+(h)$ and the annihilation operators $a^-(h)$ ($h \in \mathcal{H}$) are bounded in $\mathcal{F}(\mathcal{H})$, see Theorem 3.1, (ii) in Ref. 8.

D. Discrete setting: The anyonic exclusion principle

We will now make several observations about the discrete setting. We may choose T to be a finite or countable set and σ to be the counting measure on T , i.e., $\sigma(\{t\}) = 1$ for each $t \in T$. Hence, the space \mathcal{H} becomes the complex ℓ^2 -space over T , i.e., $\mathcal{H} = \ell^2(T \rightarrow \mathbb{C})$. We obviously have $T^{(2)} = T^2$ so that the function $Q(s, t)$ is defined for all $(s, t) \in T^2$. Thus, we have, in particular, constructed Fock representations of the discrete commutation relations (4) with additional commutation relations between ∂_s^\dagger and ∂_t^\dagger and between ∂_s and ∂_t for those pairs $(s, t) \in T^2$ for which $|Q(s, t)| = 1$. (Note that in this case, the operators $\partial_t^\dagger, \partial_t$ have a rigorous meaning.)

Since the function Q is Hermitian, we have $Q(t, t) \in \mathbb{R}$ for each $t \in T$. Hence, $|Q(t, t)| = 1$ if and only if either $Q(t, t) = 1$ or $Q(t, t) = -1$. In the first case, we just get the tautological commutation relation $(\partial_t^\dagger)^2 = (\partial_t^\dagger)^2$. In the second case, we get $(\partial_t^\dagger)^2 = -(\partial_t^\dagger)^2$ so that $(\partial_t^\dagger)^2 = \partial_t^2 = 0$. If the latter formulas hold for all $t \in T$, then we may call the corresponding commutation relations the *discrete Q-CR of fermion type*.

For the discrete Q -CR of fermion type, the operators ∂_t^\dagger and ∂_t become bounded in $\mathcal{F}(\mathcal{H})$ and have norm equal to 1, see Ref. 8, Corollary 3.2 and Remark after it. Hence, for each $h \in \ell^1(T \rightarrow \mathbb{C})$,

$$\|a^+(h)\| = \|a^-(h)\| \leq \|h\|_{\ell^1(T \rightarrow \mathbb{C})}.$$

Let us now assume that $T \subset \mathbb{N}$ and fix $q \in \mathbb{C}$, $|q| = 1$. We consider the function

$$Q(s, t) := \begin{cases} q, & \text{if } s > t, \\ \bar{q}, & \text{if } s < t, \\ -1, & \text{if } s = t. \end{cases}$$

The corresponding Q -CR describes a *discrete anyon system of fermion type*. Note that $|Q(s, t)| = 1$ for all $(s, t) \in T^2$, hence $\mathcal{P}_n = \mathbb{P}_n$ is the projection of $\mathcal{H}^{\otimes n}$ onto $\mathcal{F}_n(\mathcal{H})$.

Theorem 18 (Anyonic exclusion principle). *Consider a discrete anyon system of fermion type. Let $m \in \mathbb{N}$, $m \geq 2$. Assume that the parameter $q \in \mathbb{C}$, $q \neq 1$, is an m th root of unity, i.e., $q^m = 1$. Then, for any $h \in \mathcal{H}$, we have*

$$a^+(h)^m = a^-(h)^m = 0. \tag{37}$$

III. PROOFS

In this section, we collect the proofs of the results from Sec. II.

Proof of Proposition 3. We start with the following crucial lemma.

Lemma 19. *Let $\rho = \pi_l \eta$ be a reduced representation of a permutation $\rho \in S_n$. Then*

$$Q_\rho(t_1, \dots, t_n) = Q(t_{\eta^{-1}(l)}, t_{\eta^{-1}(l+1)}) Q_\eta(t_1, \dots, t_n), \quad (t_1, \dots, t_n) \in T^{(n)}. \tag{38}$$

Proof. Let

$$L_\rho := Q_\rho(t_1, \dots, t_n) = \prod_{\substack{1 \leq i < j \leq n \\ \rho(i) > \rho(j)}} Q(t_i, t_j), \quad L_\eta := Q_\eta(t_1, \dots, t_n) = \prod_{\substack{1 \leq i < j \leq n \\ \eta(i) > \eta(j)}} Q(t_i, t_j).$$

Let $1 \leq u < v \leq n$. We consider the following cases.

- If $\eta(u), \eta(v) \notin \{l, l + 1\}$, then both $\eta(u), \eta(v)$ are fixed points for π_l . Consequently, $\rho(u) = \eta(u)$ and $\rho(v) = \eta(v)$ so that $\rho(u) > \rho(v)$ if and only if $\eta(u) > \eta(v)$. Hence, the term $Q(t_u, t_v)$ appears in L_ρ if and only if it appears in L_η .
- If $\eta(u) \in \{l, l + 1\}$ and $\eta(v) \notin \{l, l + 1\}$, then $\rho(v) = v \notin \{l, l + 1\}$ and, since $\rho(u) = (\pi_l \eta)(u) \in \{l, l + 1\}$, the order between $\eta(u)$ and $\eta(v)$ is the same as between $\rho(u)$ and $\rho(v)$. Thus, the term $Q(t_u, t_v)$ appears in L_ρ if and only if it appears in L_η .
- The case $\eta(u) \notin \{l, l + 1\}$ and $\eta(v) \in \{l, l + 1\}$ is analogous to the previous one.
- Consider the case $\eta(u) = l$ and $\eta(v) = l + 1$. Then the term $Q(t_u, t_v)$ does not appear in L_η . Further, $\rho(u) = (\pi_l \eta)(u) = \pi_l(l) = l + 1$ and $\rho(v) = (\pi_l \eta)(v) = \pi_l(l + 1) = l$ so that $\rho(u) > \rho(v)$. Hence, the term $Q(t_u, t_v)$ appears in L_ρ . But we also have $Q(t_{\eta^{-1}(l)}, t_{\eta^{-1}(l+1)}) = Q(t_u, t_v)$ on the right hand side of equality (38).
- Finally, consider the case $\eta(u) = l + 1$ and $\eta(v) = l$. But then $\rho(u) = (\pi_l \eta)(u) = l$ and $\rho(v) = (\pi_l \eta)(v) = l + 1$. Thus, η changes the order of the pair (u, v) , while ρ does not. Therefore, η has more inversions than ρ : $|\eta| > |\rho|$. But this contradicts the assumption that ρ is in the reduced form. Thus, this case is impossible. \square

We will now prove the proposition by induction on the length of a permutation π . If $|\pi| = 1$, then $\pi = \pi_k$ for some $k \in \{1, \dots, n - 1\}$. In this case, the statement trivially follows from the definition of Ψ_k , see (10). Assume that the statement holds for each permutation of length m . Let π be a permutation of length $m + 1$, and let $\pi = \varphi \pi_l$ be a reduced representation of π . Hence, the length of the permutation φ

is m . Denote $\eta := \varphi^{-1}$ and $\rho := \pi^{-1}$ so that $\rho = \pi_l \eta$. Then, for each $f^{(n)} \in \mathcal{H}^{\otimes n}$, by using the induction's assumption and Lemma 19, we get

$$\begin{aligned} (\Psi_\pi f^{(n)})(t_1, \dots, t_n) &= (\Psi_\varphi \Psi_l f^{(n)})(t_1, \dots, t_n) \\ &= Q_\eta(t_1, \dots, t_n) (\Psi_l f^{(n)})(t_{\varphi(1)}, \dots, t_{\varphi(n)}) \\ &= Q_\eta(t_1, \dots, t_n) Q(t_{\varphi(l)}, t_{\varphi(l+1)}) f^{(n)}(t_{\varphi(1)}, \dots, t_{\varphi(l+1)}, t_{\varphi(l)}, \dots, t_{\varphi(n)}) \\ &= Q_\rho(t_1, \dots, t_n) f^{(n)}(t_{\pi(1)}, \dots, t_{\pi(n)}). \end{aligned}$$

Proof of Lemma 5. (i) Since \mathcal{P}_n is self-adjoint and $\mathcal{P}_n \geq 0$, we can write $\mathcal{P}_n = (\sqrt{\mathcal{P}_n})^2$. Let $f^{(n)} \in \mathcal{H}^{\otimes n}$ be such that \square

$$0 = (f^{(n)}, f^{(n)})_{\mathcal{F}_n(\mathcal{H})} = \|\sqrt{\mathcal{P}_n} f^{(n)}\|_{\mathcal{H}^{\otimes n}}^2.$$

Hence, $f^{(n)} \in \text{Ker}(\sqrt{\mathcal{P}_n})$. But $\text{Ker} \sqrt{\mathcal{P}_n} \subset \text{Ker} \mathcal{P}_n$, which implies

$$\{f^{(n)} \in \mathcal{H}^{\otimes n} \mid (f^{(n)}, f^{(n)})_{\mathcal{F}_n(\mathcal{H})} = 0\} \subset \text{Ker}(\mathcal{P}_n).$$

The inverse inclusion trivially follows from (17).

(ii) Let $f^{(n)} \in \overline{\text{Ran}(\mathcal{P}_n)}$ be such that $(f^{(n)}, f^{(n)})_{\mathcal{F}_n(\mathcal{H})} = 0$. By part (i), $f^{(n)} \in \text{Ker}(\mathcal{P}_n)$. But $\overline{\text{Ran}(\mathcal{P}_n)} \perp \text{Ker}(\mathcal{P}_n)$. Hence, $\overline{\text{Ran}(\mathcal{P}_n)} \cap \text{Ker}(\mathcal{P}_n) = \{0\}$, and so $f^{(n)} = 0$. \square

Proof of Theorem 7. Using (18), we have

$$\overline{\text{Ran}(\mathcal{P}_n)} = \left(\sum_{k=1}^{n-1} \text{Ker}(\mathbf{1} + \Psi_k) \right)^\perp = \bigcap_{k=1}^{n-1} \text{Ker}(\mathbf{1} + \Psi_k)^\perp = \bigcap_{k=1}^{n-1} \overline{\text{Ran}(\mathbf{1} + \Psi_k)}. \tag{39}$$

For $l \in \mathbb{N}$ and $k \in \{1, \dots, n - 1\}$, we denote

$$T_{k,l}^{(n)} := \left\{ (t_1, \dots, t_n) \in T^{(n)} : \frac{l-1}{l} \leq |Q(t_k, t_{k+1})| < \frac{l}{l+1} \right\}$$

and recall the definition of $T_k^{(n)}$, see (20). Then, for each $k \in \{1, \dots, n - 1\}$, we have the orthogonal decomposition

$$\mathcal{H}^{\otimes n} = \left(\bigoplus_{l=1}^{\infty} L^2(T_{k,l}^{(n)} \rightarrow \mathbb{C}, \sigma^{\otimes n}) \right) \oplus L^2(T_k^{(n)} \rightarrow \mathbb{C}, \sigma^{\otimes n}). \tag{40}$$

Each of the spaces on the right-hand side of (40) is invariant for the operator $\mathbf{1} + \Psi_k$. On each space $L^2(T_{k,l}^{(n)} \rightarrow \mathbb{C}, \sigma^{\otimes n})$, the norm of the operator Ψ_k is bounded by $\frac{l}{l+1} < 1$. Hence, the operator $\mathbf{1} + \Psi_k$ is invertible in this space. Therefore the kernel of the operator $\mathbf{1} + \Psi_k$ restricted to $L^2(T_{k,l}^{(n)} \rightarrow \mathbb{C}, \sigma^{\otimes n})$ is trivial

$$\text{Ker}(\mathbf{1} + \Psi_k) \cap L^2(T_{k,l}^{(n)} \rightarrow \mathbb{C}, \sigma^{\otimes n}) = \{0\} \quad \text{for each } l \in \mathbb{N}.$$

Let $f^{(n)} \in L^2(T_k^{(n)} \rightarrow \mathbb{C}, \sigma^{\otimes n})$. Consider the decomposition $f^{(n)} = f_{k,+}^{(n)} + f_{k,-}^{(n)}$ with

$$f_{k,\pm}^{(n)}(t_1, \dots, t_n) := \frac{1}{2} [f^{(n)}(t_1, \dots, t_n) \pm Q(t_k, t_{k+1}) f^{(n)}(t_1, \dots, t_{k+1}, t_k, \dots, t_n)].$$

One can easily see that $f_{k,+}^{(n)}$ and $f_{k,-}^{(n)}$ are orthogonal and $f_{k,+}^{(n)} \in \text{Ran}(\mathbf{1} + \Psi_k)$. Hence $f_{k,-}^{(n)} \in \text{Ker}(\mathbf{1} + \Psi_k)$. Therefore, the orthogonal projection of $L^2(T_k^{(n)} \rightarrow \mathbb{C}, \sigma^{\otimes n})$ onto $\text{Ker}(\mathbf{1} + \Psi_k)$, denoted by $D_k^{(n)}$, is given by

$$(D_k^{(n)} f^{(n)})(t_1, \dots, t_n) = \frac{1}{2} [f^{(n)}(t_1, \dots, t_n) - Q(t_k, t_{k+1}) f^{(n)}(t_1, \dots, t_{k+1}, t_k, \dots, t_n)].$$

Hence, the orthogonal projection of $\mathcal{H}^{\otimes n}$ onto $\text{Ker}(\mathbf{1} + \Psi_k)$, denoted by $E_k^{(n)}$, is given by

$$\begin{aligned} (E_k^{(n)} f^{(n)})(t_1, \dots, t_n) &= \frac{1}{2} \chi_{T_k^{(n)}}(t_1, \dots, t_n) [f^{(n)}(t_1, \dots, t_n) - Q(t_k, t_{k+1}) f^{(n)}(t_1, \dots, t_{k+1}, t_k, \dots, t_n)], \end{aligned}$$

where χ_A denotes the indicator function of a set A . Therefore, the orthogonal projection of $\mathcal{H}^{\otimes n}$ onto $\text{Ker}(\mathbf{1} + \Psi_k)^\perp = \overline{\text{Ran}(\mathbf{1} + \Psi_k)}$, denoted by $F_k^{(n)}$, is given by

$$(F_k^{(n)} f^{(n)})(t_1, \dots, t_n) = \chi_{T^{(n)} \setminus T_k^{(n)}}(t_1, \dots, t_n) f^{(n)}(t_1, \dots, t_n) + \frac{1}{2} \chi_{T_k^{(n)}}(t_1, \dots, t_n) [f^{(n)}(t_1, \dots, t_n) + Q(t_k, t_{k+1}) f^{(n)}(t_1, \dots, t_{k+1}, t_k, \dots, t_n)].$$

Thus, the set $\overline{\text{Ran}(\mathbf{1} + \Psi_k)}$ consists of all functions from $\mathcal{H}^{\otimes n}$ that are Q -quasisymmetric in the t_k, t_{k+1} -variables on the set $T_k^{(n)}$, i.e., for $\sigma^{\otimes n}$ -a.a. $(t_1, \dots, t_n) \in T_k^{(n)}$, equality (7) holds. From here and formula (39), the theorem follows. \square

Proof of Theorem 8. We start with the following lemma.

- Lemma 20.* (i) Let $\mathbf{t}^{(n)} \in T^{(n)}$. Then $\pi \in S_n^1(\mathbf{t}^{(n)})$ if and only if $\pi^{-1} \in S_n^1(\mathbf{t}_\pi^{(n)})$.
 (ii) Let $\mathbf{t}^{(n)} \in T^{(n)}$, let $\pi \in S_n^1(\mathbf{t}^{(n)})$, and let $\nu \in S_n^1(\mathbf{t}_\pi^{(n)})$. Then $\varphi := \pi\nu \in S_n^1(\mathbf{t}^{(n)})$.
 (iii) For each $\mathbf{t}^{(n)} \in T^{(n)}$ and $\pi \in S_n^1(\mathbf{t}^{(n)})$, we have $c_n(\mathbf{t}^{(n)}) = c_n(\mathbf{t}_\pi^{(n)})$.

Proof. (i) By (22),

$$R_{\pi^{-1}}(\mathbf{t}^{(n)}) = \overline{R_\pi(\mathbf{t}_\pi^{(n)})}. \tag{41}$$

From here the statement follows.

- (ii) Assume that $\varphi \notin S_n^1(\mathbf{t}^{(n)})$. Then there exist $i < j$ such that $\varphi^{-1}(i) > \varphi^{-1}(j)$ and $R(t_i, t_j) = 0$. Let us consider two cases.

Case 1: $\pi^{-1}(i) > \pi^{-1}(j)$. But then (22) implies that $R_{\pi^{-1}}(\mathbf{t}^{(n)}) = 0$, hence $\pi \notin S_n^1(\mathbf{t}^{(n)})$, which is a contradiction.

Case 2: $\pi^{-1}(i) < \pi^{-1}(j)$. We then have

$$\nu^{-1}(\pi^{-1}(i)) = \varphi^{-1}(i) > \varphi^{-1}(j) = \nu^{-1}(\pi^{-1}(j)).$$

By (22),

$$R_{\nu^{-1}}(\mathbf{t}_\pi^{(n)}) := \prod_{\substack{1 \leq a < b \leq n \\ \nu^{-1}(a) > \nu^{-1}(b)}} R(t_{\pi(a)}, t_{\pi(b)}).$$

Choose $a = \pi^{-1}(i)$ and $b = \pi^{-1}(j)$. Then $a < b$, $\nu^{-1}(a) > \nu^{-1}(b)$, and

$$R(t_{\pi(a)}, t_{\pi(b)}) = R(t_i, t_j) = 0.$$

Therefore, $R_{\nu^{-1}}(\mathbf{t}_\pi^{(n)}) = 0$, which implies $\nu \notin S_n^1(\mathbf{t}_\pi^{(n)})$. This is again a contradiction. Thus, we must have $\varphi \in S_n^1(\mathbf{t}^{(n)})$.

- (iii) By part (ii), if $\nu \in S_n^1(\mathbf{t}_\pi^{(n)})$, then $\pi\nu \in S_n^1(\mathbf{t}^{(n)})$. Hence, $c_n(\mathbf{t}_\pi^{(n)}) \leq c_n(\mathbf{t}^{(n)})$. On the other hand, by part (i), $\pi^{-1} \in S_n^1(\mathbf{t}_\pi^{(n)})$. Hence, by part (i), if $\mu \in S_n^1(\mathbf{t}^{(n)})$ then $\pi^{-1}\mu \in S_n^1(\mathbf{t}_\pi^{(n)})$. Hence, $c_n(\mathbf{t}^{(n)}) \leq c_n(\mathbf{t}_\pi^{(n)})$. \square

We first show that the operator \mathbb{P}_n is self-adjoint. By (21)–(24), we can write the operator \mathbb{P}_n in the form

$$(\mathbb{P}_n f^{(n)})(\mathbf{t}^{(n)}) = \frac{1}{c_n(\mathbf{t}^{(n)})} \sum_{\pi \in S_n} R_{\pi^{-1}}(\mathbf{t}^{(n)}) f^{(n)}(\mathbf{t}_\pi^{(n)}).$$

Hence using Lemma 20, (iii) and (41), we get, for any $f^{(n)}, g^{(n)} \in \mathcal{H}^{\otimes n}$,

$$\begin{aligned}
 (\mathbb{P}_n f^{(n)}, g^{(n)})_{\mathcal{H}^{\otimes n}} &= \sum_{\pi \in \mathcal{S}_n} \int_{T^{(n)}} \frac{1}{c_n(\mathbf{t}^{(n)})} \overline{R_{\pi^{-1}}(\mathbf{t}^{(n)}) f^{(n)}(\mathbf{t}_\pi^{(n)})} g^{(n)}(\mathbf{t}^{(n)}) \sigma^{\otimes n}(d\mathbf{t}^{(n)}) \\
 &= \sum_{\pi \in \mathcal{S}_n} \int_{T^{(n)}} \frac{1}{c_n(\mathbf{t}_{\pi^{-1}}^{(n)})} \overline{R_{\pi^{-1}}(\mathbf{t}_{\pi^{-1}}^{(n)}) f^{(n)}(\mathbf{t}^{(n)})} g^{(n)}(\mathbf{t}_{\pi^{-1}}^{(n)}) \sigma^{\otimes n}(d\mathbf{t}^{(n)}) \\
 &= \sum_{\pi \in \mathcal{S}_n} \int_{T^{(n)}} \frac{1}{c_n(\mathbf{t}_\pi^{(n)})} \overline{R_\pi(\mathbf{t}_\pi^{(n)}) f^{(n)}(\mathbf{t}^{(n)})} g^{(n)}(\mathbf{t}_\pi^{(n)}) \sigma^{\otimes n}(d\mathbf{t}^{(n)}) \\
 &= \sum_{\pi \in \mathcal{S}_n} \int_{T^{(n)}} \frac{1}{c_n(\mathbf{t}_\pi^{(n)})} R_{\pi^{-1}}(\mathbf{t}^{(n)}) \overline{f^{(n)}(\mathbf{t}^{(n)})} g^{(n)}(\mathbf{t}_\pi^{(n)}) \sigma^{\otimes n}(d\mathbf{t}^{(n)}) \\
 &= \int_{T^{(n)}} \overline{f^{(n)}(\mathbf{t}^{(n)})} \sum_{\pi \in \mathcal{S}_n^1(\mathbf{t}^{(n)})} \frac{1}{c_n(\mathbf{t}_\pi^{(n)})} R_{\pi^{-1}}(\mathbf{t}^{(n)}) g^{(n)}(\mathbf{t}_\pi^{(n)}) \sigma^{\otimes n}(d\mathbf{t}^{(n)}) \\
 &= \int_{T^{(n)}} \overline{f^{(n)}(\mathbf{t}^{(n)})} \frac{1}{c_n(\mathbf{t}^{(n)})} \sum_{\pi \in \mathcal{S}_n^1(\mathbf{t}^{(n)})} R_{\pi^{-1}}(\mathbf{t}^{(n)}) g^{(n)}(\mathbf{t}_\pi^{(n)}) \sigma^{\otimes n}(d\mathbf{t}^{(n)}) \\
 &= (f^{(n)}, \mathbb{P}_n g^{(n)})_{\mathcal{H}^{\otimes n}}.
 \end{aligned} \tag{42}$$

Thus, $\mathbb{P}_n^* = \mathbb{P}_n$.

Our next aim is to prove that $\mathbb{P}_n^2 = \mathbb{P}$, which will imply that \mathbb{P}_n is an orthogonal projection in $\mathcal{H}^{\otimes n}$. For $f^{(n)} \in \mathcal{H}^{\otimes n}$, we have, by Lemma 20, (ii) and (iii),

$$\begin{aligned}
 (\mathbb{P}_n^2 f^{(n)})(\mathbf{t}^{(n)}) &= \frac{1}{c_n(\mathbf{t}^{(n)})} \sum_{\pi \in \mathcal{S}_n^1(\mathbf{t}^{(n)})} \frac{1}{c_n(\mathbf{t}_\pi^{(n)})} \sum_{\nu \in \mathcal{S}_n^1(\mathbf{t}_\pi^{(n)})} (\Phi_\pi \Phi_\nu f^{(n)})(\mathbf{t}^{(n)}) \\
 &= \frac{1}{c_n(\mathbf{t}^{(n)})^2} \sum_{\pi \in \mathcal{S}_n^1(\mathbf{t}^{(n)})} \sum_{\nu \in \mathcal{S}_n^1(\mathbf{t}_\pi^{(n)})} (\Phi_\pi \Phi_\nu f^{(n)})(\mathbf{t}^{(n)}) \\
 &= \frac{1}{c_n(\mathbf{t}^{(n)})^2} \sum_{\varphi \in \mathcal{S}_n^1(\mathbf{t}^{(n)})} \sum_{\substack{\pi \in \mathcal{S}_n^1(\mathbf{t}^{(n)}) \\ \pi\nu = \varphi}} (\Phi_\pi \Phi_\nu f^{(n)})(\mathbf{t}^{(n)}).
 \end{aligned} \tag{43}$$

Let $\varphi \in \mathcal{S}_n^1(\mathbf{t}^{(n)})$ and $\pi \in \mathcal{S}_n^1(\mathbf{t}^{(n)})$. By Lemma 20, (i), we have $\pi^{-1} \in \mathcal{S}_n^1(\mathbf{t}_\pi^{(n)})$. Hence, by Lemma 20, (ii), we get $\nu := \pi^{-1}\varphi \in \mathcal{S}_n^1(\mathbf{t}_\pi^{(n)})$. From here and (43), we get

$$(\mathbb{P}_n^2 f^{(n)})(\mathbf{t}^{(n)}) = \frac{1}{c_n(\mathbf{t}^{(n)})^2} \sum_{\varphi \in \mathcal{S}_n^1(\mathbf{t}^{(n)})} \sum_{\pi \in \mathcal{S}_n^1(\mathbf{t}^{(n)})} (\Phi_\pi \Phi_{\pi^{-1}\varphi} f^{(n)})(\mathbf{t}^{(n)}). \tag{44}$$

Lemma 21. Let $\mathbf{t}^{(n)} \in T^{(n)}$, $\pi \in \mathcal{S}_n^1(\mathbf{t}^{(n)})$, and $\nu \in \mathcal{S}_n^1(\mathbf{t}_\pi^{(n)})$. Then, for each $f^{(n)} \in \mathcal{H}^{\otimes n}$,

$$(\Phi_\pi \Phi_\nu f^{(n)})(\mathbf{t}^{(n)}) = (\Phi_{\pi\nu} f^{(n)})(\mathbf{t}^{(n)}). \tag{45}$$

Proof. We first note that equality (45) explicitly means that

$$R_{\pi^{-1}}(\mathbf{t}^{(n)}) R_{\nu^{-1}}(\mathbf{t}_\pi^{(n)}) f^{(n)}(\mathbf{t}_{\pi\nu}^{(n)}) = R_{\nu^{-1}\pi^{-1}}(\mathbf{t}^{(n)}) f^{(n)}(\mathbf{t}_{\pi\nu}^{(n)}),$$

which is equivalent to the equality

$$R_{\pi^{-1}}(\mathbf{t}^{(n)}) R_{\nu^{-1}}(\mathbf{t}_\pi^{(n)}) = R_{\nu^{-1}\pi^{-1}}(\mathbf{t}^{(n)}). \tag{46}$$

Since $\pi \in \mathcal{S}_n^1(\mathbf{t}^{(n)})$, $\nu \in \mathcal{S}_n^1(\mathbf{t}_\pi^{(n)})$, and $\pi\nu \in \mathcal{S}_n^1(\mathbf{t}^{(n)})$, we have

$$|R_{\pi^{-1}}(\mathbf{t}^{(n)})| = 1, \quad |R_{\nu^{-1}}(\mathbf{t}_\pi^{(n)})| = 1, \quad |R_{\nu^{-1}\pi^{-1}}(\mathbf{t}^{(n)})| = 1. \tag{47}$$

We define a Hermitian function $G : T^{(2)} \rightarrow \mathbb{C}$ by

$$G(s, t) := \begin{cases} R(s, t), & \text{if } |R(s, t)| = 1, \\ 1, & \text{if } R(s, t) = 0. \end{cases} \tag{48}$$

For each $\pi \in S_n$, similarly to the operator Ψ_π defined for the function Q and to the operator Φ_π defined for the function R , we define an operator $\Gamma_\pi : \mathcal{H}^{\otimes n} \rightarrow \mathcal{H}^{\otimes n}$ for the function G . Thus,

$$(\Gamma_\pi f^{(n)})(\mathbf{t}^{(n)}) = G_{\pi^{-1}}(\mathbf{t}^{(n)})f^{(n)}(\mathbf{t}_\pi^{(n)}),$$

where

$$G_\pi(\mathbf{t}^{(n)}) := \prod_{\substack{1 \leq i < j \leq n \\ \pi(i) > \pi(j)}} G(t_i, t_j), \quad \mathbf{t}^{(n)} \in T^{(n)}. \tag{49}$$

For an adjacent transposition $\pi_j = (j, j + 1)$, we denote $\Gamma_j := \Gamma_{\pi_j}$. By Lemma 2, the operators Γ_j satisfy the braid relations. Furthermore, since $|G(s, t)| = 1$ for all $(s, t) \in T^{(2)}$, we get $\Gamma_j^2 = \mathbf{1}$. Using, e.g., Ref. 9, we therefore conclude that the operators Γ_π with $\pi \in S_n$ form a unitary representation of S_n , i.e., for any $\pi, \nu \in S_n$, it holds that $\Gamma_\pi \Gamma_\nu = \Gamma_{\pi\nu}$, and in fact, for all $\mathbf{t}^{(n)} \in T^{(n)}$,

$$G_{\pi^{-1}}(\mathbf{t}^{(n)})G_{\nu^{-1}}(\mathbf{t}_\pi^{(n)}) = G_{\nu^{-1}\pi^{-1}}(\mathbf{t}^{(n)}). \tag{50}$$

But if $\mathbf{t}^{(n)} \in T^{(n)}$, $\pi \in S_n^1(\mathbf{t}^{(n)})$, and $\nu \in S_n^1(\mathbf{t}_\pi^{(n)})$, then formulas (47)–(50) imply (46). □

Now formula (44) and Lemma 21 yield the equality

$$\begin{aligned} (\mathbb{P}_n^2 f^{(n)})(\mathbf{t}^{(n)}) &= \frac{1}{c_n(\mathbf{t}^{(n)})^2} \sum_{\varphi \in S_n^1(\mathbf{t}^{(n)})} \sum_{\pi \in S_n^1(\mathbf{t}_\varphi^{(n)})} (\Phi_\varphi f^{(n)})(\mathbf{t}^{(n)}) \\ &= \frac{1}{c_n(\mathbf{t}^{(n)})} \sum_{\varphi \in S_n^1(\mathbf{t}^{(n)})} (\Phi_\varphi f^{(n)})(\mathbf{t}^{(n)}) = (\mathbb{P}_n f^{(n)})(\mathbf{t}^{(n)}). \end{aligned} \tag{51}$$

Thus, \mathbb{P}_n is an orthogonal projection in $\mathcal{H}^{\otimes n}$.

It remains to prove that $\text{Ran}(\mathbb{P}_n) = \mathcal{F}_n(\mathcal{H})$. Let $f^{(n)} \in \mathcal{F}_n(\mathcal{H})$. Theorem 7 and the construction of the Φ_π operators imply that for $\sigma^{\otimes n}$ -a.a. $\mathbf{t}^{(n)} \in T^{(n)}$ and for each $\pi \in S_n^1(\mathbf{t}^{(n)})$, we have $(\Phi_\pi f^{(n)})(\mathbf{t}^{(n)}) = f^{(n)}(\mathbf{t}^{(n)})$. Hence, by (24), $\mathbb{P}_n f^{(n)} = f^{(n)}$, i.e., $f^{(n)} \in \text{Ran}(\mathbb{P}_n)$.

Finally, we have to prove the inclusion $\text{Ran}(\mathbb{P}_n) \subset \mathcal{F}_n(\mathcal{H})$. This means that for any $f^{(n)} \in \mathcal{H}^{\otimes n}$ and $k \in \{1, \dots, n - 1\}$,

$$(\Phi_k \mathbb{P}_n f^{(n)})(\mathbf{t}^{(n)}) = (\mathbb{P}_n f^{(n)})(\mathbf{t}^{(n)}) \quad \text{for } \sigma^{\otimes n} - \text{a.a. } \mathbf{t}^{(n)} \in T_k^{(n)}. \tag{52}$$

The proof of (52) is similar to the proof of the equality $\mathbb{P}_n^2 = \mathbb{P}$ [formulas (43), (44), and (51)], so we omit it. □

Proof of Corollary 10. We start with the following lemma

Lemma 22. For each $n \in \mathbb{N}$, we have

$$\mathbb{P}_{n+1}(\mathbb{P}_n \otimes \mathbf{1}) = \mathbb{P}_{n+1}, \tag{53}$$

$$\mathbb{P}_{n+1}(\mathbf{1} \otimes \mathbb{P}_n) = \mathbb{P}_{n+1}. \tag{54}$$

Proof. We will only prove equality (53), since the proof of (54) is similar. For a permutation $\nu \in S_n$, we denote by $\nu \otimes \text{id}$ the permutation from S_{n+1} defined by $(\nu \otimes \text{id})(i) := \nu(i)$ for $i \in \{1, \dots, n\}$ and $(\nu \otimes \text{id})(n + 1) := n + 1$. Analogously to the proof of Theorem 8, we get, for any $f^{(n+1)} \in \mathcal{H}^{\otimes(n+1)}$,

$$\begin{aligned} &(\mathbb{P}_{n+1}(\mathbb{P}_n \otimes \mathbf{1})f^{(n+1)})(\mathbf{t}^{(n+1)}) \\ &= \frac{1}{c_{n+1}(\mathbf{t}^{(n+1)})} \sum_{\pi \in S_{n+1}^1(\mathbf{t}^{(n+1)})} \frac{1}{c_n(t_{\pi(1)}, \dots, t_{\pi(n)})} \sum_{\nu \in S_n^1(t_{\pi(1)}, \dots, t_{\pi(n)})} (\Phi_{\pi(\nu \otimes \text{id})} f^{(n+1)})(\mathbf{t}^{(n+1)}) \\ &= \frac{1}{c_{n+1}(\mathbf{t}^{(n+1)})} \sum_{i=1}^{n+1} \sum_{\substack{\varphi \in S_{n+1}^1(\mathbf{t}^{(n+1)}) \\ \varphi(n+1)=i}} \sum_{\substack{\pi \in S_{n+1}^1(\mathbf{t}^{(n+1)}) \\ \pi(n+1)=i \\ \nu \in S_n^1(t_{\pi(1)}, \dots, t_{\pi(n)}) \\ \pi(\nu \otimes \text{id})=\varphi}} \frac{1}{c_n(t_{\pi(1)}, \dots, t_{\pi(n)})} (\Phi_\varphi f^{(n+1)})(\mathbf{t}^{(n+1)}). \end{aligned} \tag{55}$$

Let $\varphi, \pi \in S_{n+1}^1(\mathbf{t}^{(n+1)})$ be such that $\pi(n+1) = \nu(n+1) = i$. Then $\nu' := \pi^{-1}\varphi \in S_{n+1}^1(\mathbf{t}_\pi^{(n+1)})$ and $\nu'(n+1) = n+1$. Therefore, $\nu' = \nu \otimes \text{id}$, where $\nu \in S_n^1(t_{\pi(1)}, \dots, t_{\pi(n)})$. Hence, by (55),

$$\begin{aligned} & (\mathbb{P}_{n+1}(\mathbb{P}_n \otimes \mathbf{1})f^{(n+1)})(\mathbf{t}^{(n+1)}) \\ &= \frac{1}{c_{n+1}(\mathbf{t}^{(n+1)})} \sum_{i=1}^{n+1} \sum_{\substack{\varphi \in S_{n+1}^1(\mathbf{t}^{(n+1)}) \\ \varphi(n+1)=i}} (\Phi_{\varphi}f^{(n+1)})(\mathbf{t}^{(n+1)}) \sum_{\substack{\pi \in S_{n+1}^1(\mathbf{t}^{(n+1)}) \\ \pi(n+1)=i}} \frac{1}{c_n(t_{\pi(1)}, \dots, t_{\pi(n)})}. \end{aligned}$$

Therefore, it is sufficient to prove that for any $\mathbf{t}^{(n+1)} \in T^{(n+1)}$ and $i \in \{1, \dots, n+1\}$,

$$\sum_{\substack{\pi \in S_{n+1}^1(\mathbf{t}^{(n+1)}) \\ \pi(n+1)=i}} \frac{1}{c_n(t_{\pi(1)}, \dots, t_{\pi(n)})} = 1. \tag{56}$$

To this end, we denote

$$S_{n,i}^1(\mathbf{t}^{(n+1)}) := \{\pi \in S_{n+1}^1(\mathbf{t}^{(n+1)}) : \pi(n+1) = i\}$$

and let $c_{n+1,i}(\mathbf{t}^{(n+1)}) := |S_{n,i}^1(\mathbf{t}^{(n+1)})|$. We state that for any $\mathbf{t}^{(n+1)} \in T^{(n+1)}$ and $\pi \in S_{n,i}^1(\mathbf{t}^{(n+1)})$,

$$c_{n+1,i}(\mathbf{t}^{(n+1)}) = c_n(t_{\pi(1)}, \dots, t_{\pi(n)}). \tag{57}$$

Indeed, if $\nu \in S_n^1(t_{\pi(1)}, \dots, t_{\pi(n)})$, then $\nu \otimes \text{id} \in S_{n+1}^1(\mathbf{t}_\pi^{(n+1)})$. Therefore, $\pi(\nu \otimes \text{id}) \in S_{n+1}^1(\mathbf{t}^{(n+1)})$ and

$$(\pi(\nu \otimes \text{id}))(n+1) = \pi(n+1) = i.$$

Hence, $\pi(\nu \otimes \text{id}) \in S_{n+1,i}^1(\mathbf{t}^{(n+1)})$. So $c_n(t_{\pi(1)}, \dots, t_{\pi(n)}) \leq c_{n+1,i}(\mathbf{t}^{(n+1)})$. On the other hand, take any $\varphi \in S_{n+1,i}^1(\mathbf{t}^{(n+1)})$. Let $\nu' := \pi^{-1}\varphi$. As shown above, $\nu' = \nu \otimes \text{id}$, where $\nu \in S_n^1(t_{\pi(1)}, \dots, t_{\pi(n)})$. Hence, $c_{n+1,i}(\mathbf{t}^{(n+1)}) \leq c_n(t_{\pi(1)}, \dots, t_{\pi(n)})$, and formula (57) is proven. Finally, formula (57) implies (56). \square

Lemma 23. For $k \in \mathbb{N}$, we denote by $\mathbf{1}_k$ the identity operator in $\mathcal{H}^{\otimes k}$. Then, for each $n \geq 2$,

$$\mathbb{P}_{n+k}(\mathbb{P}_n \otimes \mathbf{1}_k) = \mathbb{P}_{n+k}, \tag{58}$$

$$\mathbb{P}_{n+k}(\mathbf{1}_k \otimes \mathbb{P}_n) = \mathbb{P}_{n+k}. \tag{59}$$

Proof. We will again only prove the first formula, (58), the proof of (59) being similar. We prove (58) by induction on k . For $k = 1$, formula (58) becomes (53). Let $k \geq 2$ and assume that formula (53) holds for $k - 1$. We then have

$$\begin{aligned} \mathbb{P}_{n+k}(\mathbb{P}_n \otimes \mathbf{1}_k) &= \mathbb{P}_{n+k}(\mathbb{P}_{n+k-1} \otimes \mathbf{1})(\mathbb{P}_n \otimes \mathbf{1}_{k-1} \otimes \mathbf{1}) \\ &= \mathbb{P}_{n+k}[(\mathbb{P}_{n+k-1}(\mathbb{P}_n \otimes \mathbf{1}_{k-1})) \otimes \mathbf{1}] \\ &= \mathbb{P}_{n+k}(\mathbb{P}_{n+k-1} \otimes \mathbf{1}) = \mathbb{P}_{n+k}. \end{aligned} \tag{60}$$

Using Lemma 23, we get, for $n \geq 2$ and $k \in \{1, \dots, n-1\}$,

$$\mathbb{P}_n(\mathbb{P}_k \otimes \mathbb{P}_{n-k}) = \mathbb{P}_n(\mathbb{P}_k \otimes \mathbf{1}_{n-k})(\mathbf{1}_k \otimes \mathbb{P}_{n-k}) = \mathbb{P}_n(\mathbf{1}_k \otimes \mathbb{P}_{n-k}) = \mathbb{P}_n. \tag{61}$$

Proof of Proposition 12. The result below was shown in the proof of Theorem 3.1 in Ref. 8.

Lemma 24 (Ref. 8). Let a bounded linear operator $\mathcal{R}_n : \mathcal{H}^{\otimes n} \rightarrow \mathcal{H}^{\otimes n}$ be defined by

$$\mathcal{R}_n := \mathbf{1}_n + \Psi_1 + \Psi_1\Psi_2 + \dots + \Psi_1\Psi_2 \dots \Psi_{n-1}. \tag{60}$$

Then, for $n \in \mathbb{N}$,

$$(n+1)\mathcal{P}_{n+1} = (\mathbf{1} \otimes \mathcal{P}_n)\mathcal{R}_{n+1}. \tag{61}$$

Analogously to $\mathcal{F}_{\text{fin}}(\mathcal{H})$, we define a linear space $\mathbf{F}_{\text{fin}}(\mathcal{H})$ that consists of finite sequence $(f^{(0)}, f^{(1)}, \dots, f^{(n)}, \dots)$ with $f^{(i)} \in \mathcal{H}^{\otimes i}$. For $h \in \mathcal{H}$, we define a linear operator $A^-(h) : \mathbf{F}_{\text{fin}}(\mathcal{H}) \rightarrow \mathbf{F}_{\text{fin}}(\mathcal{H})$ by setting

$$(A^-(h)f^{(n)})(t_1, \dots, t_{n-1}) := \int_T \overline{h(s)} f^{(n)}(s, t_1, \dots, t_{n-1}) \sigma(ds),$$

for $f^{(n)} \in \mathcal{H}^{\otimes n}$, $n \in \mathbb{N}$, and $A^-(h)(1, 0, 0, \dots) := 0$.

Let $f^{(n)} \in \mathcal{F}_n(\mathcal{H})$, $g^{(n+1)} \in \mathcal{H}^{\otimes(n+1)}$, and $h \in \mathcal{H}$. Then, by (26) and Lemmas 9 and 24, we get

$$\begin{aligned} & (a^+(h)f^{(n)}, \mathbb{P}_{n+1}g^{(n+1)})_{\mathcal{F}_{n+1}(\mathcal{H})} (n+1)! \\ &= (\mathcal{P}_{n+1}\mathbb{P}_{n+1}(h \otimes f^{(n)}), \mathbb{P}_{n+1}g^{(n+1)})_{\mathcal{H}^{\otimes(n+1)}} (n+1)! \\ &= (\mathcal{P}_{n+1}(h \otimes f^{(n)}), g^{(n+1)})_{\mathcal{H}^{\otimes(n+1)}} (n+1)! \\ &= (\mathcal{R}_{n+1}^*(\mathbf{1} \otimes \mathcal{P}_n)(h \otimes f^{(n)}), g^{(n+1)})_{\mathcal{H}^{\otimes(n+1)}} n! \\ &= (h \otimes (\mathcal{P}_n f^{(n)}), \mathcal{R}_{n+1}g^{(n+1)})_{\mathcal{H}^{\otimes(n+1)}} n! \\ &= (\mathcal{P}_n f^{(n)}, A^-(h)\mathcal{R}_{n+1}g^{(n+1)})_{\mathcal{H}^{\otimes n}} n! \\ &= (f^{(n)}, \mathbb{P}_n A^-(h)\mathcal{R}_{n+1}g^{(n+1)})_{\mathcal{F}_n(\mathcal{H})} n!. \end{aligned}$$

From here both formulas (27) and (28) follow. □

Remark 25. Note that formula (28) can now be written in the form

$$a^-(h)\mathbb{P}_n g^{(n)} = \mathbb{P}_{n-1} A^-(h)\mathcal{R}_n g^{(n)}, \tag{62}$$

for $h \in \mathcal{H}$ and $g^{(n)} \in \mathcal{H}^{\otimes n}$.

Proof of Theorem 14. By choosing an orthonormal basis $(e_n)_{n \in \mathbb{N}}$ of \mathcal{H} and writing the infinite matrix of the operator Ψ [see (9)] in terms of the orthonormal basis $(e_n \otimes e_m)_{n,m \in \mathbb{N}}$ of $\mathcal{H}^{\otimes 2}$, one can derive the commutation relation (34) from Sec. 3 of Ref. 8. For the reader's convenience, we will now present a complete proof of this commutation relation without use of an orthonormal basis.

Let $g, h \in \mathcal{H}$ and $f^{(n)} \in \mathcal{F}_n(\mathcal{H})$. By formulas (26) and (62), we get

$$a^-(g)a^+(h)f^{(n)} = \mathbb{P}_n A^-(g)\mathcal{R}_{n+1}(h \otimes f^{(n)}). \tag{63}$$

By (60),

$$\mathcal{R}_{n+1} = \mathbf{1}_{n+1} + \Psi_1(\mathbf{1} \otimes \mathcal{R}_n). \tag{64}$$

Formulas (63) and (64) yield

$$a^-(g)a^+(h)f^{(n)} = (g, h)_{\mathcal{H}} f^{(n)} + \mathbb{P}_n u^{(n)}, \tag{65}$$

where

$$u^{(n)} := A^-(g)\Psi_1(h \otimes (\mathcal{R}_n f^{(n)})).$$

A direct calculation shows that

$$u^{(n)}(t_1, \dots, t_n) = \int_T \sigma(ds) \overline{g(s)} h(t_1) Q(s, t_1) (\mathcal{R}_n f^{(n)})(s, t_2, \dots, t_n). \tag{66}$$

On the other hand, using additionally (54), we get

$$a^+(h)a^-(g)f^{(n)} = \mathbb{P}_n (h \otimes (\mathbb{P}_{n-1} A^-(g)\mathcal{R}_n f^{(n)})) = \mathbb{P}_n v^{(n)}, \tag{67}$$

where

$$v^{(n)} := h \otimes (A^-(g)\mathcal{R}_n f^{(n)}).$$

Note that

$$v^{(n)}(t_1, \dots, t_n) = \int_T \sigma(ds) \overline{g(s)} h(t_1) (\mathcal{R}_n f^{(n)})(s, t_2, \dots, t_n). \tag{68}$$

Formulas (65)–(68) prove (34).

Corollary 10 and formula (31) show that for each $\varphi^{(2)} \in \mathcal{H}^{\otimes 2}$ and $f^{(n)} \in \mathcal{F}_n(\mathcal{H})$,

$$\int_{T^2} \sigma(ds) \sigma(dt) \varphi^{(2)}(s, t) \partial_s^\dagger \partial_t^\dagger f^{(n)} = \mathbb{P}_{n+2}((\mathbb{P}_2 \varphi^{(2)}) \otimes f^{(n)}). \tag{69}$$

By Theorem 7, since $\varphi^{(2)}$ has support in Θ , we get $\mathbb{P}_2 \Psi \varphi^{(2)} = \mathbb{P}_2 \varphi^{(2)}$. Hence, formulas (32) and (69) imply

$$\begin{aligned} \int_{T^2} \sigma(ds) \sigma(dt) \varphi^{(2)}(s, t) \partial_s^\dagger \partial_t^\dagger f^{(n)} &= \int_{T^2} \sigma(ds) \sigma(dt) Q(s, t) \varphi^{(2)}(t, s) \partial_s^\dagger \partial_t^\dagger f^{(n)} \\ &= \int_{T^2} \sigma(ds) \sigma(dt) \varphi^{(2)}(s, t) Q(t, s) \partial_t^\dagger \partial_s^\dagger f^{(n)}, \end{aligned}$$

which gives (35).

Finally, formula (36) is obtained by taking the adjoint operators on the left and right hand sides of formula (35), see (33). □

Proof of Theorem 18. Using Corollary 10, we get, for each $f^{(n)} \in \mathcal{F}_n(\mathcal{H})$ and $h \in \mathcal{H}$,

$$a^+(h)^m f^{(n)} = \mathbb{P}_{m+n}(h^{\otimes m} \otimes f^{(n)}) = \mathbb{P}_{m+n}((\mathbb{P}_m h^{\otimes m}) \otimes f^{(n)}).$$

Hence, it suffices to prove that $\mathbb{P}_m(h^{\otimes m}) = 0$.

Denote by $(e_t)_{t \in T}$ the canonical orthonormal basis in $\mathcal{H} = \ell^2(T \rightarrow \mathbb{C})$, i.e., $e_t(s) = 1$ if $s = t$ and $e_t(s) = 0$ if $s \neq t$. In view of (9), we get

$$\Psi e_s \otimes e_t = Q(t, s) e_t \otimes e_s, \quad (s, t) \in T^2.$$

Note that the operators $(\Psi_\pi)_{\pi \in S_m}$ form a unitary representation of the group S_m , see the proof of Lemma 21. Therefore, for each $k \in \{1, \dots, m - 1\}$, we have $\mathbb{P}_m \Psi_k = \mathbb{P}_m$. Hence, for any $t_1, \dots, t_m \in T$,

$$\mathbb{P}_m(e_{t_1} \otimes \dots \otimes e_{t_{k-1}} \otimes e_{t_{k+1}} \otimes e_{t_k} \otimes e_{t_{k+2}} \otimes \dots \otimes e_{t_m}) = Q(t_k, t_{k+1}) \mathbb{P}_m(e_{t_1} \otimes \dots \otimes e_{t_m}).$$

This implies that

$$\mathbb{P}_m(e_{t_1} \otimes \dots \otimes e_{t_m}) = 0 \quad \text{if } |\{t_1, \dots, t_m\}| < m \tag{70}$$

(i.e., if some index t_i appears twice or more times). Analogously, for any $(t_1, \dots, t_m) \in T^m$ and $\pi \in S_m$,

$$\mathbb{P}_m(e_{t_{\pi(1)}} \otimes \dots \otimes e_{t_{\pi(m)}}) = Q_\pi(t_1, \dots, t_m) \mathbb{P}_m(e_{t_1} \otimes \dots \otimes e_{t_m}). \tag{71}$$

Let $h = \sum_{t \in T} h_t e_t \in \mathcal{H}$. We get, by (70) and (71),

$$\begin{aligned} \mathbb{P}_m h^{\otimes m} &= \sum_{t_1, \dots, t_m \in T} h_{t_1} \dots h_{t_m} \mathbb{P}_m(e_{t_1} \otimes \dots \otimes e_{t_m}) \\ &= \sum_{\substack{t_1, \dots, t_m \in T \\ t_i \neq t_j \text{ if } i \neq j}} h_{t_1} \dots h_{t_m} \mathbb{P}_m(e_{t_1} \otimes \dots \otimes e_{t_m}) \\ &= \sum_{\substack{t_1, \dots, t_m \in T \\ t_1 < t_2 < \dots < t_m}} \sum_{\pi \in S_m} h_{t_{\pi(1)}} \dots h_{t_{\pi(m)}} \mathbb{P}_m(e_{t_{\pi(1)}} \otimes \dots \otimes e_{t_{\pi(m)}}) \\ &= \sum_{\substack{t_1, \dots, t_m \in T \\ t_1 < t_2 < \dots < t_m}} h_{t_1} \dots h_{t_m} \sum_{\pi \in S_m} \mathbb{P}_m(e_{t_{\pi(1)}} \otimes \dots \otimes e_{t_{\pi(m)}}) \\ &= \sum_{\substack{t_1, \dots, t_m \in T \\ t_1 < t_2 < \dots < t_m}} h_{t_1} \dots h_{t_m} \left(\sum_{\pi \in S_m} Q_\pi(t_1, \dots, t_m) \right) \mathbb{P}_m(e_{t_1} \otimes \dots \otimes e_{t_m}), \end{aligned} \tag{72}$$

where we used that $h_{t_{\pi(1)}} \dots h_{t_{\pi(m)}} = h_{t_1} \dots h_{t_m}$ for any permutation $\pi \in S_m$. It can be easily proven by induction on m that for any $t_1, \dots, t_m \in T$ with $t_1 < t_2 < \dots < t_m$, we have

$$\sum_{\pi \in S_m} Q_{\pi}(t_1, \dots, t_m) = [m]_q!. \quad (73)$$

Here we used the notation, for $m \in \mathbb{N}$ and $q \neq 1$,

$$[m]_q! := \prod_{i=1}^m [i]_q, \quad \text{where } [i]_q := 1 + q + q^2 + \dots + q^{i-1} = \frac{1 - q^i}{1 - q}.$$

Since $q^m = 1$, we get $[m]_q! = 0$. Hence, the theorem follows from (72) and (73). \square

ACKNOWLEDGMENTS

The authors acknowledge the financial support for this work by the Polish National Science Center, Grants No. 2012/05/B/ST1/00626 and 2016/21/B/ST1/00628 (M.B. and J.W.). M.B. was partially supported by the MAESTRO Grant No. DEC-2011/02/A/ST1/00119. M.B. and E.L. acknowledge the financial support of the SFB 701 ‘‘Spectral structures and topological methods in mathematics,’’ Bielefeld University.

- ¹ Anshelevich, M., ‘‘Partition-dependent stochastic measures and q -deformed cumulants,’’ *Documenta Math.* **6**, 343–384 (2001), <http://www.math.uiuc.edu/documenta/vol-06/14.html>.
- ² Bożejko, B., ‘‘Deformed Fock spaces, Hecke operators and monotone Fock space of Muraki,’’ *Demonstratio Math.* **45**, 399–413 (2012).
- ³ Bożejko, M., Ejsmont, W., and Hasebe, T., ‘‘Fock space associated to Coxeter groups of type B,’’ *J. Funct. Anal.* **269**, 1769–1795 (2015).
- ⁴ Bożejko, M., Kümmerer, B., and Speicher, R., ‘‘ q -Gaussian processes: Non-commutative and classical aspects,’’ *Commun. Math. Phys.* **185**, 129–154 (1997).
- ⁵ Bożejko, M. and Lytvynov, E., ‘‘Meixner class of non-commutative generalized stochastic processes with freely independent values I. A characterization,’’ *Commun. Math. Phys.* **292**, 99–129 (2009).
- ⁶ Bożejko, M., Lytvynov, E., and Wysoczański, J., ‘‘Noncommutative Lévy processes for generalized (particularly anyon) statistics,’’ *Commun. Math. Phys.* **313**, 535–569 (2012).
- ⁷ Bożejko, M. and Speicher, R., ‘‘An example of a generalized Brownian motion,’’ *Commun. Math. Phys.* **137**, 519–531 (1991).
- ⁸ Bożejko, M. and Speicher, R., ‘‘Completely positive maps on Coxeter groups, deformed commutation relations, and operator spaces,’’ *Math. Ann.* **300**, 97–120 (1994).
- ⁹ Coxeter, H. S. M. and Moser, W. O. J., *Generators and Relations for Discrete Groups*, 4th ed. (Springer-Verlag, Berlin, New York, 1980).
- ¹⁰ Dabrowski, Y., ‘‘A free stochastic partial differential equation,’’ *Ann. Inst. Henri Poincaré Probab. Stat.* **50**, 1404–1455 (2014).
- ¹¹ Dykema, K. and Nica, A., ‘‘On the Fock representation of the q -commutation relations,’’ *J. Reine Angew. Math.* **440**, 201–212 (1993).
- ¹² Frappat, L., Sciarrino, A., Sciuto, S., and Sorba, P., ‘‘Anyonic realizations of the quantum affine Lie algebra $U_q(\widehat{A}_{N-1})$,’’ *Phys. Lett. B* **369**, 313–324 (1996).
- ¹³ Goldin, G. A. and Majid, S., ‘‘On the Fock space for nonrelativistic anyon fields and braided tensor products,’’ *J. Math. Phys.* **45**, 3770–3787 (2004).
- ¹⁴ Goldin, G. A., Menikoff, R., and Sharp, D. H., ‘‘Particle statistics from induced representations of a local current group,’’ *J. Math. Phys.* **21**, 650–664 (1980).
- ¹⁵ Goldin, G. A., Menikoff, R., and Sharp, D. H., ‘‘Representations of a local current algebra in nonsimply connected space and the Aharonov–Bohm effect,’’ *J. Math. Phys.* **22**, 1664–1668 (1981).
- ¹⁶ Goldin, G. and Sharp, D., ‘‘Diffeomorphism groups, anyon fields, and q commutators,’’ *Phys. Rev. Lett.* **76**, 1183–1187 (1996).
- ¹⁷ Guionnet, A. and Shlyakhtenko, D., ‘‘Free monotone transport,’’ *Inventiones Math.* **197**, 613–661 (2014).
- ¹⁸ Jørgensen, P. E. T., Schmitt, L. M., and Werner, R. F., ‘‘ q -canonical commutation relations and stability of the Cuntz algebra,’’ *Pac. J. Math.* **165**, 131–151 (1994).
- ¹⁹ Jørgensen, P. E. T., Schmitt, L. M., and Werner, R. F., ‘‘Positive representations of general commutation relations allowing Wick ordering,’’ *J. Funct. Anal.* **134**, 33–99 (1995).
- ²⁰ Jørgensen, P. E. T., Proskurin, D., and Samoilenko, Y. S., ‘‘The kernel of Fock representations of Wick algebras with braided operator of coefficients,’’ *Pac. J. Math.* **198**, 109–123 (2001).
- ²¹ Kennedy, M. and Nica, A., ‘‘Exactness of the Fock space representations of the q -commutation relations,’’ *Commun. Math. Phys.* **308**, 115–132 (2011).
- ²² Królak, L., ‘‘Wick product for commutation relations connected with Yang–Baxter operators and new constructions of factors,’’ *Commun. Math. Phys.* **210**, 685–701 (2000).
- ²³ Lerda, A. and Sciuto, S., ‘‘Anyons and quantum groups,’’ *Nucl. Phys. B* **401**, 613–643 (1993).

- ²⁴ Liguori, A. and Mintchev, M., “Fock representations of quantum fields with generalized statistics,” *Commun. Math. Phys.* **169**, 635–652 (1995).
- ²⁵ Lust-Piquard, F., “Riesz transform on deformed Fock space,” *Commun. Math. Phys.* **205**, 519–549 (1999).
- ²⁶ Lytvynov, E., “Gauge-invariant quasi-free states on the algebra of the anyon commutation relations,” *Comm. Math. Phys.* **351**, 653–687 (2017).
- ²⁷ Merberg, A., “Some results on continuous deformed free group factors,” e-print [arXiv:1203.2176](https://arxiv.org/abs/1203.2176).
- ²⁸ Nelson, B. and Zeng, Q., “An application of free transport to mixed q -Gaussian algebras,” *Proc. Amer. Math. Soc.* **144**, 4357–4366 (2016).
- ²⁹ Nelson, B. and Zeng, Q., “Free monotone transport for infinite variables,” *Int. Math. Res. Notices* rnx060 (2017).
- ³⁰ Nica, A. and Speicher, R., *Lectures on the Combinatorics of Free Probability*, London Mathematical Society Lecture Note Series Vol. 335 (Cambridge University Press, London, 2006).
- ³¹ Nou, A., “Non-injectivity of the q -deformed von Neumann algebra,” *Math. Ann.* **330**, 17–38 (2004).
- ³² Proskurin, D. P and Samoilenko, Yu. S., “Deformations of canonical commutation relations, their $*$ -representations, and enveloping C^* -algebras,” *J. Math. Sci.* **164**, 648–657 (2010).
- ³³ Ricard, E., “Factoriality of q -Gaussian von Neumann algebras,” *Commun. Math. Phys.* **257**, 659–665 (2005).
- ³⁴ Shlyakhtenko, D., “Some estimates for non-microstates free entropy dimension with applications to q -semicircular families,” *Int. Math. Res. Not.* **2004**, no. 51, 2757–2772.
- ³⁵ Śniady, P., “Factoriality of Bożejko–Speicher von Neumann algebras,” *Commun. Math. Phys.* **246**, 561–567 (2004).
- ³⁶ Speicher, R., “Generalized statistics of macroscopic fields,” *Lett. Math. Phys.* **27**, 97–104 (1993).