



Swansea University
Prifysgol Abertawe



Cronfa - Swansea University Open Access Repository

This is an author produced version of a paper published in:

Injury

Cronfa URL for this paper:

<http://cronfa.swan.ac.uk/Record/cronfa37128>

Paper:

Eftaxiopoulou, T., Barnett-Vanes, A., Arora, H., Macdonald, W., Nguyen, T., Itadani, M., Sharrock, A., Britzman, D., Proud, W., et. al. (2016). Prolonged but not short-duration blast waves elicit acute inflammation in a rodent model of primary blast limb trauma. *Injury*, 47(3), 625-632.



<http://dx.doi.org/10.1016/j.injury.2016.01.017>

This item is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Copies of full text items may be used or reproduced in any format or medium, without prior permission for personal research or study, educational or non-commercial purposes only. The copyright for any work remains with the original author unless otherwise specified. The full-text must not be sold in any format or medium without the formal permission of the copyright holder.


Permission for multiple reproductions should be obtained from the original author.

Authors are personally responsible for adhering to copyright and publisher restrictions when uploading content to the repository.


<http://www.swansea.ac.uk/library/researchsupport/ris-support/>



(A)



(B)



(C)

Figure 3: (A) Shows the number of circulating neutrophils and CD43Lo/His48Hi monocytes observed at 6hrs and 24hrs for all groups. Representative flow cytometry plots are shown for Sham and Group III at 6hrs. (B) Proportions of circulating CD43Hi/His48Int-Lo monocytes, NK Cells and T Cells were examined. (C) Representative bar graphs showing the concentration of circulating IL-6 and CXCL1 in the plasma. Data are n=4-6 from 2-3 separate experiments, *p<0.05, **p<0.01.