



Swansea University
Prifysgol Abertawe



Cronfa - Swansea University Open Access Repository

This is an author produced version of a paper published in:

Genomics Data

Cronfa URL for this paper:

<http://cronfa.swan.ac.uk/Record/cronfa38371>

Paper:

Ascunce, M., Huguet-Tapia, J., Braun, E., Ortiz-Urquiza, A., Keyhani, N. & Goss, E. (2016). Whole genome sequence of the emerging oomycete pathogen *Pythium insidiosum* strain CDC-B5653 isolated from an infected human in the USA. *Genomics Data*, 7, 60-61.

<http://dx.doi.org/10.1016/j.gdata.2015.11.019>

Released under the terms of a Creative Commons Attribution 4.0 International license (CC-BY).

This item is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Copies of full text items may be used or reproduced in any format or medium, without prior permission for personal research or study, educational or non-commercial purposes only. The copyright for any work remains with the original author unless otherwise specified. The full-text must not be sold in any format or medium without the formal permission of the copyright holder.

Permission for multiple reproductions should be obtained from the original author.

Authors are personally responsible for adhering to copyright and publisher restrictions when uploading content to the repository.

<http://www.swansea.ac.uk/library/researchsupport/ris-support/>



Whole genome sequence of the emerging oomycete pathogen *Pythium insidiosum* strain CDC-B5653 isolated from an infected human in the USA



Marina S. Ascunce^{a,b,1}, Jose C. Huguet-Tapia^{a,1}, Edward L. Braun^c, Almudena Ortiz-Urquiza^d, Nemat O. Keyhani^d, Erica M. Goss^{a,b,*}

^a Emerging Pathogens Institute, University of Florida, Gainesville, FL, United States

^b Department of Plant Pathology, University of Florida, Gainesville, FL, United States

^c Department of Biology and Genetics Institute, University of Florida, Gainesville, FL, United States

^d Department of Microbiology and Cell Science, University of Florida, Gainesville, FL, United States

ARTICLE INFO

Article history:

Received 12 November 2015

Accepted 22 November 2015

Available online 23 November 2015

Keywords:

Oomycete

Pythium insidiosum

Pythiosis

Human emerging pathogen

Genome sequencing

ABSTRACT

Pythium insidiosum ATCC 200269 strain CDC-B5653, an isolate from necrotizing lesions on the mouth and eye of a 2-year-old boy in Memphis, Tennessee, USA, was sequenced using a combination of Illumina MiSeq (300 bp paired-end, 14 millions reads) and PacBio (10 Kb fragment library, 356,001 reads). The sequencing data were assembled using SPAdes version 3.1.0, yielding a total genome size of 45.6 Mb contained in 8992 contigs, N₅₀ of 13 Kb, 57% G + C content, and 17,867 putative protein-coding genes. This Whole Genome Shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession JRRHR00000000.

© 2015 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Specifications	
Organism/cell line/tissue	<i>Pythium insidiosum</i> strain CDC-B5653
Sex	Not applicable
Sequencer or array type	Illumina MiSeq and PacBio
Data format	Assembled
Experimental factors	CDC sample originally isolated from necrotizing lesions on the mouth and eye of a 2-year-old boy
Experimental features	Whole genome shotgun sequencing followed by genome assembly and gene description
Consent	Not applicable
Sample source location	ATCC 200269

1. Direct link to deposited data

<http://www.ncbi.nlm.nih.gov/nuccore/JRRHR00000000.1>.

2. Experimental design, materials and methods

The oomycete genus *Pythium* comprises more than 250 described species [1], most of which are saprobes or facultative plant pathogens

* Corresponding author at: Emerging Pathogens Institute, University of Florida, Gainesville, FL, United States.

E-mail address: emgoss@ufl.edu (E.M. Goss).

¹ M.S.A. and J.C. H-T. contributed equally to this work.

that cause seed rot and damping-off, root, stem and fruit rot, foliar blight, and postharvest decay [2]. *P. insidiosum* is the only *Pythium* species that causes disease in mammals. It is the causal agent of pythiosis, a deadly disease of horses, dogs, and other mammals in tropical and subtropical regions [3,4]. Pythiosis also affects humans, and was first reported in Thailand in 1985 [5,6].

Whole genome sequencing was applied to *P. insidiosum* ATCC 200269 strain CDC-B5653, which was originally isolated from necrotizing lesions on the mouth and eye of a 2-year-old boy in Memphis, Tennessee, USA. A combination of Illumina MiSeq (300 bp paired-end, 14 millions reads) and PacBio (10 Kb fragment library, 356,001 reads) sequencing data were used to assemble the genome using SPAdes version 3.1.0 [7], yielding a total size of 45.6 Mb contained in 8992 contigs, N₅₀ of 13 Kb, maximum contig length of 148 Kb, and 57% G + C content. We used Augustus version 3.0.1 [8] to predict genes *ab initio*, using a gene model previously described for *Pythium* [9]. This genome contains 225 tRNA and 17,867 putative protein-coding genes. To create a representative set of orthologous groups for *P. insidiosum* and its closest relatives, genomes from the following seven *Pythium* species were included: *P. ultimum* var. *ultimum*, *P. arrhenomanes*, *P. irregulare*, *P. aphanidermatum*, *P. iwayamai*, *P. ultimum* var. *sporangiferum*, and *P. vexans* (now *Phytophythium vexans* [10]). These genomes were downloaded from the *Pythium* Genome Database (<http://pythium.plantbiology.msu.edu/download.shtml>) [9,11]. Reciprocal BLAST analysis on all genomes indicated that *P. insidiosum* shares 5922 unique

orthologous genes with the other *Pythium* genomes and has 649 taxon specific genes. *P. insidiosum* shares more orthologs (233) with *P. aphanidermatum* and *P. arrhenomanes* than with the remaining *Pythium* species. These findings indicate that the three species are evolutionarily close to each other, which is consistent with estimates of *Pythium* phylogeny based on ITS sequences [1]. Further analysis will examine genes and gene families that distinguish the *P. insidiosum* genome from those of plant other pathogenic oomycetes.

Acknowledgments

We wish to thank Jackson Presser for technical support and Jeff Rollins for guidance and laboratory resources. This research was funded by the University of Florida Research Opportunity Incentive Seed Fund from the UF Office of Research, the UF Emerging Pathogens Institute, and the UF IFAS Department of Plant Pathology.

References

- [1] C.A. Lévesque, A.W.A.M. de Cock, Molecular phylogeny and taxonomy of the genus *Pythium*. *Mycol. Res.* 108 (2004) 1363–1383.
- [2] A.J. Van der Plaats-Niterink, Monograph of the genus *Pythium*. *Stud. Mycol.* 21 (1981) 1–242.
- [3] A.W. De Cock, L. Mendoza, A.A. Padhye, L. Ajello, L. Kaufman, *Pythium insidiosum* sp. nov., the etiologic agent of pythiosis. *J. Clin. Microbiol.* 25 (1987) 344–349.
- [4] L. Mendoza, *Pythium insidiosum* and mammalian hosts. in: K. Lamour, S. Kamoun (Eds.), *Oomycete Genetics and Genomics: Diversity, Interactions, and Research Tools*, in: *Oomycete Genet. Genomics Divers. Interact. Res. Tools*, John Wiley & Sons, Inc., Hoboken, NJ 2009, pp. 387–405.
- [5] M. Thianprasit, Fungal infection in Thailand. *Jpn. J. Dermatol.* 96 (1986) 1343–1345.
- [6] T. Krajaejun, B. Sathapatayavongs, R. Prachartam, P. Nitiyanant, P. Leelachaikul, W. Wanachiwanawin, et al., Clinical and epidemiological analyses of human pythiosis in Thailand. *Clin. Infect. Dis.* 43 (2006) 569–576, <http://dx.doi.org/10.1086/506353>.
- [7] S. Nurk, A. Bankevich, D. Antipov, A. Gurevich, A. Korobeynikov, A. Lapidus, et al., Assembling Genomes and Mini-Metagenomes from Highly Chimeric Reads. in: M. Deng, R. Jiang, F. Sun, X. Zhang (Eds.), *Res. Comput. Mol. Biol.* Springer Berlin, Heidelberg 2013, pp. 158–170, http://dx.doi.org/10.1007/978-3-642-37195-0_13.
- [8] M. Stanke, M. Diekhans, R. Baertsch, D. Haussler, Using native and syntenically mapped cDNA alignments to improve de novo gene finding. *Bioinforma. Oxf. Engl.* 24 (2008) 637–644, <http://dx.doi.org/10.1093/bioinformatics/btn013>.
- [9] C.A. Lévesque, H. Brouwer, L. Cano, J.P. Hamilton, C. Holt, E. Huitema, et al., Genome sequence of the necrotrophic plant pathogen *Pythium ultimum* reveals original pathogenicity mechanisms and effector repertoire. *Genome Biol.* 11 (2010) R73, <http://dx.doi.org/10.1186/gb-2010-11-7-r73>.
- [10] A.W.A.M. de Cock, A.M. Lodhi, T.L. Rintoul, K. Bala, G.P. Robideau, Z.G. Abad, et al., *Phytopythium*: molecular phylogeny and systematics. *Persoonia Mol. Phylogeny Evol. Fungi.* 34 (2015) 25–39, <http://dx.doi.org/10.3767/003158515X685382>.
- [11] B.N. Adhikari, J.P. Hamilton, M.M. Zerillo, N. Tisserat, C.A. Lévesque, C.R. Buell, Comparative genomics reveals insight into virulence strategies of plant pathogenic oomycetes. *PLoS One* 8 (2013), e75072, <http://dx.doi.org/10.1371/journal.pone.0075072>.