



Swansea University
Prifysgol Abertawe



Cronfa - Swansea University Open Access Repository

This is an author produced version of a paper published in:
Journal of Science and Medicine in Sport

Cronfa URL for this paper:
<http://cronfa.swan.ac.uk/Record/cronfa38856>

Paper:

Williams, N., Russell, M., Cook, C. & Kilduff, L. (2018). The effect of lower limb occlusion on recovery following sprint exercise in academy rugby players. *Journal of Science and Medicine in Sport*
<http://dx.doi.org/10.1016/j.jsams.2018.02.012>

This item is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Copies of full text items may be used or reproduced in any format or medium, without prior permission for personal research or study, educational or non-commercial purposes only. The copyright for any work remains with the original author unless otherwise specified. The full-text must not be sold in any format or medium without the formal permission of the copyright holder.

Permission for multiple reproductions should be obtained from the original author.

Authors are personally responsible for adhering to copyright and publisher restrictions when uploading content to the repository.

<http://www.swansea.ac.uk/library/researchsupport/ris-support/>

Accepted Manuscript

Title: The effect of lower limb occlusion on recovery following sprint exercise in academy rugby players

Authors: N. Williams, M. Russell, C.J. Cook, L.P. Kilduff

PII: S1440-2440(18)30064-1

DOI: <https://doi.org/10.1016/j.jsams.2018.02.012>

Reference: JSAMS 1818

To appear in: *Journal of Science and Medicine in Sport*

Received date: 6-10-2017

Revised date: 4-1-2018

Accepted date: 22-2-2018

Please cite this article as: Williams N, Russell M, Cook CJ, Kilduff L.P. The effect of lower limb occlusion on recovery following sprint exercise in academy rugby players. *Journal of Science and Medicine in Sport* <https://doi.org/10.1016/j.jsams.2018.02.012>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Title: The effect of lower limb occlusion on recovery following sprint exercise in academy rugby players

Authors: Williams, N.^{a, b} Russell, M.^c, Cook, C.J.^{d, e}, Kilduff, L.P.^{a, d}

^a Applied Sports Technology Exercise and Medicine Research Centre (A-STEM), Swansea University, Swansea, United Kingdom

^b Sport Wales, Welsh Institute of Sport, Sophia Gardens, Cardiff, United Kingdom

^c School of Social and Health Sciences, Leeds Trinity University, Leeds, United Kingdom

^d Welsh Institute for Performance Solutions

^e University of Canberra Research Institute for Sport and Exercise, University of Canberra, Australia

Corresponding Author: Liam P.Kilduff

L.Kilduff@swansea.ac.uk

Article type: Original investigation

Abstract word count: 249

Manuscript word count: 2936

Figures: 2

Abstract

Objectives: The effects of vascular occlusion on recovery of physiological and neuromuscular markers over 24h, and hormonal reactivity to subsequent exercise were investigated.

Design: Counterbalanced, randomised, crossover

Methods: Academy rugby players ($n=24$) completed six 50-m sprints (five-min inter-set recovery) before occlusion cuff application (thighs) and intermittent inflation to 171-266

mmHg (Recovery) or 15 mmHg (Con) for 12-min (two sets, three-min repetitions, three-min non-occluded reperfusion). Countermovement jumps, blood (lactate, creatine kinase), saliva (testosterone, cortisol), and perceptual (soreness, recovery) responses were measured before (baseline) and after (post, +2h, +24h) sprinting. Saliva was sampled after a 30-min resistance exercise session performed 24h after sprinting.

Results: Although sprinting (total: 40.0 ± 2.8 s, $p=0.238$; average: 6.7 ± 0.5 s, $p=0.674$) influenced creatine kinase ($p<0.001$, $+457.1 \pm 327.3 \mu\text{L}^{-1}$, at 24h), lactate ($p<0.001$, $6.8 \pm 2.3 \text{ mmol}\cdot\text{L}^{-1}$, post), testosterone ($p<0.001$, $-55.9 \pm 63.2 \text{ pg}\cdot\text{ml}^{-1}$, at 2h) and cortisol ($p<0.001$, $-0.3 \pm 0.3 \mu\text{g}\cdot\text{dl}^{-1}$, at 2h) concentrations, countermovement jump power output ($p<0.001$, -409.6 ± 310.1 W; -5.4 ± 3.4 cm, post), perceived recovery ($p<0.001$, -3.0 ± 2.3 , post), and muscle soreness ($p<0.001$; 1.5 ± 1.1 , at 24h), vascular occlusion had no effect (all $p>0.05$) on recovery. In response to subsequent exercise performed 24h after vascular occlusion, testosterone increased pre-to-post-exercise (Recovery: $p=0.031$, $21.6 \pm 44.9 \text{ pg}\cdot\text{ml}^{-1}$; Con: $p=0.178$, $10.6 \pm 36.6 \text{ pg}\cdot\text{ml}^{-1}$) however $\Delta\text{testosterone}$ was not significantly different ($p=0.109$) between conditions.

Conclusions: Vascular occlusion had no effect on physiological or neuromuscular markers 2h or 24h after sprinting or in response to a physical stress test.

Keywords: Occlusion, sprint, hormonal reactivity

Introduction

Physical and metabolic disturbances result from team sport match-play¹. Accordingly, various measures are applied to indicate the presence of exercise-induced muscle damage (EIMD) and the efficacy of recovery interventions². Elevated Creatine Kinase (CK) concentrations³, disruption in the hormonal milieu (i.e., testosterone; T, cortisol; C)⁴, and impairments in neuromuscular function (NMF) occur post-match⁵; with perturbations

occurring for at least 48h^{3,5}. The use of hormonal responses, such as changes in T and C indicate anabolic and catabolic balance⁶ and thus readiness-to-train or competition-preparedness, is an emerging concept in the fatigue-recovery paradigm⁷.

As short-term post-match fatigue impairs subsequent performance, recovery strategies (e.g. cold-water immersion and active recovery) are an integral component of weekly training practices that have been extensively investigated (for review see⁸). More recently, vascular occlusion, the use of blood pressure cuffs applied on specific limbs to restrict blood-flow, has been suggested as a recovery strategy^{9,10}. While exact mechanisms are unclear, vascular occlusion is purported to elevate adenosine concentrations and activation of adenosine triphosphate (ATP) sensitive potassium channels (K_{ATP}), increasing blood flow¹¹ and benefiting oxygen and nutrient delivery via vasculature dilation; a response likely exaggerated during reperfusion, possibly improving substrate re-synthesis¹¹. Alternatively, attenuated inflammatory responses¹⁰ and reduction of muscle oedema and intramuscular pressure decrease nociceptor stimulation, potentially reducing muscle soreness¹².

Unfortunately, the evidence for vascular occlusion as an effective post-exercise recovery modality is currently inconsistent. Two investigations identified improved recovery^{10,11} whereas others^{13,14,15} disagree. Methodological differences exist when implementing vascular occlusion, for example standardised cuff pressures have been implemented despite recommendations regarding individualised application relative to thigh girth and resting blood measurements¹⁶. Similarly, inconsistent timings of recovery assessments (i.e., 1-72h) post-occlusion exist between studies. Previous research has also required players to remain rested for the duration of post-exercise recovery; however, this has limited application to applied practice where football or rugby players are frequently required to play multiple games within a week (i.e., <72h separating games), and train when complete physical recovery may not be achieved¹⁷. Accordingly, identification of the physiological response to a subsequent physical

stressor may denote if players are adequately recovered to return to training. Notably, T is a stress biomarker; consequently, the monitoring of T in response to a physical stress test could provide information on readiness to train/compete¹⁸.

The primary aim of the study was therefore to investigate the effects of post-exercise vascular occlusion (using individualised cuff pressures) on recovery (2h and 24h) of physiological and performance markers following maximal sprint exercise whilst also considering the hormonal reactivity to a subsequent exercise challenge performed at 24h. It was hypothesised that vascular occlusion implemented post-exercise would facilitate the recovery of biochemical, neuromuscular and hormonal markers measured after 24h.

Method

Following institutional ethical approval and informed written consent, 24 male Academy rugby union players (age: 21.8 ± 3.0 y, mass: 96.9 ± 10.1 kg, stature: 1.85 ± 0.09 m) participated in the study during pre-season (1–2 sessions per day 4–5 days a week; strength, power, speed training). All participants were informed of the experimental procedures, the purpose of the study, and possible risks.

Participants attended the testing venue four times. Two main trials (Vascular occlusion: Recovery; Control: Con), seven days apart, were completed on an indoor 3G surface (temperature: 20°C; humidity: 41%) in a randomised, counter-balanced, crossover design. Measurement timings were consistent between trials to limit circadian variation, and participants refrained from alcohol and intense physical exercise in the 24h preceding trials.

On arrival for main trials, participants rested for 10-min before recording blood pressure (Omron Healthcare, Europe; systolic >140 mmHg and diastolic >90 mmHg precluded further study involvement). Thigh girth, physiological (capillary blood and saliva) and perceptual

(soreness and recovery) assessments followed. After a 10-min standardised warm-up, participants performed two maximal countermovement jumps (CMJ) separated by 90 s (portable force platform: Type 92866AA, Kistler, Germany) to assess NMF^{5,19}. A further 10-min warm-up followed (20 m dynamic exercises and accelerations, two-50 m sprints at 80% and 100% effort) with five-min of enforced rest before six-50 m (each separated by five-min rest) timed sprints (Brower Timing System, Salt Lake City, Utah, USA) were performed to induce muscle damage^{20,21} (Figure 1). As per pre-exercise instructions, maximal effort was required across all six sprints, but no encouragement was provided during exercise. Participants and coaches were blinded from sprint timings and feedback was not provided until all trials were completed. Average, and cumulative sprint times were recorded for the six sprints.

Immediately post-exercise, baseline measures were repeated before occlusion cuffs were applied to the proximal point of the thighs while participants lay supine. The cuff (11 cm; Sports Rehab Tourniquet, Sportsrehab) was manually inflated to 15 mmHg (Con), reflecting previous research⁹, or to 60% of individually calculated pressures (171-266 mmHg; Recovery), determined from thigh girth and blood pressure measurements¹⁶. Cuffs were applied for a total of 12-min (two cycles of three-min occlusion, three-min reperfusion)⁹ as reports suggest that three-min cycles of occlusion fulfil the duration threshold and a total ischemic stimulus of at least four-min is required to elicit a protective effect in human myocardium, irrespective of the number of ischemic cycles²². After 2h and 24h baseline measures were repeated; timings which are consistent with previous research^{20,21} and represent the duration between competition and return to training.

To assess hormonal reactivity to a subsequent exercise stimulus performed after 24h, saliva was collected five-min before and immediately after a 30-min physical stress test (three sets of power cleans and back squats, four sets of bench press and bench pull at relative loads of

60-85% 1RM; one and three-min rest between sets and exercises, respectively). Participants were accustomed to the resistance exercise and testing procedures employed and these protocols were sufficient to elicit a stress response¹⁸. Session supervisors provided technical support only and were not aware of the condition that players were in. Feedback was not provided regarding within-session performance.

***** INSERT FIGURE 1 NEAR HERE *****

Saliva collection required passive drooling (~2 ml) into a sterile vial (SalivaBio, Salimetrics LLC, USA) after refraining from brushing of teeth, drinking hot fluids, or eating hard foods 2h beforehand. All samples were stored (-15°C) immediately after collection and transferred to -80°C within 4h of collection. Post thawing, centrifugation (Micro Centaur, MSE, London, United Kingdom; five-min at 3000 revolutions·min⁻¹) preceded duplicate analysis of T and C concentrations due to known reliable reflections of gonadal function²³ using indirect enzyme-linked immunosorbent assay (ELISA) kits (Salimetrics Europe Ltd., Suffolk, U.K.). The lowest detection limits for T and C were 6.1 pg·ml⁻¹ and 0.012 µg·dl⁻¹ respectively, and inter-assay CV values were <10% in both cases.

Participants provided a 20 µL fingertip capillary blood sample (analysed retrospectively for lactate concentrations; Biosen C-Line Clinic, EKF Diagnostic GmgH, Barleben, Germany). Additionally, 120 µL sample was collected, immediately centrifuged (3000 revolutions·min⁻¹ for 10-min; Labofuge 400R, Kendro Laboratories, Germany) for the extraction of plasma, and stored (-80°C) until later analysis. Plasma samples thawed before 6 µL was used for CK analysis (automated analyser; ABX Pentra 400, Horiba ABX, Montpellier, France). Sample testing was carried out in duplicate, intra sample CV values were <2.0%.

Perceived lower body muscle soreness was assessed using a 7-point Likert scale ranging from zero (complete absence of soreness) to six (severe pain limiting movement) which is reliable

and valid²⁴. Perception of recovery status was assessed using a 11-point likert scale²⁵ from zero (very poorly recovered/extremely tired) to 10 (very well recovered) which reflects changes in total sprint time relative to prior exercise²¹. Participants were familiar with the scales and were asked to base scores on perceived soreness during normal movement and were alone when recording scores to reduce influences of peers.

Assessment of CMJ was completed on a portable force platform (Kistler instrument Ltd., Farnborough, UK) sampling at 1000 Hz. Peak power output (PPO) of the lower body was calculated as previously described¹⁹. The vertical component of the ground reaction force and participants' body weight were used to determine instantaneous velocity and displacement of the centre of gravity¹⁹. Instantaneous power output was determined using Equation 1 and the highest value produced was deemed PPO. Jump height (JH) was defined as the difference in vertical displacement of the center of gravity between take-off (toes leave the force plate) and maximum displacement¹⁹.

Equation 1:

Power (W) = Vertical ground reaction force (N) x vertical velocity of centre of gravity (m.s.⁻¹)

Statistical analysis

All data is presented as mean \pm standard deviation (SD). Following confirmation of parametric assumptions, multivariate analysis of variance (MANOVA) with Bonferroni adjustment assessed between-trial differences for variables with multiple time points per trial (i.e. T, C, T:C ratio, perception muscle soreness and recovery, CK, blood lactate, PPO and JH). A one-way ANOVA was performed to assess between-trial differences in response to the physical stress test (T, C, T:C ratio and perception muscle soreness). Paired samples t-tests were performed for between-trial comparisons of data expressed over a single time point within a trial (i.e. mean and total sprint times, T and C pre-and post-stress test). Statistical

analyses were carried out using SPSS (SPSS Chicago, IL) with significance being accepted at $p \leq 0.05$.

Results

There were no significant differences between conditions for total (Recovery: 39.86 ± 2.87 s; Con: 40.26 ± 2.77 s, $p=0.238$) or average (Recovery: 6.69 ± 0.47 s; Con: 6.71 ± 0.46 s, $p=0.674$) sprint times. The Δ lactate concentrations from pre-to-post sprinting showed no significant difference between conditions (Recovery: 6.88 ± 2.53 $\text{mmol}\cdot\text{l}^{-1}$, $85 \pm 6\%$; Con: 6.76 ± 2.04 $\text{mmol}\cdot\text{l}^{-1}$, $86 \pm 5\%$, $p=0.807$, Figure 2).

There was a significant time effect for CK ($F_{(1,48)}=72.928$, $p<0.001$, Figure 2) with increases at 24h compared to pre-sprints in both Recovery (408.19 ± 291.45 $\mu\cdot\text{L}^{-1}$, $55 \pm 33\%$, $p<0.001$) and Con (506.02 ± 359.14 $\mu\cdot\text{L}^{-1}$, $56 \pm 34\%$ $p<0.001$). However, there was no significant interaction effects between condition and time ($F_{(1,48)}=1.157$, $p=0.293$).

***** INSERT FIGURE 2 NEAR HERE *****

Muscle soreness ($F_{(2,95)}=7.714$, $p<0.001$) and perception of recovery ($F_{(2,88)}=70.931$, $p<0.001$) were affected by sprinting, with the greatest change in muscle soreness occurring 24h post exercise (Recovery: 1.5 ± 1.0 ; Con: 1.6 ± 1.1). There was no significant interaction effects between time and condition for muscle soreness ($F_{(2,95)}=0.009$, $p=0.993$) or perception of recovery ($F_{(2,88)}=0.158$, $p=0.924$).

Sprint exercise affected PPO ($F_{(2,96)}=42.141$, $p<0.001$) and JH ($F_{(2,82)}=58.353$, $p<0.001$) with PPO (Recovery: 417.74 ± 293.09 W, $8 \pm 4\%$, $p<0.001$; Con: 401.46 ± 332.69 W, $7 \pm 5\%$, $p<0.001$) and JH (Recovery: 5.49 ± 3.24 cm, $13 \pm 7\%$, $p<0.001$; Con: 5.38 ± 3.66 cm, $13 \pm 8\%$,

$p < 0.001$) decreasing post-sprints (Figure 3). No further timing effects and no effect of Recovery on PPO ($F_{(2, 96)} = 0.304$, $p = 0.757$) or JH ($F_{(2, 82)} = 0.304$, $p = 0.436$) occurred.

Sprinting increased T ($F_{(2, 100)} = 20.127$, $p < 0.001$, Figure 2), T:C ratio ($F_{(2, 95)} = 19.200$, $p < 0.001$) and decreased C ($F_{(2, 89)} = 32.651$, $p < 0.001$, Figure 2). However, condition did not affect the recovery of T ($F_{(2, 100)} = 2.159$, $p = 0.114$), C ($F_{(2, 89)} = 0.640$, $p = 0.531$) or T:C ratio ($F_{(2, 95)} = 0.299$, $p = 0.759$).

Testosterone significantly increased in response to the physical stress test in the Recovery ($+21.58 \pm 44.90$ $\text{pg}\cdot\text{ml}^{-1}$, $7 \pm 17\%$, $p = 0.031$) but not the Con ($+10.62 \pm 36.57$ $\text{pg}\cdot\text{ml}^{-1}$, $4 \pm 13\%$, $p = 0.178$) with no differences in baseline values between conditions ($p = 0.232$); however, ΔT was similar between conditions ($p = 0.109$). Cortisol declined over time ($F_{(1, 46)} = 7.806$, $p < 0.001$), pre-to post physical stress test (Recovery: -0.14 ± 0.23 $\mu\text{g}\cdot\text{dl}^{-1}$, $38 \pm 72\%$, $p = 0.007$; Con: -0.17 ± 0.27 $\mu\text{g}\cdot\text{dl}^{-1}$, $50 \pm 94\%$, $p = 0.006$), with similar results for T:C ratio ($F_{(1, 46)} = 29.836$, $p < 0.001$). Recovery had no impact on hormonal response as no differences were observed between conditions (T; $p = 0.226$, C; $p = 0.679$, T:C; $p = 0.421$).

***** INSERT FIGURE 3 NEAR HERE *****

Discussion

This study aimed to investigate the effects of individualised vascular occlusion on recovery (2h and 24h) of physiological and neuromuscular indices following sprint exercise while also considering hormonal reactivity to subsequent training. Vascular occlusion did not influence the physiological or neuromuscular markers measured 2h or 24h after sprint exercise in

Academy rugby players. Perception of muscle soreness was not different between conditions, sprinting, increased muscle soreness 24h post-exercise. As similar between-condition responses to a physical stress test occurred at 24h, vascular occlusion did not facilitate recovery following 2h or 24h of rest, nor change the hormonal response to a subsequent physical stress test. Likewise, vascular occlusion did not detrimentally affect any measures assessed, recovery rate was not negatively influenced in comparison to Con, thus alleviating concerns about using this strategy, acutely, within season.

Total ($p=0.238$) and average ($p=0.674$) sprint times were consistent between conditions with similar physiological responses being observed. Johnston et al.²¹ highlighted CK values increased by $570 \mu\cdot\text{L}^{-1}$ (current results $+506.02 \mu\cdot\text{L}^{-1}$). Compared to match responses, increases of $586.6 \mu\cdot\text{L}^{-1}$ (⁴; 24h post-soccer match) and $431 \mu\cdot\text{L}^{-1}$ (²⁶; 16h post-rugby match) have been reported. However, time and distance of maximal sprint completed in a match cannot be controlled and varies depending on position, therefore comparison against match outcome is difficult as many factors may influence performance on match day. Nevertheless, previous reports suggest that six 50m sprints reflect normal training sessions²⁷.

The current study individualised cuff pressure (171-226 mmHg) as a standard pressure applied to different individuals may non-uniformly influence the pressure exerted on the vasculature and thus impact the degree of blood flow restriction. Loenneke et al.²⁸ suggested that pressures aiming to restrict blood flow of the lower body should be determined by limb circumference; findings supported by observations that thigh circumference is the biggest predictor of arterial occlusion in the lower body ($\beta=0.570$)¹⁶ with brachial systolic blood pressure being an additional significant predictor ($\beta=0.231$)¹⁶. That said, in contradiction to the findings of Beaven et al.⁹ who identified improved peak power recovery 24h after occlusion, we observed no effects on recovery. Although comparable vascular occlusion timings were used, cuff pressures differed between the studies (i.e., a standardised 220 mmHg

versus individualised application). Alternative methodological factors may be influencing the resulting outcome of vascular occlusion as a recovery modality; cuff pressure was determined from a regression equation¹⁶ based on non-elite participants, although this protocol considers thigh girth, tissue type (muscle or fat) may impact the level of blood flow restriction, a variable which will vary between elite-trained and untrained participants, training status may also impact effectiveness of vascular occlusion used for recovery.

An alternative methodological discrepancy existing between previous research is the training status of participants. Research completed by Beaven et al.⁹ investigated healthy, active non-sport participating males and found vascular occlusion improved recovery. However, Northey et al.¹⁴ and the current study investigated the use of vascular occlusion in well-trained individuals, identifying no effect on recovery. This potentially suggests that training status may mediate the efficacy of vascular occlusion as a recovery strategy. However, even with a group of active or well-trained participants, high inter-individual variability in EIMD marker responses has been found even when participants perform the same exercise²⁹. Research suggests that the large inter-individual variability including non-modifiable factors (ethnicity, age, and gender) could be responsible for most of the equivocal findings and uncertainties regarding EIMD etiology²⁹; another variable potentially explaining variation between the current study and previous work. Therefore, it may be important to examine the use of vascular occlusion on an individual basis as it may offer a practical recovery option for some athletes/players and at some levels of performance.

A novel aspect of this research was the response of a physical stress test as an indicator of recovery status and readiness-to-train following sprint exercise at 24h. Within the literature, previous recovery studies have assessed response to a recovery strategy when the athlete is rested, however players are frequently required to return to training 24h after a match. Therefore, hormonal response to a physical stress test may indicate if players are recovered and able to return to training. In the current study, T increased pre-to post stress test in both conditions (Recovery: 21.58 ± 44.90 pg·ml⁻¹, $7 \pm 17\%$, $p=0.031$; Con: 10.62 ± 36.57 pg·ml⁻¹,

$4 \pm 13 \%$, $p=0.178$), but ΔT was not significantly different between conditions, suggesting Recovery had no impact on hormonal response. Furthermore, C showed a significant decline from pre-to post-stress test in both the Recovery ($0.14 \pm 0.23 \mu\text{g}\cdot\text{dl}^{-1}$, $38 \pm 72\%$, $p=0.007$) and Con ($0.17 \pm 0.27 \mu\text{g}\cdot\text{dl}^{-1}$, $50 \pm 94\%$, $p=0.006$). Therefore, consistent with rested results, there was no difference between conditions regarding rate of recovery following a physical stress test.

Limitations of the research should be acknowledged; the placebo effect is not accounted for, which may be important to consider. Within Con, cuffs were still applied and therefore is difficult to blind participants from the conditions due to the obvious difference in cuff pressure. Therefore, a further condition would be optimal to identify the impact of vascular occlusion, determining whether there is a placebo or physiological effect on performance when compared against a control in which cuffs are absent as completed by Marocolo et al.³⁰. Similarly, it is difficult to blind testers to conditions due to variation in cuff pressure, however, all testers and coaches were informed that no verbal encouragement was to be given during exercise (sprints, jumps, stress test).

Conclusions

The data presented in this investigation highlight application of two cycles of vascular occlusion administered intermittently after sprinting did not influence, either positively or negatively, physiological or neuromuscular markers of recovery assessed after 2h or 24h of rest or after a subsequent physical stress test.

Practical Implications

- Individualised vascular occlusion applied post-exercise didn't influence recovery or readiness-to-train after 24h

- An absence of negative effects for physiological and performance markers alleviates concerns about the use of vascular occlusion within season
- Methodological variations (such as exercise protocol, cuff pressure, duration of occlusion and training status) may be modulating the efficacy of vascular occlusion as a recovery strategy

Acknowledgements

None to declare. There was no financial support for this study. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

1. Bangsbo, J., Mohr, M. & Krstrup, P. Physical and metabolic demands of training and match-play in the elite football player. *J Sports Sci*, 2006, 24(7), pp.665–674.
2. Ferguson, R.A., Dodd, M.J. & Paley, V.R. Neuromuscular electrical stimulation via the peroneal nerve is superior to graduated compression socks in reducing perceived muscle soreness following intense intermittent endurance exercise. *Eur J Appl Physiol*, 2014, pp.2223–2232.

3. Russell, M. et al. Relationships between match activities and peak power output and Creatine Kinase responses to professional reserve team soccer match-play. *Hum Mov Sci*, 2016, 45, pp.96–101.
4. Shearer, D.A. et al. Measuring Recovery in Elite Rugby Players: The Brief Assessment of Mood, Endocrine Changes, and Power. *Res Q Exerc Sport*, 2015, 86(4), pp.379–86.
5. West, D.J. et al. Neuromuscular function, hormonal, and mood responses to a professional rugby union match. *J Strength Cond Res*, 2014, 28(1), pp.194–200.
6. Fry, R.W. et al., 1991. Monitoring exercise stress by changes in metabolic and hormonal responses over a 24-h period. *Eur J Appl Physiol Occup Physiol*, 63(3–4), pp.228–234.
7. Crewther, B.T. et al. Can salivary testosterone and cortisol reactivity to a mid-week stress test discriminate a match outcome during international rugby union competition? *J Sci Med Sport*, 2017.
8. Nédélec, M. et al. Recovery in soccer: Part II-recovery strategies. *Sports Med*, 2013, 43(1), pp.9–22.
9. Beaven, C.M. et al. Intermittent lower-limb occlusion enhances recovery after strenuous exercise. *Appl Physiol Nutr Metab*, 2012, 37(6), pp.1132–9.
10. Page, W., Swan, R. & Patterson, S.D., 2017. The effect of intermittent lower limb occlusion on recovery following exercise-induced muscle damage: A randomized controlled trial. *J Sci Med Sport*, 2017.
11. Liu, G.S. et al., 1994. Evidence that the adenosine A₃ receptor may mediate the protection afforded by preconditioning in the isolated rabbit heart. *Cardio Res*, 28(7), pp.1057–1061.
12. Howatson, G. & van Someren, K.A., 2008. The prevention and treatment of exercise-induced muscle damage. *Sports Med (Auckland, N.Z.)*, 38(6), pp.483–503.
13. Garcia, C.A. et al. Ischemic Preconditioning and Acute Recovery of Performance in Rugby Union Players. *Sports Med Int Open*, 2017, 1, pp.107–112.
14. Northey, M. et al. Vascular Occlusion and Sequential Compression for Recovery Post Resistance Exercise. *J Strength Cond Res*, 2016, 30(2), pp.533–539.

15. Borne, R., Hausswirth, C. & Bieuzen, F. Relationship Between Blood Flow and Performance Recovery: A Randomized, Placebo-Controlled Study. *I J Sports Physiol Perform*, 2016, (May), pp.1–44.
16. Loenneke, J.P. et al. Blood flow restriction in the upper and lower limbs is predicted by limb circumference and systolic blood pressure. *Eur J Appl Physiol*, 2015, 115(2), pp.397–405.
17. Nédélec, M. et al., 2012. Recovery in soccer: Part 1. *Sports Med*, 43(1), pp.9–22.
18. Crewther, B.T. & Cook, C.J. Effects of different post-match recovery interventions on subsequent athlete hormonal state and game performance. *Physiol Behav*, 106(4), 2012, pp.471–475.
19. West, D.J. et al. Relationships between force-time characteristics of the isometric midthigh pull and dynamic performance in professional rugby league players. *J Strength Cond Res*, 2011, 2(12), pp.3070–3075.
20. Taylor, T. et al. The impact of neuromuscular electrical stimulation on recovery after intensive, muscle damaging, maximal speed training in professional team sports players. *J Sci Med Sport*, 2014, 18(3), pp.328–332.
21. Johnston, M. et al., 2015. Neuromuscular, physiological and endocrine responses to a maximal speed training session in elite games players. *Eur J Sport Sci*, pp.1–7.
22. Ghosh, S., Standen, N. B. Galiñanes, M., 2000. Preconditioning the human myocardium by simulated ischemia: Studies on the early and delayed protection. *Cardio Res*, 45(2), 339-350.
23. Dabbs, J.M. Salivary testosterone measurements: reliability across hours, days, and weeks. *Physiol Behav*, 1990, pp.83-86.
24. Impellizzeri, F.M. & Maffiuletti, N. A. Convergent evidence for construct validity of a 7-point likert scale of lower limb muscle soreness. *Clinical J Sports Med*, 2007, 17(6), pp.494–496.
25. Laurent, C.M. et al. A practical approach to monitoring recovery: development of a perceived recovery status scale. *J Strength Cond Res*, 2011, 25(3), pp.620–628.


26. Jones, M.R. et al., 2014. Match play performance characteristics that predict post-match creatine kinase responses in professional rugby union players. *BMC Sports Sci Med Rehab*, 6(1), p.38.
27. Francis, C. (2008). The structure of training for speed. Retrieved from www.Charliefrancis.com
28. Loenneke, J.P. et al. Effects of cuff width on arterial occlusion: implications for blood flow restricted exercise. *Eur J Appl Physiol*, 2012, 112(8), pp.2903–2912.
29. Damas, F. et al. Susceptibility to Exercise-Induced Muscle Damage : a Cluster Analysis with a Large Sample. *Int J Sports Med*, 2016, 37(8), 633-640.
30. Marocolo, M. et al., 2015. Are the Beneficial Effects of Ischemic Preconditioning on Performance Partly a Placebo Effect? *Int J Sports Med*, 36(10), pp.822–825.

Figure 1: Timeline of data collection. Con; control trial, Recovery; occlusion trial, RE; resistance exercise. Measurements: salivary testosterone, salivary cortisol (●), blood sampling for blood lactate and Creatine Kinase (□), perception muscle soreness questionnaires (◆), countermovement jump (◇)


Figure 2: Physiological responses (a) Testosterone (b) cortisol (c) Creatine Kinase (d) Blood

lactate collected pre, post, 2 h and 24 h post sprint protocol (* $p < 0.05$)


Figure 3: (a) Peak power output and (b) jump height, determined from countermovement jump collected pre, post, 2 h and 24 h post sprint




a




b

2
0
8
6
4
2
0

c




d




a

Peak Power Output (W)



b

Jump Height (cm)



— Occlusion - - - Control