



Swansea University
Prifysgol Abertawe



Cronfa - Swansea University Open Access Repository

This is an author produced version of a paper published in:
Proceedings of Science

Cronfa URL for this paper:
<http://cronfa.swan.ac.uk/Record/cronfa50083>

Conference contribution :

Lee, J., Bennett, E., Hong, D., Lin, C., Lucini, B., Piai, M. & VDACCHINO, D. (2019). *Progress in the lattice simulations of $Sp(2N)$ gauge theories*. Proceedings of Science, (pp. 192 Michigan State University, East Lansing, Michigan, USA: The 36th Annual International Symposium on Lattice Field Theory (LATTICE2018).
<http://dx.doi.org/10.22323/1.334.0192>

Released under the terms of a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC-BY-NC-ND).

This item is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Copies of full text items may be used or reproduced in any format or medium, without prior permission for personal research or study, educational or non-commercial purposes only. The copyright for any work remains with the original author unless otherwise specified. The full-text must not be sold in any format or medium without the formal permission of the copyright holder.

Permission for multiple reproductions should be obtained from the original author.

Authors are personally responsible for adhering to copyright and publisher restrictions when uploading content to the repository.

<http://www.swansea.ac.uk/library/researchsupport/ris-support/>

Progress in the lattice simulations of $Sp(2N)$ gauge theories

Jong-Wan Lee^{*,ab†§}, Ed Bennett^{c‡}, Deog Ki Hong^{a§}, C.-J. David Lin^{de¶}, Biagio Lucini^{f||**},
Maurizio Piai^{g**} and Davide Vadacchino^h

^aDepartment of Physics, Pusan National University, Busan 46241, Korea

^bExtreme Physics Institute, Pusan National University, Busan 46241, Korea

^cSwansea Academy of Advanced Computing, Swansea University, Singleton Park, Swansea, SA2 8PP, UK

^dInstitute of Physics, National Chiao-Tung University, Hsinchu 30010, Taiwan

^eCentre for High Energy Physics, Chung-Yuan Christian University, Chung-Li 32032, Taiwan

^fDepartment of Mathematics, Computational Foundary, Bay Campus, Swansea University, Swansea, SA1 8EN, UK

^gDepartment of Physics, Swansea University, Singleton Park, Swansea, SA2 8PP, UK

^hINFN, Sezione di Pisa, Largo Pontecorvo 3, 56127 Pisa, Italy

E-mail: jwlee823@pusan.ac.kr, e.j.bennett@swansea.ac.uk,
dkhong@pusan.ac.kr, dlin@mail.nctu.edu.tw, b.lucini@swansea.ac.uk,
m.piai@swansea.ac.uk, davide.vadacchino@pi.infn.it

We report on the status of our programme to simulate $Sp(2N)$ gauge theories on the lattice. Motivated by the potential realization of an $SU(4)/Sp(4) \sim SO(6)/SO(5)$ composite Higgs model and the applications to self interacting dark matter, we first perform dynamical simulations of $Sp(4)$ theories with two Dirac flavors in the fundamental representation. Preliminary results of the meson spectrum are presented, along with discussion of the lattice systematics. We also introduce two-index anti-symmetric Dirac fermions. Such fermions are relevant in the context of partial top compositeness, provided they carry $SU(3)$ color quantum numbers, and hence we introduce three (Dirac) copies of them. We present the quenched meson spectrum and explore the phase space of bare lattice parameters. For all the numerical simulations we use the standard Wilson lattice gauge and fermion actions.

The 36th Annual International Symposium on Lattice Field Theory - LATTICE2018
22-28 July, 2018

Michigan State University, East Lansing, Michigan, USA.

*Speaker.

†Supported in part by the National Research Foundation of Korea grant funded by the Korea government(MSIT) (NRF-2018R1C1B3001379).

‡Funded by the Supercomputing Wales project, which is part-funded by the European Regional Development Fund (ERDF) via Welsh Government.

§Supported in part by Korea Research Fellowship program funded by the Ministry of Science, ICT and Future Planning through the National Research Foundation of Korea (2016H1D3A1909283).

¶Supported by Taiwanese MoST grant 105-2628-M-009-003-MY4.

||Supported in part by the Royal Society and the Wolfson Foundation.

**Supported in part by the STFC Consolidated Grants ST/L000369/1 and ST/P00055X/1.

1. Introduction

Many phenomenological models in physics beyond the Standard Model (BSM), either addressing the (little) hierarchy problem in the Standard Model (SM) or searching for theories of dark matter, or both, rely on the existence of novel strong dynamics. Some of them may be realised at short distances by $Sp(2N)$ gauge theories with fermionic matter fields. In particular, the theory with two Dirac fermions in the fundamental representation in which the (enhanced) global symmetry $SU(4)$ is broken to $Sp(4)$ in the presence of the fermion condensate and/or finite fermion mass naturally realizes the $SO(6)/SO(5)$ composite Higgs model, where the Higgs doublets of the conventional SM are identified by the four out of five pseudo Nambu-Goldstone (NG) bosons corresponding to the coset space [1]. If we add fermions in the two-index anti-symmetric representation to this model, we might also explain the mass hierarchy of quarks through the mixing with the fermionic composite operators composed of fermions from both representations, a mechanism referred to as partial compositeness [2]. In the context of dark matter phenomenology, furthermore, $Sp(2N)$ theory with N_f fundamental Dirac flavors ($N_f \geq 2$, but sufficiently small to be in the asymptotically free range) has been considered as one of the candidates for strongly coupled gauge theories realising the strongly-interacting-massive-particle (SIMP) mechanism [3].

$Sp(2N)$ theories with $N \geq 2$ are new territory for the lattice community. While pure gauge theories were investigated in [4] a while ago, the theories with fermions were only started being studied by the current authors very recently. In an earlier publication [5] we described the basic lattice set-up in details, restricted attention to $Sp(4)$ and calculated the spectrum of glueballs and mesons constructed by fundamental fermions in the quenched approximation. In this work we extend our studies of the meson spectrum by replacing the quenched fermions with dynamical ones. In addition, we perform first explorations of the theory with antisymmetric fermionic matter.

2. The model

Within $Sp(2N)$, the global symmetry structure of the phenomenologically interesting composite-Higgs models can be in principle simulated on the lattice without major technical difficulties: it can be realised for the minimal number of hypercolours $N = 2$ by a mixed representation content of two fundamental and three anti-symmetric flavours of Dirac fermions [6]. Since the representations are pseudoreal and real, respectively, the global symmetries of the matter fields are enhanced to $SU(4) \times SU(6)$. The field content of the model in terms of two-component spinor fields is summarized in Table 1. The continuum and lattice theories, as well as the low-energy effective theory (EFT)—at the next-to-the-leading order in the (degenerate) fermion mass, within the context of the hidden-local-symmetry (HLS)—were extensively discussed in [5].

In our numerical simulations we use the standard Wilson lattice action for the gauge links and fundamental Dirac fermions. Gauge configurations are generated by employing the Heat-Bath (HB) algorithm for the pure gauge model and the hybrid Monte Carlo (HMC) algorithm for the model with dynamical fermions. We use a variant of HiRep code [7] appropriately modified to implement the main features of $Sp(4)$ theory [5]. In this work we have the anti-symmetric two-index irreducible representation of $Sp(2N)$: we follow the prescriptions for $SU(N)$ in [7] except subtracting the omega-trace term, where our convention for the $2N \times 2N$ symplectic matrix is

Table 1: The field contents of the theory. The three columns indicate the transformation laws under $Sp(4)$ gauge, global $SU(4)$ and $SU(6)$ symmetries, respectively. q_F and q_{AS} are 2-component fermions.

Fields	$Sp(4)$	$SU(4)$	$SU(6)$
V_μ	10	1	1
q_F	4	4	1
q_{AS}	5	1	6




Figure 1: Masses of pseudoscalar mesons measured on various lattice volumes for $m_0 = -0.77$ (left) and -0.79 (right) with $\beta = 7.2$. The dashed blue lines are the results of exponential fits to the data.

$\Omega = \begin{bmatrix} 0 & \mathbb{I}_{N \times N} \\ -\mathbb{I}_{N \times N} & 0 \end{bmatrix}$. All of the quantities measured on the lattice are converted to physical ones by utilizing the Lüscher's gradient flow technique [8, 9]. Based on the numerical studies of the scale-setting procedure in [5], we decided to use the derivative of the action density $\mathcal{E}(t)$ at nonzero flow time t [10], $\mathcal{W}(t) = t d\mathcal{E}(t)/dt$, as it shows small cut-off dependence than the case of $\mathcal{E}(t)$ itself. The scale is set by $\mathcal{W}(t)|_{t/a^2=(\omega_0/a)^2} = \mathcal{W}_0$ with an optimal choice of $\mathcal{W}_0 = 0.35$.

3. Mesons in $Sp(4)$ with $N_f = 2$ (dynamical) fundamental Dirac fermions

Our observables of interest are the masses and decay constants of flavored pseudoscalar (PS), vector (V) and axial-vector (AV) mesons. Consider the Euclidean two-point correlation functions

$$C_{\mathcal{O}}(\tau) = \frac{1}{L^3} \sum_{\vec{x}} \langle 0 | \mathcal{O}_M(\vec{x}, \tau) \mathcal{O}_M^\dagger(\vec{0}, 0) | 0 \rangle \xrightarrow{\tau \rightarrow \infty} \frac{\langle 0 | \mathcal{O}_M | M \rangle \langle 0 | \mathcal{O}_M | M \rangle^*}{m_M} \left[e^{-m_M \tau} + e^{-m_M(T-\tau)} \right]. \quad (3.1)$$

$\mathcal{O}_M(x) = \bar{u}(x) \Gamma d(x)$ is the mesonic interpolating operator built with fundamental Dirac fermions u and d , where the Dirac structures result in operators which overlap with PS, V and AV meson states. As the matrix elements are proportional to decay constants, we parameterize them by

$$\langle 0 | \bar{u} \gamma_5 \gamma_\mu d | PS \rangle = f_{PS} p_\mu, \quad \langle 0 | \bar{u} \gamma_\mu d | V \rangle = f_V m_V \epsilon_\mu, \quad \langle 0 | \bar{u} \gamma_5 \gamma_\mu d | AV \rangle = f_{AV} m_{AV} \epsilon_\mu, \quad (3.2)$$

where ϵ_μ is the unit polarization transverse to the four-momentum p_μ . Note that our definitions of the meson states $|M\rangle$ involve the self-adjoint isospin fields rather the charged meson fields, where the analogous pion decay constant in QCD is $f_\pi \simeq 93$ MeV. As the measured decay constants




Figure 2: Lattice spacings in units of the inverse of the gradient flow scale w_0 . Red circles, purple squares and blue diamonds denote currently available ensembles for $\beta = 6.9$ with $m_0 = [-0.85, -0.924]$, $\beta = 7.2$ with $m_0 = [-0.6, -0.79]$ and $\beta = 7.5$ with $m_0 = [-0.65, -0.69]$, respectively.

receive multiplicative renormalization, we use the lattice perturbation theory at the one-loop to match the lattice data to the continuum. For further details, we refer the reader to Ref. [5].

Before we present the numerical results of the mass spectrum, we discuss the systematic errors due to the lattice artefacts. First of all, we investigate the finite volume effects. In Fig. 1, we show the masses of the PS meson in lattice units for $\beta = 7.2$ with $m_0 = -0.77$ and -0.79 on three different lattice volumes, $L/a = 16, 20, 24$. As the Sp(4) theory with $N_f = 2$ fundamental Dirac fermions is expected to exhibit color confinement, meson masses will receive finite volume corrections in the form of an exponential fall-off whose characteristic decay rate will be the mass of the lightest excitation in the theory, i.e. the mass of PS meson. We therefore perform a fit of the data to the function, $m_{PS}(L) = m_{PS}^\infty + Ae^{-m_{PS}^\infty L}$, where the fit results are denoted by blue dashed lines. With a statistical uncertainty of $\sim 0.3\%$ the size of FV effects become compatible with the statistical errors at $m_{PS}L \sim 7.5$, and thus we choose the lattice volume for all ensembles to satisfy the condition $m_{PS}L \gtrsim 7.5$.

As we discussed above we achieve the scale setting by using the Lüscher's gradient flow techniques. In contrast to the lattice QCD, we have already showed that the gradient flow scale w_0/a rapidly changes as we vary the fermion mass [5], see also [11, 12]. In Fig. 2 we present the lattice spacings with respect to the PS mass-squared in units of w_0 for currently available ensembles. The dependence on the fermion mass becomes milder as we approach the continuum. On the other hand, the fermion-mass dependence seems to persist in the chiral limit. In this article we do not attempt to take the continuum limit due to the limited number of ensembles, but postpone it to our future work in which we will fully take advantage of the low-energy EFT developed in [5].

We now present the preliminary results of the dynamical spectrum of the mesons for $\beta = 6.9$ and 7.2 with various values of fermion mass in Fig. 3. The masses (left) and the decay constants (right) are shown with respect to the mass squared of the PS meson, which is physical and thus scheme-independent. From the comparison of two values of the lattice coupling, we find that the V meson masses systematically receive sizable corrections due to the discretization. In the case of V and AV mesons, we also find that with sufficiently small m_{PS} the decay constants are consistent with each other for given statistical errors, and insensitive to m_{PS} and β . From the fact that the V




Figure 3: Masses and decay constants of PS (circle), V (square) and AV (diamond) mesons in the Sp(4) theory with two dynamical Dirac fermions in the fundamental representation. We generated ensembles for $\beta = 6.9$ on a 32×16^3 lattice, except for the two lightest-mass ensembles on a 32×24^3 lattice, and for $\beta = 7.2$ on a 36×16^3 lattice, except for the three lightest-mass ensembles on a 36×24^3 lattice.




Figure 4: The Sum of the squared decay constants of the PS, V, AV mesons with dynamical $N_f = 2$ fundamental Dirac fermions. For reference, we also present the fit results by the colored bands for quenched Sp(4) simulations with $\beta = 7.62$ (green) and 8.0 (purple).

meson mass increases for smaller lattice spacing, we expect that the S parameter, defined in the framework of the low-energy EFT only with the lightest mesons, decreases in the continuum limit.

The low-energy NLO EFT predicts that the sum of the squared decay constants of PS, V and AV mesons in the zero momentum limit, $f_0^2 = f_{PS}^2 + f_V^2 + f_{AV}^2$, is independent from the fermion mass, which was evidenced by the quenched spectrum [5]. In Fig. 4 we show the results of f_0^2 measured from dynamical simulations for various masses, where no significant mass dependence is found over the wide ranges. The resulting values are also not statistically far from the results in the quenched cases. Such numerical results strongly support predictions of the EFT indicating the insensitivity of f_0^2 on the fermion mass.

4. Mesons in quenched Sp(4) with anti-symmetric Dirac fermions

In Sp(2N) theory the anti-symmetric two-index representation is real, hence with N_f Dirac flavours the global symmetry is enhanced to $SU(2N_f)$ and broken to $SO(2N_f)$ in the presence of




Figure 5: Masses and decay constants of pseudoscalar (circle), vector (square) and axial-vector (diamond) mesons constructed from fermions in the anti-symmetric (red) and fundamental (blue) representation in the quenched $Sp(4)$ theory at $\beta = 8.0$. The lattice volume was 48×24^3 .




Figure 6: Mass scan of the $Sp(4)$ theory with $N_f = 3$ anti-symmetric Wilson fermions at $\beta = 6.4, 6.5$ and 6.6 from left to right, respectively. The red and blue symbols denote the expectation values of the plaquette $\langle P \rangle$ obtained from random (hot) and unit (cold) initial configurations on a 8^4 lattice.

non-zero (symmetric) condensate. Pseudo NG bosons correspond to $2N_f^2 + N_f - 1$ broken generators which belongs to the coset $SU(2N_f)/SO(2N_f)$. In terms of Dirac flavors, the NG bosons are $N_f^2 - 1$ mesons in the adjoint representation, and $N_f(N_f + 1)/2$ diquarks and anti-diquarks in the symmetric representation. As in the case with fundamental fermions, we focus on the spectrum of flavored PS, V, and AV mesons which are degenerate with the corresponding diquark and anti-diquark states transformed in the same way under the global symmetry in the massless limit. As dynamical ensembles are not available yet, we calculate the masses and decay constants in the quenched limit from the same ensembles used for the fundamental fermions in [5]. The results for $\beta = 8.0$ are shown as red symbols in Fig. 5. For a comparison we also present the results for the quenched spectrum with fundamental fermions denoted by blue symbols. The masses and decay constants for both representations show similar dependence on the PS meson mass, but the overall scale is substantially different.

Toward the dynamical simulation with anti-symmetric fermions, the primary task is to search for any singularity associated with the bulk-phase transition by exploring the bare lattice parameter

space. Using 8^4 lattices with $\beta = 6.4, 6.5, 6.6$, we calculate the expectation values of the plaquette $\langle P \rangle$ by varying the bare fermion masses. We focus on the vicinity of the region in which the plaquette values change abruptly. The results are presented in Fig. 6, where red squares and blue circles are obtained from the random (hot) and unit (cold) initial configurations. We find that hot and cold results are consistent with each other at $\beta = 6.6$, while they are well separated over the range of $m_0 = [-1.145, -1.135]$ at $\beta = 6.4$. Such strong hysteresis at small β provides strong evidence for the existence of a first-order bulk phase transition. We therefore estimate our conservative value of the phase boundary as $\beta \gtrsim 6.6$ at which the continuum extrapolation can be taken correctly.

5. Conclusion

We presented preliminary results of a first lattice calculation of the meson spectrum for the $Sp(4)$ gauge theory with two dynamical fundamental Dirac fermions. We first investigated the systematic effects associated the finite volume and the finite lattice spacing. Although the extrapolation to the continuum limit has not been carried out yet, our numerical results show consistency with the low-energy EFT expectations. We also presented the spectrum for the theory with Dirac fermions in the anti-symmetric representation in the quenched limit. Toward the dynamical simulation, we explored the phase structure in the lattice parameter space and identified the phase boundary for the first order bulk phase transition.

Acknowledgements

We are very grateful to Michele Mesiti and Jarno Rantaharju who assisted us in modifying and improving the HiRep code for this project.

References

- [1] D.B. Kaplan, H. Georgi and S. Dimopoulos, *Phys. Lett. B* **136** (1984) 187.
- [2] D.B. Kaplan, *Nucl. Phys. B* **365** (1991) 259-278.
- [3] Y. Hochberg, E. Kuflik, H. Murayama, T. Volansky and J.G. Wacker, *Phys. Rev. Lett.* **115** (2015) no.2, 021301, [arXiv:1411.3727].
- [4] K. Holland, M. Pepe and U.J. Wiese, *Nucl. Phys. B* **694** (2004) 35, [hep-lat/0312022].
- [5] E. Bennett, D. K. Hong, J.-W. Lee, C.-J. D. Lin, B. Lucini, M. Piai and D. Vadacchino, *JHEP* **03** (2018) 185, [arXiv:1712.04220].
- [6] J. Barnard, T. Gherghetta and T.S. Ray, *JHEP* **02** (2014) 002, [arXiv:1311.6562].
- [7] L. Del Debbio, A. Patella and C. Pica, *Phys. Rev. D* **81** (2010) 094503, [arXiv:0805.2058].
- [8] M. Lüscher, *JHEP* **08** (2010) 071, [arXiv:1006.4518].
- [9] M. Lüscher and P. Weisz, *JHEP* **02** (2011) 051, [arXiv:1101.0963].
- [10] S. Borsányi et al., *JHEP* **09** (2012) 010, [arXiv:1203.4469].
- [11] R. Arthur, V. Drach, M. Hansen, A. Hietanen, C. Pica and F. Sannino, *Phys. Rev. D* **94** (2016) 094507, [arXiv:1602.06559].
- [12] V. Ayyar et al., *Phys. Rev. D* **97** (2018) 074505, [arXiv:1710.00806].