

Article

Household Composition and Inequalities in COVID-19 Vaccination in Wales, UK

Alex Lench ^{1,2,*}, Malorie Perry ^{1,2,†}, Rhodri D. Johnson ², Richard Fry ², Gill Richardson ³, Ronan A. Lyons ², Ashley Akbari ², Adrian Edwards ⁴, Brendan Collins ⁵, Natalie Joseph-Williams ⁴, Alison Cooper ⁴ and Simon Cottrell ¹

¹ Vaccine Preventable Disease Programme and Communicable Disease Surveillance Centre, Public Health Wales, 2 Capital Quarter, Tyndall Street, Cardiff CF10 4BZ, UK

² Population Data Science, Health Data Research UK, Swansea University Medical School, Swansea SA2 8PP, UK

³ Policy, Research and International Development, Public Health Wales, 2 Capital Quarter, Tyndall Street, Cardiff CF10 4BZ, UK

⁴ Wales COVID-19 Evidence Centre, PRIME Centre Wales, Division of Population Medicine, School of Medicine, Cardiff University, 8th floor, Neuadd Meirionnydd, Heath Park, Cardiff CF14 4XN, UK

⁵ Health and Social Services Group, Finance Directorate, Welsh Government, Cardiff CF10 3NQ, UK

* Correspondence: alex.lench@wales.nhs.uk

† These authors contributed equally to this work.

Abstract: The uptake of COVID-19 vaccination in Wales is high at a population level but many inequalities exist. Household composition may be an important factor in COVID-19 vaccination uptake due to the practical, social, and psychological implications associated with different living arrangements. In this study, the role of household composition in the uptake of COVID-19 vaccination in Wales was examined with the aim of identifying areas for intervention to address inequalities. Records within the Wales Immunisation System (WIS) COVID-19 vaccination register were linked to the Welsh Demographic Service Dataset (WSDS; a population register for Wales) held within the Secure Anonymised Information Linkage (SAIL) databank. Eight household types were defined based on household size, the presence or absence of children, and the presence of single or multiple generations. Uptake of the second dose of any COVID-19 vaccine was analysed using logistic regression. Gender, age group, health board, rural/urban residential classification, ethnic group, and deprivation quintile were included as covariates for multivariable regression. Compared to two-adult households, all other household types were associated with lower uptake. The most significantly reduced uptake was observed for large, multigenerational, adult group households (aOR 0.45, 95%CI 0.43–0.46). Comparing multivariable regression with and without incorporation of household composition as a variable produced significant differences in odds of vaccination for health board, age group, and ethnic group categories. These results indicate that household composition is an important factor for the uptake of COVID-19 vaccination and consideration of differences in household composition is necessary to mitigate vaccination inequalities.

Keywords: COVID-19; vaccines; vaccination; immunisation; households; household composition; inequities; inequalities



Citation: Lench, A.; Perry, M.; Johnson, R.D.; Fry, R.; Richardson, G.; Lyons, R.A.; Akbari, A.; Edwards, A.; Collins, B.; Joseph-Williams, N.; et al. Household Composition and Inequalities in COVID-19 Vaccination in Wales, UK. *Vaccines* **2023**, *11*, 604. <https://doi.org/10.3390/vaccines11030604>

Academic Editors: Alika K. Maunakea, Ruben Juarez, May Okihiro and Juan De la Torre

Received: 30 January 2023

Revised: 22 February 2023

Accepted: 5 March 2023

Published: 7 March 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The COVID-19 pandemic was characterised by diverse inequalities and following the development of vaccines, this included widespread differences in the uptake of immunisation. Previous work by this group in 2021 characterised a number of predictors for COVID-19 vaccine uptake [1]. This study, and others, identified multiple inequalities in COVID-19 vaccination with differences by ethnic group being amongst the most severe [1,2]. Whilst previous research has demonstrated inequalities in COVID-19 vaccine uptake, they

have also indicated that further work was necessary to characterise patterns of coverage and provide insights for potential interventions to address the inequalities identified.

The role of housing as an important factor in health and wellbeing is well established and the characterisation of households in terms of the number and relative characteristics of the residents (household composition) is a key consideration in this area [3]. In particular, isolation is well understood as a risk factor for a broad range of adverse health and social issues [4], though other considerations relating to household composition, such as overcrowding and multi-generational living are also important and can be associated with factors such as ethnic group [5,6].

Methodological advances developed by Rodgers et al. [7] have allowed anonymous individual-level data to be linked using unique residence identifiers within the Secure Anonymised Information Linkage (SAIL) databank in Wales [8]. While this technique cannot currently provide information on the type of relationships between different residents of a property, it can be used to effectively characterise households by household size, age of residents, and generational composition [9]. In the present study, we aimed to use this methodology to examine the influence of household composition on COVID-19 vaccination coverage and inequalities.

2. Materials and Methods

Analyses were completed within the SAIL Databank, hosted by Swansea University, as part of the Con-COV (Controlling COVID-19) project [10,11], using the software package R (version 4.1.3). Con-COV is a total population electronic cohort derived from linking many databases [10]. All individuals alive and resident in Wales on 1 January 2022 were included in the initial cohort, identified from the Welsh Demographic Service dataset (WDSD). In Wales, vaccination for COVID-19 began on 8 December 2020, and delivery of first and second doses in the adult population plateaued by one year.

Vaccination status, sex, urban/rural location of residence, health board of residence, ethnic group, and deprivation quintile of Lower-Layer Super Output Area were assigned as previously described [1]. Ethnic group categories were defined by the Office for National Statistics 2011 Census [12]. Individuals with learning disabilities were identified using standard guidance on clinical coding for identifying vaccine eligibility, monitoring/reporting of vaccine uptake, and call/recall purposes, provided by the University of Nottingham PRIMIS group [13].

Household size and generational composition were calculated using an encrypted Unique Property Reference Number, known as a Residential Anonymised Linkage Field (RALF), as previously described [7,9]. Generational composition was calculated using the 'Relative Age to Youngest' (AtY) method [9]. Where the age between the oldest and youngest people in the household is 18 years or less, the household was classified as having one generation and where the age between the oldest and youngest people in the household is over 18 years, the household was classified as having multiple generations.

Mutually exclusive household composition groups (referred to as household types) were calculated based on household size, generational composition, and the presence/absence of at least one child (person < 18 years old) in the household as described in Table 1. Household types were sole occupancy (1), two adults (2), adults/children (3), large adults/children (4), adult group single/multiple generations (5/6), and large adult group single/multiple generations (7/8).

Adults living in households of size 10 or above were excluded from the analysis as these households are likely to contain a substantial proportion of communal residences, such as care homes or supported living where immunisation is coordinated and monitored directly via the residence. Persons with immunocompromised status were also excluded due to this (relatively small) group having a three-dose primary course.

Table 1. Description of household categories used to assess COVID-19 vaccine uptake.

Household Type	Definition	Relevant Housing Classifications
1. Single occupancy	Size 1	Lone dwelling.
2. Two adults	Size 2, no <18	Partnership. Cohabitation.
3. Adults/children	Size 2–5, <18 present	Family ^a
4. Large adults/children	Size 6–9, <18 present	Large family ^a
5. Adult group, single generation	Size 3–5, no <18, age difference <19 years	Communal residence.
6. Adult group, multiple generations	Size 3–5, no <18, age difference >18 years	Family with adult children. Communal residence.
7. Large adult group, single generation	Size 6–9, no <18, 1 generation, age difference <19 years	Large communal residence.
8. Large adult group, multiple generations	Size 6–9, no <18, >1 generation, age difference >18 years	Large family with adult children. Large communal residence.

^a Adults/children and large adults/children households may also include a low proportion of communal residences.

Vaccination of adults (aged 18 years or over as of 31 March 2022) with a complete primary course (i.e., first two doses) of any COVID-19 vaccine, received as of 1 January 2022, was analysed. Uptake figures were calculated overall and stratified by sex, age group, rural/urban classification, learning disability, and ethnic group. The odds of vaccination with a complete primary course were estimated using univariable and multivariable logistic regression. Forward step-wise selection, based on optimising the Akaike information criterion value and consideration of likely confounders, was used to select the adjustment variables for the multivariable model with sex, age group, rural/urban classification, health board of residence, ethnic group, and deprivation quintile included in the analysis. The potential influence of correlations within households was assessed by duplication of analyses using cluster-robust regression with RALF assigned as a clustered factor but no significant differences were detected [14].

3. Results

The study population included 2,317,275 individuals aged 18 years or older and alive and resident in Wales on the 1st of January 2022. Differences in uptake for two doses of the COVID-19 vaccine were observed across household types when calculated overall and for multiple subpopulations (Figure 1). Denominator figures are shown in Supplementary Table S2. The highest percentage uptake was observed for people aged 70 and over living in two-adult households (97%, $n = 213,092$), and the lowest percentage uptake was for people from an Asian ethnic group living in large adult group, single-generation households (29%, $n = 482$).


The highest number of unvaccinated people overall were in adult/children households (97,176) and multi-generation adult groups (79,872). For people with learning disabilities, the largest number of unvaccinated people lived in multi-generation adult groups (400) and two-adult (311) households. For those aged 70 years and older, the highest numbers of unvaccinated people were in single occupancy (4763) and two-adult (5712) household types. Asian, Black, and Other ethnic group categories had the highest numbers of unvaccinated people in adults/children (1936, 1030, and 905, respectively) and large adults/children (2352, 934, and 925, respectively) household types.

In a univariable analysis, the odds of having received a complete primary course were significantly less for individuals living in all other household types compared with the two-adult household type (Table 2). The odds of a sole occupant being vaccinated were lower than for individuals living in a two-adult household type but higher than for individuals in all other household types. In general, lower odds ratios for vaccination were generally associated with large household types and/or household types with smaller total populations.

Table 2. Uptake of two doses of COVID-19 vaccine (any type) by household type and odds of being vaccinated, Wales ^a.

Household Type	Population (n)	Vaccinated (n)	Uptake (%)	OR (95% CI)
1. Single occupancy	327,629	286,963	87.6%	0.70 (0.69–0.71)
2. Two adults	639,882	582,346	91.0%	Baseline
3. Adults/children	549,758	452,582	82.3%	0.46 (0.46–0.47)
4. Large adults/children	135,043	94,799	70.2%	0.23 (0.23–0.24)
5. Adult group, single generation	48,892	33,993	69.5%	0.23 (0.22–0.23)
6. Adult group, multiple generations	577,057	497,185	86.2%	0.62 (0.61–0.62)
7. Large adult group, single generation	6553	3806	58.1%	0.14 (0.13–0.14)
8. Large adult group, multiple generations	32,461	22,472	69.2%	0.22 (0.22–0.23)

^a Data from the Wales Immunisation System (WIS) as of 1 January 2022.



(a)

(b)

Figure 1. Vaccination with a complete two-dose course of COVID-19 vaccine (any type) by household type, stratified for sex, learning disability, age group, and Asian, Black, Mixed, and Other ethnic groups, Wales 2020–2021. (a) Heat map for percentage uptake with light-dark shading for higher to lower percentages across stratified categories. (b) Heat map for number of non-vaccinated people with light-dark shading for lower to higher numbers within stratified categories. Data sourced from the all Wales Immunisation System (WIS) in SAIL, COVID-19 Vaccination Data (CVVD), as of 1st January 2022. Ranges are given where necessary to mask low numbers.

These findings were consistent with results from multivariable regression, which included gender, age group, health board, rural/urban location, ethnic group, and deprivation quintile as covariates (Table 3). In this multivariable analysis, the size of the effects of household type on vaccination were generally around half those in the univariable regression, with three exceptions. Firstly, the single-person household type (1) had a vaccination odds ratio of 0.70 (0.69–0.71) in both univariable and multivariable analyses. Secondly,

the difference for the multi-generational adult group household type (6) was less than a factor of two, with a univariable OR of 0.62 (0.61–0.62) and a multivariable OR of 0.89 (0.88–0.90). Thirdly, the difference for the single-generation, large, adult group household type (7) was greater than a factor of two, with a univariable OR of 0.14 (0.13–0.14) and a multivariable OR of 0.52 (0.49–0.55). In addition to the effect of household type, the odds of being vaccinated were lower in younger age groups, minority ethnic background groups, if residing in a more deprived or urban area, if male, or if resident in certain health boards in the multivariable analysis (Table 3).

Table 3. Multivariable regression model estimates for vaccinations with two doses of COVID-19 vaccine (any type) Wales ^{a,b}.

Variable	Category	Population (n)	Vaccinated (n)	aOR (95% CI)
Household Type	1. Single occupancy	327,629	286,963	0.70 (0.69–0.71)
	2. Two adults	639,882	582,346	Baseline
	3. Adults/children	549,758	452,582	0.83 (0.82–0.84)
	4. Large adults/children	135,043	94,799	0.48 (0.47–0.49)
	5. Adult group, single generation	48,892	33,993	0.53 (0.52–0.54)
	6. Adult group, multiple generations	577,057	497,185	0.89 (0.88–0.90)
	7. Large adult group, single generation	6553	3806	0.52 (0.49–0.55)
	8. Large adult group, multiple generations	32,461	22,472	0.45 (0.43–0.46)
Gender	Female	1,161,191	1,016,737	Baseline
	Male	1,156,084	957,409	0.73 (0.72–0.74)
Age Group	18–29	402,223	298,465	0.30 (0.30–0.30)
	30–49	731,813	575,472	0.39 (0.39–0.40)
	50–69	760,289	694,063	Baseline
	70+	422,950	406,146	2.05 (2.01–2.08)
Health Board	Health Board 1	438,803	380,953	Baseline
	Health Board 2	502,623	427,760	0.72 (0.71–0.73)
	Health Board 3	369,837	303,933	0.86 (0.85–0.88)
	Health Board 4	336,450	294,497	1.04 (1.03–1.06)
	Health Board 5	281,646	241,400	0.73 (0.72–0.74)
	Health Board 6	94,642	81,317	0.67 (0.65–0.68)
	Health Board 7	293,274	244,286	0.74 (0.73–0.75)
Location Classification	Rural	712,806	626,082	1.15 (1.14–1.16)
	Urban	1,604,469	1,348,064	Baseline
Ethnic Group	Asian	44,767	34,775	0.75 (0.73–0.77)
	Black	11,773	7475	0.39 (0.37–0.40)
	Mixed	31,523	21,598	0.46 (0.45–0.47)
	Other	10,432	6740	0.40 (0.39–0.42)
	Unknown	235,552	158,880	0.33 (0.32–0.33)
	White	1,983,228	1,744,678	Baseline
Deprivation Quintile	1 (Most deprived)	452,962	356,673	0.47 (0.46–0.47)
	2	461,071	386,911	0.60 (0.59–0.61)
	3	468,108	397,891	0.68 (0.67–0.69)
	4	465,345	409,365	0.80 (0.79–0.81)
	5 (Least deprived)	469,789	423,306	Baseline

^a Data from the Wales Immunisation System as of 1 January 2022. ^b Deprivation quintile was calculated using Lower-layer Super Output Areas of residence, ranked according to the Welsh Index of Multiple Deprivation.

A multivariable regression was also conducted with the exclusion of household type as a variable to examine the impact of accounting for household composition on the odds of vaccination associated with other variables (Supplementary Table S1). Significant differences between adjusted odds ratios produced in multivariable analysis with and without the inclusion of the household type variable were seen for the 18–29 and 30–49 age groups, one health board of residence, and ethnic group variables. Compared to the White ethnic

group, the adjusted odds ratio for the Asian ethnic group was 0.66 (0.64–0.67) without the inclusion of household type and 0.75 (0.73–0.77) with the inclusion of household type. In addition, the adjusted odds ratio for the Black ethnic group was 0.35 (0.33–0.36) without the inclusion of household type and 0.39 (0.37–0.40) with the inclusion of household type. Adjusted odds ratios for Other and Unknown ethnic group categories were also significantly lower in the multivariable regression which did not include household type.

4. Discussion

In the present study it was observed that, compared to two-adult households, all other household types were associated with a lower probability of vaccination. With the exception of the single occupancy household type (which was consistent between analyses) odds ratios from multivariable regression were less significant than those from univariable regression, indicating that in most cases, the relationship between household type and vaccination is partly mediated by other factors such as age, gender, and deprivation. In general, the lowest probabilities of vaccination were observed for the large (6–9 people) and/or uncommon household types.

Previous studies examining vaccine uptake and household composition have generally focused on household size, particularly with consideration to solitary housing. A meta-analysis by Jain et al. in 2017 [15] included an assessment of living alone as a variable for the uptake of multiple vaccines by older people in European countries and found that whilst individual studies had shown mixed results, overall living alone was associated with decreased vaccine uptake. This is consistent with the results observed here. Further studies examining influenza and zoster vaccination have also found a negative association between living alone and vaccine uptake [16,17].

Research examining inequalities of vaccine uptake based on ethnic groups has identified behavioural drivers, such as distrust of information sources, but has also revealed structural barriers [18,19]. In Wales, people of Black, Asian, or Minority ethnic background are more likely to live in an overcrowded households and this has been linked to worse outcomes for COVID-19 in these groups [20,21]. In the present study, household composition was seen to impact differences in vaccine uptake based on ethnic group and Asian, Black, and Other ethnic groups were associated with a relatively high number of unvaccinated people in large adults/children households. Although our analysis accounted for the impact of deprivation on uptake on the basis of a residential area, households are also likely to vary in level of deprivation within a residential area. Therefore, the relationships between vaccine uptake, ethnic group, large household size, and overcrowding could reflect an underlying pattern of deprivation between households.

The approach of using linkage with a residential identifier for measuring household composition has some limitations compared to other methods such as surveys or census data, particularly in relation to the level of detail. However, as this approach enables the ascertainment of full residential information at an adjustable point in time it is useful for examining variation in vaccination uptake, where phases of an immunisation programme determine the relevant time period. As household composition and vaccine uptake has not been extensively studied previously, this study may provide useful information for other countries wanting to develop vaccine equality studies.

Consideration of people living alone is of particular importance as solitary living confers a vulnerability due to social isolation [4]. In our study, living alone was associated with a decreased likelihood of vaccination compared to two-adult households. A recommendation to improve uptake for those living alone could be for vaccination programmes and primary care systems to routinely assess uptake for people who live in sole occupancy households to ensure the protection of this group. However, the probability of vaccination if living alone was not as low as in large household types. Our study also indicates that housing is a factor for differences in vaccination related to ethnic background and implicates family households in these inequalities. Although such findings could reflect an underlying pattern of deprivation, policies to improve uptake based on living in large

and/or family households may be effective in reducing inequalities for these groups. These include enabling joint appointing for cohabitantes where appropriate and allowing clinic opening times to accommodate those with care duties.

The findings from this study should be taken into consideration when planning the rollout of further COVID-19 booster vaccinations to avoid further widening of inequalities. Moreover, the findings from this study may be relevant to other vaccination programmes in Wales as well as across the UK and other developed countries. Further mixed-methods research should explore the causal mechanisms of differences in vaccine uptake between different types of households and whether these mechanisms are structural in nature or relate predominantly to behaviour and interpersonal relationships.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/vaccines11030604/s1>, Table S1: Multivariate regression model estimates for vaccination with two doses of COVID-19 vaccine (any type), not including household type as a variable, Wales; Table S2: Study population by household type, number of people stratified for sex, age group and Asian, Black, Mixed and Other ethnic group categories, Wales 2020–2021; Table S3: Study population by household type, percentage of people stratified for sex, age group and Asian, Black, Mixed and Other ethnic group categories, Wales 2020–2021.

Author Contributions: A.L. and M.P. led the conception and design of this work with input from S.C., A.L. and M.P. performed the analysis with guidance from R.D.J. and R.F., R.A.L., A.A., G.R. and B.C. provided interpretation of results and wider context on interpretation. A.E., N.J.-W. and A.C. are members of the Wales COVID-19 Evidence Centre and supported the prioritisation of this project and provided comments on preliminary results. A.L. and M.P. drafted the first iteration of the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: The data used for this research were made available through the Con-COV collaboration. Controlling COVID-19 through enhanced population surveillance and intervention (Con-COV): a platform approach, funded by the UKRI Medical Research Council, grant reference: MR/V028367/1. This work was supported by Health Data Research (HDR) UK [HDR-9006] which receives its funding from the UK Medical Research Council, Engineering and Physical Sciences Research Council, Economic and Social Research Council, Department of Health and Social Care (England), Chief Scientist Office of the Scottish Government Health and Social Care Directorates, Health and Social Care Research and Development Division (Welsh Government), Public Health Agency (Northern Ireland), British Heart Foundation (BHF) and the Wellcome Trust; Administrative Data Research (ADR) UK which is funded by the Economic and Social Research Council [grant ES/S007393/1]. This specific project around vaccine equity was funded by the Wales COVID-19 Evidence Centre (WCEC).

Institutional Review Board Statement: All proposals to use SAIL data are subject to review by an independent Information Governance Review Panel (IGRP). Before any data can be accessed, approval must be given by the IGRP. The IGRP gives careful consideration to each project to ensure proper and appropriate use of SAIL data. When access has been approved, it is gained through a privacy protecting safe haven and remote access system referred to as the SAIL Gateway.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data used in this study is available from the SAIL Databank at Swansea University, Swansea, UK, which is part of the national e-health records research infrastructure for Wales. SAIL has established an application process to be followed by anyone who would like to access data via SAIL at: www.saildatabank.com/application-process.

Acknowledgments: Alex Lench led this work as part of a wider project looking at vaccine equity of COVID-19 vaccination led by Malorie Perry. The wider project oversight group includes Simon Cottrell, Gill Richardson, Ronan Lyons as well as Rowena Bailey and Sudha Arumugam of Swansea University, Alison Cooper, Liz Doe, Helen Freegard, Micaela Gal and Adrian Edwards of the Wales COVID-19 Evidence Centre, Jonathan Jones from Welsh Government and Nigel Pearson as a public representative. Enhanced surveillance of COVID-19 vaccination equity is carried out in collaboration with members of the Wales COVID-19 Vaccination Board Data and Epidemiology Sub-Group, including: Public Health Wales (VPDP and CDSC), DHCW and Swansea University (SAIL). Authors of this report would like to thank the DHCW WIS, Data Warehouse and IS teams, the WIS Data

Quality Group and all those carrying out data entry for vaccinations into WIS and associated QA work. We wish to acknowledge the collaborative partnership that enabled acquisition and access to the de-identified data, which contributed to this analysis. The collaboration was led by the Swansea University Health Data Research UK team under the direction of the Welsh Government Technical Advisory Cell (TAC) and includes the following groups and organisations: the SAIL Databank, Administrative Data Research (ADR) Wales, NHS Wales Informatics Service (NWIS), Public Health Wales, NHS Shared Services Partnership and the Welsh Ambulance Service Trust (WAST). All research conducted has been completed under the permission and approval of the SAIL independent Information Governance Review Panel (IGRP) project number 0911. The team at Swansea University who support and maintain the One Wales collaboration, Con-COV project and associated COVID-19 e-cohorts not listed as authors are, Gareth Davies, Rowena Griffiths, Jane Lyons, Laura North.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Perry, M.; Akbari, A.; Cottrell, S.; Gravenor, M.B.; Roberts, R.; Lyons, R.A.; Bedston, S.; Torabi, F.; Griffiths, L. Inequalities in coverage of COVID-19 vaccination: A population register based cross-sectional study in Wales, UK. *Vaccine* **2021**, *39*, 6256–6261. [CrossRef]
2. Viswanath, K.; Bekalu, M.; Dhawan, D.; Pinnamaneni, R.; Lang, J.; Mcloud, R. Individual and social determinants of COVID-19 vaccine uptake. *BMC Public Health* **2021**, *21*, 818. [CrossRef] [PubMed]
3. Whitehead, M.; Dahlgren, G. What can be done about inequalities in health? *Lancet* **1991**, *338*, 1059–1063. [CrossRef] [PubMed]
4. Holt-Lunstad, J. Social isolation and health. *Health Aff.* **2020**. [CrossRef]
5. Muennig, P.; Jiao, B.; Singer, E. Living with parents or grandparents increases social capital and survival: 2014 General Social Survey-National Death Index. *SSM Popul. Health* **2017**, *4*, 71–75. [CrossRef] [PubMed]
6. Cabinet Office. Ethnicity Facts and Figures: Overcrowded Households. 2020. Available online: <https://www.ethnicity-facts-figures.service.gov.uk/housing/housing-conditions/overcrowded-households/latest> (accessed on 11 January 2023).
7. Rodgers, S.E.; Lyons, R.A.; Dsilva, R.; Jones, K.H.; Brooks, C.J.; Ford, D.V.; John, G.; Verplancke, J.-P. Residential Anonymous Linking Fields (RALFs): A novel information infrastructure to study the interaction between the environment and individuals' health. *J. Public Health* **2009**, *31*, 582–588. [CrossRef] [PubMed]
8. Ford, D.V.; Jones, K.H.; Verplancke, J.P.; Lyons, R.A.; John, G.; Brown, G.; Brooks, C.J.; Thompson, S.; Bodger, O.; Couch, T.; et al. The SAIL Databank: Building a national architecture for e-health research and evaluation. *BMC Health Serv. Res.* **2009**, *9*, 157. [CrossRef] [PubMed]
9. Johnson, R.D.; Griffiths, L.J.; Hollinghurst, J.P.; Akbari, A.; Lee, A.; Thompson, D.A.; Lyons, R.A.; Fry, R. Deriving household composition using population-scale electronic health record data—A reproducible methodology. *PLoS ONE* **2021**, *16*, e0248195. [CrossRef] [PubMed]
10. Jones, K.H.; Ford, D.V.; Thompson, S.; Lyons, R.A. A Profile of the SAIL Databank on the UK Secure Research Platform. *Int. J. Popul. Data Sci.* **2019**, *4*, 1134. [CrossRef] [PubMed]
11. Lyons, J.; Akbari, A.; Torabi, F.; Davies, G.I.; North, L.; Griffiths, R.; Bailey, R.; Hollinghurst, J.; Fry, R.; Turner, S.L.; et al. Understanding and responding to COVID-19 in Wales: Protocol for a privacy-protecting data platform for enhanced epidemiology and evaluation of interventions. *BMJ Open* **2020**, *10*, e043010. [CrossRef] [PubMed]
12. Office for National Statistics. 2011 Census Variable and Classification Information: Part 6. 2013. Available online: <https://www.ons.gov.uk/census/2011census/2011censusdata/2011censususerguide/variablesandclassifications> (accessed on 11 January 2023).
13. Open Code Lists, Wider Learning Disability. Available online: <https://www.opencodelist.org/codelist/primis-covid19-vacc-uptake/learndis/v.1.5.3/> (accessed on 29 January 2023).
14. lm.cluster: Cluster Robust Standard Errors for Linear Models and General Linear Models. Available online: <https://www.rdocumentation.org/packages/miceadds/versions/3.16-18/topics/lm.cluster> (accessed on 29 January 2023).
15. Jain, A.; van Hoek, A.J.; Boccia, D.; Thomas, S.L. Lower vaccine uptake amongst older individuals living alone: A systematic review and meta-analysis of social determinants of vaccine uptake. *Vaccine* **2017**, *35*, 2315–2328. [CrossRef] [PubMed]
16. Endrich, M.M.; Blank, P.R.; Szucs, T.D. Influenza vaccination uptake and socioeconomic determinants in 11 European countries. *Vaccine* **2009**, *27*, 4018–4024. [CrossRef] [PubMed]
17. Jain, A.; Walker, J.L.; Mathur, R.; Forbes, H.J.; Langan, S.M.; Smeeth, L.; van Hoek, A.J.; Thomas, S.L. Zoster vaccination inequalities: A population based cohort study using linked data from the UK Clinical Practice Research Datalink. *PLoS ONE* **2018**, *13*, e0207183. [CrossRef] [PubMed]
18. Epperson, A.E.; Carson, S.L.; Garcia, A.N.; Casillas, A.; Castellon-Lopez, Y.; Brown, A.F.; Nanibaa'A, G. A qualitative study of COVID-19 vaccine decision making among urban Native Americans. *Vaccine X* **2022**, *12*, 100212. [CrossRef] [PubMed]
19. Balasuriya, L.; Santilli, A.; Morone, J.; Ainooson, J.; Roy, B.; Njoku, A.; Mendiola-Iparraguirre, A.; O'Connor Duffany, K.; Macklin, B.; Higginbottom, J.; et al. COVID-19 Vaccine Acceptance and Access Among Black and Latinx Communities. *JAMA Netw. Open* **2021**, *4*, e2128575. [CrossRef] [PubMed]

20. Clifford, S. Coronavirus (COVID-19) and the Black, Asian and Minority Ethnic (BAME) Population in Wales. 2020. Available online: <https://www.gov.wales/sites/default/files/statistics-and-research/2020-06/coronavirus-COVID-19-and-the-black-asian-and-minority-ethnic-population-154.pdf> (accessed on 11 January 2023).
21. Addison, K.; Batt, B.; Stock, K. Housing Overcrowding and Ethnicity: A Literature Review. 2022. Available online: <https://gov.wales/housing-overcrowding-and-ethnicity-literature-review> (accessed on 11 January 2023).

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.