

When should an athlete retire or discontinue participating in contact or collision sports following sport-related concussion? A systematic review

Michael Makdissi^{1,2}

Meghan Critchley³

Robert C. Cantu⁴

Jeffrey G. Caron^{5,6}

Gavin A. Davis⁷

Ruben J. Echemendia⁸

Pierre Frémont⁹

K. Alix Hayden¹⁰

Stanley A. Herring¹¹

Sidney R. Hinds¹²

Barry Jordan¹³

Simon Kemp^{14,15}

Mike McNamee^{16,17}

David Maddocks¹⁸

Shinji Nagahiro¹⁹

Jon Patricios²⁰

Margot Putukian²¹

Stacy Sick³

Michael Turner^{22,23}

Kathryn J. Schneider³

1. Australian Football League, Melbourne Australia
2. Florey Institute of Neuroscience and Mental Health, University of Melbourne, Australia
3. Sport Injury Prevention Research Centre, Faculty of Kinesiology, University of Calgary, Canada
4. Boston University School of Medicine, USA
5. School of Kinesiology and Physical Activity Sciences, University of Montreal, Canada
6. Centre for Interdisciplinary Research in Rehabilitation of Greater Montreal, Canada
7. Murdoch Children's Research Institute, Australia

8. University Orthopedics Center Concussion Care Clinic, State College, PA; University of Missouri- Kansas City, Kansas, USA
9. Faculty of Medicine, Laval University, Québec, Canada
10. Libraries and Cultural Resources, University of Calgary, Canada
11. Department of Rehabilitation Medicine, University of Washington, Seattle, Washington, USA
12. Uniformed Services University, Bethesda, MD, USA
13. Keck School of Medicine, University of Southern California, USA
14. Rugby Football Union, London, UK
15. London School of Tropical Medicine, London, UK
16. Department of Movement Sciences, KU Leuven, Belgium
17. School of Sport and Exercise Sciences, Swansea University, UK
18. Perry Maddocks Trollope Lawyers, Australia
19. Department of Neurosurgery, Yoshinogawa Hospital, Japan
20. Wits Sport and Health (WiSH), School of Clinical Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa
21. Major League Soccer, Princeton, NJ, USA
22. University College London, UK
23. International Concussion and Head Injury Research Foundation ICHIRF

Corresponding author

Dr. Michael Makdissi

Australian Football League, Melbourne Australia

E-mail: michael.makdissi@afl.com.au

Key words: brain concussion, sport, retire, post-concussion syndrome

Word count: 4927

Abstract word count: 254

Author Note

To the editor-in-chief of BJSM, Dr. Jonathan Drezner: this paper is submitted for your consideration as part of the concussion in sport-themed issues for BJSM, May 2023. This manuscript has been internally reviewed by the expert group represented at the Amsterdam Consensus meeting on Concussion in Sport in October 2022.

Acknowledgements

The authors are grateful for the help of Zahra Premji with the search strategy for this systematic review.

Disclosures

All authors have submitted an International Committee of Medical Journal Editors (ICMJE) Disclosure form.

Funding

This systematic review was prepared for presentation at the 6th International Conference on Concussion in Sport in Amsterdam, October 2022. An education grant was received from the Concussion in Sport International Consensus Conference Organizing Committee through Publi Creations for partial administrative and operational costs associated with the writing of the systematic reviews.

Registration: PROSPERO CRD42022155121

ABSTRACT

Objective: To systematically review the scientific literature regarding factors to consider when providing advice or guidance to athletes about retirement from contact or collision sport following sport-related concussion (SRC), and to define contra-indications to children/adolescent athletes entering or continuing with contact or collision sports after SRC.

Data sources: Medline, Embase, SPORTSDiscus, APA PsycINFO, CINAHL, and Cochrane Central Register of Controlled Trials were searched systematically.

Study eligibility criteria: Studies were included if they were i) original research, ii) reported on SRC as the primary source of injury, iii) evaluated the history, clinical assessment and/or investigation of findings that may preclude participation in sport, and iv) evaluated mood disturbance and/or neurocognitive deficits, evidence of structural brain injury, or risk factors for increased risk of subsequent SRC or prolonged recovery.

Results: Of 4355 articles identified, 93 met the inclusion criteria. None of the included articles directly examined retirement and/or discontinuation from contact or collision sport. Included studies examined factors associated with increased risk of recurrent SRC or prolonged recovery following SRC. In general, these were low-quality cohort studies with heterogeneous results and moderate risk of bias. Higher number and/or severity of symptoms at presentation, sleep disturbance, and symptom reproduction with Vestibular Ocular Motor Screen testing were associated with prolonged recovery and history of previous concussion was associated with a risk of further SRC.

Conclusion: No evidence was identified to support the inclusion of any patient-specific, injury-specific or other factors (e.g., imaging findings) as absolute indications for retirement or discontinued participation in contact or collision sport following SRC.

What is already known?

- Advice from clinicians is commonly sought regarding decisions about when an athlete should retire or discontinue participating in contact or collision sport following sport-related concussion (SRC).
- Decisions on when to cease participation in contact or collision sport are complex and multifaceted.
- Historically, several opinion-based articles have been published, however, there are no evidence-based guidelines to facilitate decision-making by athletes with clinical input from healthcare providers.

What are the new findings?

- No evidence was identified to support the inclusion of any patient-specific, injury-specific or other factors (e.g. imaging findings) as absolute indications for retirement or discontinued participation in contact or collision sport.
- Decisions regarding retirement or discontinuation from contact or collision sport in the context of SRC or repeated head trauma should involve clinicians with expertise in traumatic brain injury and sport and be individualised with consideration of psychosocial, injury-specific and sport-specific factors, any persisting symptoms, and evolving neurocognitive concerns.
- Clinicians should provide athletes (and their families or guardians) with the scientific facts and uncertainties of their condition, as well as the potential risks of returning to their previous sport and/or alternative activities with less chance of head trauma or SRC. The discussion should highlight the benefits of regular physical activity and incorporate the athlete's preferences, risk tolerance, and capability to make an informed decision.

INTRODUCTION

Advice from clinicians is increasingly being sought regarding decisions about when an athlete should retire or discontinue participating in contact or collision sports in the context of sport-related concussion (SRC) and/or repeated head trauma. Discussions may be initiated by the clinician involved in the care of the athlete, the athletes themselves, other healthcare team members, coaches or administrators, or family members/friends of the athlete. Increasingly, there is also media scrutiny of athletes who have sustained repeated concussions. The process may be prompted by the presence of factors including a history of multiple concussions, even in the absence of a recent injury, prolonged or persisting symptoms, decreasing scores on neurocognitive or other performance measures, evidence of changes on conventional structural neuroimaging, or concern about changes in mood or behaviour, or long-term risk of mood disorders or neuro-degenerative diseases that may be associated with SRC and/or repeated head trauma.¹⁻³

Decisions on when to cease participation in contact or collision sports are typically complex and multifaceted. For instance, some athletes forego formal education to pursue a career in sport, and many high-performing or elite athletes tend to strongly identify with their athlete role, which are factors that can add to the complexity of discussions surrounding athletic retirement in the context of SRC.^{4,5} Furthermore, mental health symptoms and disorders are often reported in elite athletes^{6,7} and injury, as well as retirement, have been shown to be significant athlete-specific triggers.⁸ It is important to consider incorporating a shared decision-making model that includes providing athletes with the scientific facts and uncertainties of their condition and considers the athlete's preferences and risk tolerance as well as psychological readiness to make a shared decision.⁹⁻¹³

Historically, several opinion-based articles have been published in the literature,¹⁴⁻¹⁷ however, to date, there have been no definitive evidence-based guidelines to facilitate decision-making by athletes with clinical input from healthcare providers. The lack of clear evidence-based guidance may lead to an athlete who has been given the advice to cease participation in contact or collision sports, consulting several different doctors or "doctor-shopping". In addition, the decision may generate concern among clinicians about their

medico-legal exposure when providing advice that may “restrict the trade” of athletes, particularly in the case of professional athletes. It is also widely acknowledged that participation in exercise or physical activity confers a wide range of physical, cognitive and psychological health benefits¹⁸⁻²⁰ as well as financial rewards, particularly at the elite or professional level. In some individuals, these benefits may be mitigated or outweighed by potential risks related to SRC and/or repeated head trauma, including possible long-term effects on neurocognitive or brain health.

The aim of this study was to conduct a systematic review of the literature regarding the factors that should be considered when providing advice or guidance to athletes about retirement from contact or collision sport following SRC, and the contra-indications to children/adolescent athletes entering or continuing with contact or collision sports.

METHODS

The review protocol was prospectively registered in the PROSPERO database for systematic reviews (protocol ID: CRD42022155121) and reported according to PRISMA guidelines.²¹

The authors for this paper were selected based on research and/or clinical expertise in providing retirement recommendations following SRC, with broad representation across specialities, sports, and countries.²² *(cite Schneider et al. methods paper)*

The search strategy was developed in Medline and utilized three main concepts: concussion, sports and retirement. Retirement was defined broadly for the search to capture all possible studies. The standardized search was developed for the concepts for concussion and sports for all the reviews and are described in the concussion consensus methods paper.²² *(cite Schneider et al. methods paper)* To help develop the search for retirement, all authors were invited to contribute keywords and reviewed the draft search. The draft search was created by the health sciences librarian (KAH) and reviewed by all authors as well as peer-reviewed by another health sciences librarian (ZP). Suggestions for additional keywords from co-authors were incorporated as required. The search included both subject headings and

keywords. Keywords were searched in the title, abstract and author-supplied keywords. When translating the search to other databases, keywords were similar, whereas subject headings were responsive to the indexing of the database. All searches were limited to 2001 – March 2022. All search results were exported as RIS files and then uploaded to Covidence for deduplication and screening.

The following databases were searched: MEDLINE® and Epub Ahead of Print, In-Process & Other Non-Indexed Citations and Daily (Ovid), Embase (Ovid), Cochrane Central Register of Controlled Trials (Ovid), CINAHL Plus with Full Text (EBSCO2022), and SPORTDiscus with Full Text (EBSCO). Searches were initially conducted October 2019, and top-up searches were run in March 2022. The complete search strategies for all databases are available in Supplementary File 1. The Medline search is annotated to provide information on search details.²³

Articles that met the following criteria were eligible for inclusion: 1) original research (including randomised controlled trials (RCTs), quasi-experimental designs, cohort, case control, cross sectional, case series, qualitative studies); 2) had SRC as the focus of the study (i.e., >50% of the study population); 3) written in English; 4) peer-reviewed; 5) evaluated mood disturbance and/or neurocognitive deficits; decreased impact threshold for future concussion; evidence of structural brain injury (clinical, radiological, biomarker, etc.); prolonged recovery from concussion [e.g. persisting (>14 days in adults, >30 days in children) or permanent concussive symptoms]; evaluated risk factors for prolonged recovery; evaluated lower threshold for concussion; evaluated history, clinical assessment and/or investigation findings that may preclude participation in sport. We defined children as 5-12 years and adolescents as 13-18 years.

An initial rapid screen was completed to exclude all articles that clearly did not meet inclusion criteria (e.g., not SRC or <50% of the study population SRC, non-human studies, conference proceedings). Each title/abstract was screened by two authors independently. Inter-rater agreement was assessed prior to initiating the title and abstract screen with a threshold of 80% agreement. Discrepancies were resolved by a third, blinded reviewer. Full-

text screening was completed by two authors independently and discrepancies were resolved by a third author.²² *(cite Schneider et al. methods paper)*

Two authors extracted data for each study using predesigned data extraction forms. Similarly, risk of bias (ROB) assessment was performed by two authors using the appropriate Scottish Intercollegiate Guidelines Network (SIGN; <https://www.sign.ac.uk/>) critical appraisal notes and checklist. All articles were rated as low (acceptable or high quality) or high (unacceptable) ROB based on the assessment. Any discrepancies were discussed between reviewing authors until a consensus was reached. Where needed, the lead author acted to resolve conflicts. Strength of Recommendation Taxonomy (SORT) was used to assess the overall quality of the evidence.²⁴

Equity, diversity, and inclusion statement

We included all eligible studies in the systematic review regardless of sample characteristics. Most studies included both males and females (85%) and no studies specifically mentioned the inclusion of parasport athletes. The authors of this review include both women and men, a variety of disciplines, and persons of colour. The team lacks the perspectives of members of the LGBTQI+ community, those in early career stages, and persons from low- and middle-income geographical regions.

RESULTS

We found 8269 records in the database searches (figure 1). After the removal of duplicates, 4355 records were screened by title and abstract during rapid review, and 1807 records were screened by title and abstract to determine eligibility for inclusion. Following full-text review of 204 studies, 93 were included in the review.

(Figure 1 here)

The data extraction table and risk of bias assessments are summarised in the supplementary online material. The studies included 38 prospective cohort studies, 42 retrospective cohort studies, 11 case-control studies and 2 cross-sectional studies. Sixteen studies included

adults only (i.e., aged > 18 years), with the remainder of the studies including children and adolescent subjects (two studies only included children i.e., <12 years, 14 studies only included adolescent subjects i.e., aged 13 to <18 years).

Ten studies included male subjects only, 79 studies included both male and female subjects, and four studies did not report on the sex of their cohort.

None of the studies directly examined the issue of retirement and/or discontinuation from contact collision sports. All included studies assessed factors associated with prolonged recovery following SRC (summarized in table 1) and/or increased risk of concussion (summarized in table 2). Five studies were rated high quality, 75 of acceptable quality and 13 had a high (unacceptable) risk of bias (Supplementary File 2).

Table 1. Factors associated with increased time to recovery/prolonged symptoms following SRC

Factor	Studies demonstrating Increased risk	No association demonstrated
<i>Age</i>	25 26 27	28 29* 30* 31 32 33 34 35 36
<i>Sex</i>	26 32 37 38 39 40 41 42 43 44 45	25 28 29* 34 35 36 46 47* 48 49 50 51 52
<i>Race</i>		28 29* 36 51
<i>Previous concussions</i>	28 36 43 44 51 53 54 55 56 57*^ 58 59	25 26 32 33 34 46 48 49 52 60 61 62 63 64 65 66
<i>Premorbid Headache syndrome/migraines</i>	26 51 59 67	25 31 43 49 64 68
<i>Mental health/mood disorders (anxiety/depression)</i>	36 69 70 71 72	25 31 37 46 48 51 73 74
<i>Learning disability</i>		26 28 29* 33 37 46 48 51 64 70
<i>ADHD</i>	28 43	26 33 36 37 46 51 60 64 65 70

<i>Continued participation in sport</i>	39 46 75	76
<i>Longer time from injury to receiving medical attention</i>	32 37 44 58 68 76 77	
<i>Total number/ severity of symptoms at initial assessment</i>	25 26 34 35 40 42 48 59 76 78 79 80 81	31 52
<i>Post-traumatic headache/migraines</i>	26 29* 37 41 82	48 64
<i>Specific symptom clusters</i>	33 36 41 42 49 52 59 83	
<i>Worsening symptoms from injury to clinic presentation</i>	41	
<i>Fear of pain questionnaire score</i>	25	
<i>Alcohol, tobacco, or marijuana use since injury</i>	51	
<i>Low resilience score (Connor-Davidson Resilience Scale)</i>	84	
<i>Dizziness</i>	64 70	
<i>Difficulty concentrating</i>	29*	
<i>Difficulty remembering</i>		29*
<i>Amnesia</i>	29* 34 66	35 36 37 43 49 64
<i>Loss of consciousness</i>	29* 34 40	35 36 37 43 49 64 66 70
<i>Sleep disturbance</i>	30 85 86	
<i>Orthostatic intolerance</i>	58	
<i>Presence of vestibular/oculomotor deficits</i>	25 58 59 70 81 87 88 89 90 91 92 93 94* 95	
<i>Presence of convergence insufficiency</i>	95 96	70
<i>Tandem gait deficits</i>	58 88 97	
<i>Presence of neuro-psychological deficits on</i>	37 80 81 98*	35

<i>computerised screening tests (e.g., IMPACT)</i>		
<i>Exercise tolerance in sub-acute phase of recovery</i>	99	
<i>Dual-task transverse plane movement and lateral step variability</i>	100	
<i>“5P clinical risk score”</i>	101	
<i>(GT)n Promoter Polymorphism</i>	102	
<i>Plasma t-tau</i>	103	
<i>Salivary microRNA biomarkers</i>	104	
<i>CSF biomarkers</i>	105	
<i>Higher perceived parental stress</i>	74	
<i>Structural neuroimaging (MRI)</i>	106*^ 107*	108
<i>Advanced neuroimaging</i>	109 110* 111 112* 113 * 114* 115	
<i>Functional Near Infra-Red Spectroscopy (Haemodynamic response)</i>	116^	

* = High (unacceptable) risk of bias; ^ = Associated with long-term risk/changes; DTI = diffusion tensor imaging

Table 2. Factors associated with risk of concussion

Factor	Studies demonstrating increased risk	No association demonstrated
<i>Sex</i>	78	45
<i>Previous concussions</i>	45 56 62 66 78 80	
<i>Level of play</i>		45
<i>Mental health disorders (anxiety/depression)</i>	69	

<i>Genetics</i>	117	118*
-----------------	-----	------

* = High (unacceptable) risk of bias

In general, the findings across studies were heterogeneous. The included studies differed widely in the clinical outcomes measured, the timing of assessments, definitions for “prolonged symptoms” and whether conducted in specialty concussion clinics.

The most consistent results for factors associated with increased time to recovery and/or risk of persisting symptoms include longer time to presentation,^{32 37 44 58 68 76 77} total number and/or severity of symptoms at initial presentation,^{25 26 34 35 40 42 48 59 76 78 80 81} sleep disturbance,^{30 85 86} and symptom reproduction with Vestibular Ocular Motor Screen (VOMS) testing.^{25 58 59 70 81 87-95}

Studies on plasma, salivary and/or CSF biomarkers or advanced imaging biomarkers were limited in design, often with small cohorts of athletes.^{103 104 109 111-115 119}

For studies assessing factors associated with an increased risk of concussion, the most consistent finding was that previous concussion was associated with a risk of further concussions.^{45 56 62 66 78 80}

Given the lack of consistent and high-quality research, the variability in outcomes selected and lack of studies evaluating the specific question related to criteria for retirement, the strength of recommendation for criteria for retirement was rated as B (“Recommendation based on inconsistent or limited quality patient-oriented evidence”) or C (“Recommendation based on consensus, usual practice, opinion, disease-oriented evidence, case series for studies of diagnosis, treatment, prevention or screening”).²⁴

DISCUSSION

There was no evidence identified by this systematic review to support the inclusion of any patient-specific, injury-specific, or other factors (e.g., imaging findings) as absolute indications for retirement or discontinued participation in contact or collision sports.

When considering our research questions, we included a broad search of factors that may lead to an increased risk of repeat concussions and/or a significant risk of adverse health outcomes if the individual was exposed to further head trauma and/or SRC. The factors evaluated in the included studies investigated common clinical presentations such as prolonged or persisting post-concussion symptoms, neurological or cognitive deficits, persistent mood or behavioural disturbance, decreased impact threshold for future concussion and radiological evidence of structural brain injury (some of which may be detected incidentally and may not be related to SRC).

The results revealed that several factors may be associated with either an increased risk of recurrent SRC or of prolonged or persisting symptoms following SRC (see tables 1 and 2). Greater time from injury to receiving medical attention has been shown to be associated with longer recovery.^{32 37 44 58 68 76 77} From a secondary prevention standpoint, recognizing the importance of early assessment and management of SRC may result in reduced risk of persisting symptoms. Similarly, athletes with initial high symptom burden⁶⁸ and children/adolescents with higher risk scores¹²⁰ could be screened and directed for evidence-informed interventions in the early period following SRC, which may reduce the risk of persisting symptoms. However, the results were inconsistent across studies, which may be due in part to the heterogenous populations, study designs and outcome measures used. Furthermore, the quality and quantity of the data were low, with a high risk of bias identified in many of the studies (see Supplementary File 2).

Studies examining factors such as age, sex, prior concussions, ADHD or learning disabilities and prior history of mood disorders or migraines demonstrated variable outcomes. This is likely due to variability in study quality, exposure and outcomes evaluated. Many studies sampled patients from specialty clinics which raises the threat of selection bias, where the participants included in the studies may represent a more severe sample and thus more likely to have suffered from persistent symptoms or functional limitations that may

overestimate the relationship between exposure and outcome. Furthermore, many studies conducted univariate analyses and failed to take into consideration important potential confounding factors. Thus, future high-quality incident cohort studies (i.e., following groups of athletes forward over time) are needed; until then, the answers to our questions remain largely unanswered.

Given the evolving nature of the evidence regarding SRC and ongoing concerns related to potential long-term adverse psychological and/or neurocognitive outcomes, it is expected that clinicians will increasingly be faced with questions regarding retirement or discontinuation of contact or collision sports. We suggest a pragmatic approach based on the collective clinical and research experience of the author group and guided by the systematic review. The advice provided is not intended as a clinical practice protocol/directive or legal standard of care and should not be interpreted as such, but rather as a set of general principles to be used by clinicians to provide structure to the complex and multifaceted decision-making process regarding retirement or discontinuation from contact or collision sports in the context of SRC or repeated head trauma.

Traditionally in sports medicine, return to play or return to sports decisions following injury rely on the identification of “absolute” contra-indications that may signify an unacceptable risk to return, the consideration of “relative” contra-indications that may be associated with an increased risk of poor outcomes, and the assessment of other important individual factors such as athlete knowledge, values, priorities, risk tolerance, etc.

There are several factors that have been suggested in the literature as indications for not returning to contact or collision sports following SRC.¹⁴⁻¹⁷ These are summarised in table 3.

Table 3. Factors suggested as possible indications for not returning to contact sport.

Persisting or prolonged
a) symptoms after concussion and/or,
b) neurological abnormalities on physical examination and/or,
c) deficits on neuropsychological testing,

despite time away from contact or collision sports, and compliance with prescribed treatments or interventions
Increasing symptom severity and/or duration with repeat concussions
Concussions precipitated by lower threshold impacts
Structural abnormalities identified on conventional neuroimaging that may be associated with an increased risk of poor outcomes with future head impact and require neurosurgical assessment and management

Based on the results of the systematic review, there is no evidence to support any one factor being an “absolute” contraindication to an athlete being able to continue participation in a contact or collision sport. Given the current limitations in knowledge, it is important that clinicians do not over-index any specific anecdotal guidelines or rely on any one clinical factor when advising athletes about when to retire or discontinue participation in contact or collision sports. Rather, an individualised and collaborative approach to shared decision-making is recommended.

Athletes with persisting symptoms, neurological abnormalities and/or neurocognitive deficits are typically not considered to be cleared for return to contact or collision sports. Given the non-specific nature of the symptoms, and considerable overlap with other syndromes (e.g., mood disorders, post-traumatic headaches/migraines), it is essential to identify and treat co-existent pathology that may respond positively to treatment.^{121 122} (*cite Schneider et al. review 5 and Yeates et al. review 6*) Consequently, it is recommended that clinicians with experience in SRC be involved in the assessment and treatment of the athlete before discussions regarding retirement from sport are initiated. Furthermore, in these instances, a cautious process-oriented approach to decision-making is suggested, where it may be more prudent to restrict contact or collision sport for a prolonged period (e.g., one season) before re-assessment and reconsideration of retirement. Obviously, in some cases, the athlete may choose not to return to contact or collision sports given the impact that symptoms may have on work, education and quality of life.

In this systematic review, we identified several factors that may be associated with an increased risk of recurrent SRC, prolonged or persisting symptoms or long-term adverse

sequelae (tables 1 and 2). These form the basis of important clinical factors that should be considered as part of a comprehensive assessment in the multi-faceted decision-making process regarding retirement or discontinuation from contact or collision sports in the context of SRC or repeated head trauma.

Based on the results of this systematic review and the collective experience of the author group, it is suggested that the following issues are also considered when discussing ongoing participation in contact or collision sport.

1. *History of prior concussion*, including frequency (decreased interval) between concussions, symptom burden and severity at the time of injury, duration of clinical recovery from previous concussions and increasing severity with each successive concussion or minimal impact producing concussion.
2. *Sport-specific factors*, including, type of sport played, years played, position and playing style. Inherent to this discussion is a detailed understanding of concussion risk as well as the head impact dynamics (i.e., the number and magnitude of acceleration, deceleration, and rotational forces) commonly observed in the sport and the implications these may have to return the athlete to the risks associated with that sport. The level of play is also important. Financial benefits exist in professional sports, which may add to the complexities of the decision-making process, as well as the legal implications for the healthcare provider who may be concerned about restriction of employment or trade. Furthermore, some sports have their own specific regulations regarding clearance for participation (e.g., retinal detachment in boxing).
3. *Psychological factors* including values, athletic identity, risk tolerance and readiness or confidence to return to contact or collision sports.^{123 124}
4. *The child or adolescent athlete*. Brain development continues throughout childhood and into young adulthood. In children, progressing to the next age group level in contact or collision sports, especially pre- to post-pubescent children may pose additional risks, especially given the variable age of growth and likelihood of mismatch of size and

strength of players in the competition. Similarly, adolescent athletes who may be progressing to higher levels of competition, including participation in elite pathway programs and/or open-age competitions, may pose higher risks with increased training loads, players of a larger size and/or higher velocity of impacts. The cognitively immature child/adolescent athlete may not yet be capable of adequately understanding the relative risks and benefits of participating in contact or collision sports, and parents/guardians may not be unanimous in their recommendations, influenced by multiple factors including cultural and socio-economic background,¹²⁵ expectations for the child's future professional sports capabilities, vicarious benefits from the child's sporting achievements, and parental anxieties. It is also imperative to appreciate that the priorities in children are cognitive and emotional development and physical maturation. Therefore, if repeat concussions in an individual child limit the ability to return to school and develop appropriately, restrictions on returning to contact or collision sports must be considered. However, considering the important health benefits of implementing a physically active lifestyle in youth,^{19 20} it is equally critical that any child or adolescent restricted from participating in contact or collision sports be encouraged to participate in other non-contact physical or sporting activities.

5. *Geographic variations.* Various legal and cultural frameworks exist in different countries and territories that may impact decisions regarding retiring or discontinuing participation in contact or collision sports.
6. *Para-athlete.* Factors can be even more complex in the assessment of para-sport athletes. Athletic identity is an important consideration as participation in sports may present an opportunity to transcend the bias and stigma the individuals may experience when going about their daily lives. The presence of a neurological finding would not necessarily exclude someone from engaging in sport, but specific circumstances (e.g., seizure disorder in cerebral palsy or for those with neuromuscular conditions and the impact on sport), should be medically cleared prior to participation in sport.

The assessment should be individualized and involve clinicians with expertise in traumatic brain injury and sport, and frequently and preferably will include an independent multi-disciplinary team. The ensuing shared decision-making process should:

- a) Include a discussion that considers the benefits of participation in physical activity including important effects on physical, psychological, social and cognitive well-being weighed against the potential long-term risks that may be associated with concussion and/or exposure to repeated head trauma.¹²⁶ (cite Iverson et al. review 9). Removal from sports even temporarily can exacerbate psychological health issues and providers need to be keenly aware to balance protection with the potential harms of removal. Given the positive benefits of exercise on health, care must be taken to avoid restricting all physical activity. Exercise and physical activity are critical in the prevention and management of many chronic diseases.¹²⁷⁻¹²⁹ Any athlete who ultimately retires from contact or collision sports, should be encouraged to continue non-contact or low-contact physical activity and the benefits explained.
- b) Address the key concerns and correct any misconceptions that the athlete or their family, care givers, or guardians, in the case of athletes under the age of legal consent, may have.
- c) Use language that is appropriate for the health literacy of the individuals based on multiple factors, including their level of education and culture, to reduce the risk that the information is misinterpreted.
- d) The assessment should be carefully documented, as with all aspects of medical care and decision-making.

In many instances, the clinician will help weigh the risks versus benefits and provide their opinion to the athlete, based on their best clinical judgement. Consequently, the advice should be provided by someone with expertise in concussion and head trauma and encompass input by clinicians from disciplines relevant to the athlete's presentation who are familiar with the sport-specific elements of any risk.

Alternatively, it has been suggested that the clinician should not provide specific guidance, but rather provide enough information for the athlete to make an informed decision.

Turner and colleagues suggested that *"the choice to return to play after a concussion or mild*

TBI injury is the athlete's decision once they have (1) recovered from their injury and have the legal capacity to make an informed decision; (2) been medically assessed and (3) been informed of any possible long-term risks in a language that they can understand".¹³⁰

Retirement or discontinuation of a sports career due to injury has potentially life-altering consequences, irrespective of the level of play, and has been associated with an increased risk for mental health symptoms and disorders.^{6 8 131} It is critical that any athlete in this situation be afforded adequate psychological support and follow-up.¹³²

Medical ethical and legal considerations

It is beyond the scope of this paper to provide a comprehensive review of the medico-legal issues associated with an athlete's decision to retire. However, consideration of the ethical and legal duties owed to athletes by medical staff in providing medical advice in the best interests of the athlete patient is an important part of such a process. Different jurisdictions may have their own legislation and case law, and different sports governing bodies may impose their own regulations or guidelines in relation to the management of concussion in their sport. Broadly from a legal perspective, in advising an athlete, it is important that medical staff:

- make the athlete aware of the role(s) they play in the athlete's care, stating clearly if they have or foresee any potential or actual conflicts of interest affecting the decision that might compound informed decision-making by the athlete,
- make best efforts to ensure the athlete is free from coercion or undue influence,
- take all reasonable steps to ensure the athlete has the capacity to make an informed decision,
- make best efforts to ensure that all parties to the discussion are aware of and respect patient confidentiality,
- have understood the athlete patient's own conception of their best interest and informed the athlete of the risks that may be associated with continuing to participate, based on current knowledge of the relevant literature/research,
- use terms and language the athlete understands,
- offer the athlete the opportunity to ask medically relevant questions,

- respond as best they reasonably can to more holistic questions and discuss concerns raised by the athlete,
- understand and respect that, ultimately, it is the competent adult athlete's decision whether to retire,
- consider the use of a cool-off period to allow time for further reflection and consultation where appropriate, and
- make and retain some form of written record that documents the process.

Specific issues arise with child and adolescent athletes. While legislation including the defined age of a "child" may vary across different jurisdictions, in general terms, a parent or guardian must consent to medical treatment of a child. Even under these conditions, however, it is good practice to seek the assent of the child. Many jurisdictions recognise that 'mature minors' may be competent to consent or refuse treatment if they can reasonably understand the information set provided and are not subject to undue influence. Nevertheless, medical staff should be aware that for more significant medical decisions, such as return to play following SRC, it would be prudent to involve a parent/guardian in the discussion and ensure the informed consent of a parent/guardian is obtained.¹³⁰

Further considerations and future research

This systematic review has revealed limited scientific evidence to specifically direct decisions regarding retirement from or participation in contact or collision sport.

To allow us to better inform such decisions future research should focus on

- Improved understanding of potential long-term risks related to concussion and repeated head trauma,
- Robust, clinically relevant biomarkers of brain recovery,
- Identification of genetic risk factors for poor long-term outcomes,
- Understanding of head impact exposure and concussion risk of various sports. This should include all levels in which the game is played as well as cover all competitors (including representation of ethnicity, sociocultural factors, sex and gender, age, disability, level of competition)

Ideally, studies should be prospective cohort designs, that include measurement of factors that may have an impact on outcome including age, sex, sport, culture, genetics, previous injury, co-existing medical conditions, playing conditions (e.g., air quality), and socio-economic status. Also, studies often lack objective and valid measures of exposure to sport and frequently rely on self-reports of exposure or years played. To date most funding is allotted for research that involve shorter durations (i.e., 1-5 year studies) and does not enable prospective follow-up of athletes over time, thus limiting our understanding of the potential factors that are important to consider in decisions related to retirement. The terminal endpoint of most studies is 'return to play clearance' where players would return to the at-risk cohort, however, the short duration of studies precludes longer evaluation of outcomes. In other fields, prospective cohort studies with long-term follow-up have significantly advanced knowledge (e.g., Framingham heart study).

Therefore, future studies should investigate factors associated with unfavourable outcomes after return to play such as early recurrence of concussion or severe injuries leading to chronic impairments and disabilities.

It is also incumbent upon individual sporting bodies, associations, and federations to implement evidence-based rule changes and regulations that assist in reducing the risk of concussion and exposure to repeated head impacts/accelerations.¹³³ (*cite Emery et al review 1*) Such regulations may, in some circumstances, mandate that the athlete does not return to a particular sport (e.g., boxing), when the risks to the individual athlete (based on expert medical opinion) are considered too great.

Review limitations

The main limitation of our systematic review was the large degree of heterogeneity and low methodological quality of the included studies. Furthermore, we only included studies that evaluated the investigation or treatment of SRC. Finally, there was an inherent limitation related to publication bias.

Key recommendations (Box)

1. Decisions regarding retirement or discontinuation from contact or collision sports in the context of SRC or repeated head trauma are complex and multifaceted and should therefore involve clinicians with expertise in traumatic brain injury and sport, and preferably include a multi-disciplinary team (SORT C).
2. The decision-making process should be individualised and incorporate a comprehensive clinical history and physical examination, sequential clinical evaluation, and neuroimaging (when clinically indicated). (SORT C) The assessment should include consideration of psychosocial factors (e.g., values, identity, risk tolerance, psychological readiness), injury-specific factors (e.g., history of prior concussions including number, worsening post-concussive symptoms with subsequent concussions, lower threshold for concussions), sport-specific factors (e.g., type of sport, level played), and any persisting symptoms and/or neuro-cognitive concerns (SORT C).
3. In athletes with persisting or prolonged post-concussion symptoms, neurological abnormalities and/or neurocognitive deficits, it is essential to identify and treat and pre-existing, coexisting or resultant pathology (e.g., mood disorders, post-traumatic headaches/migraines) (SORT B).
4. The clinician should make the athlete aware of the role(s) they play in the athlete's care, stating clearly if they have or foresee any potential or actual conflicts of interest affecting the decision that might compound informed decision-making by the athlete (SORT C).
5. The discussion should provide athletes with the scientific facts and uncertainties of their condition as well as the potential risks associated with returning to their previous sport and/or alternative physical activities with less chance of head trauma or SRC. It should highlight the benefits of regular physical activity and incorporate the athlete's preferences, risk tolerance, and capability to make an informed decision. The discussion should be communicated in a manner that considers the athlete's language preference, education and health literacy, and include others stakeholders (e.g., parents, family) as

indicated. The discussion, including the athlete's understanding of what was communicated, should be carefully documented in the medical records (SORT C).

6. In the child or adolescent athlete, the primary concerns are return to school and maintaining healthy levels of physical activity. This often requires a multidisciplinary process that includes the child/adolescent, parent/caregivers, healthcare providers, school leadership and teachers in the discussions (SORT C).
7. Given its positive benefits on health, care must be taken to avoid restricting all physical activity. Athletes who retire from contact or collision sport should be encouraged to continue non-contact or low-contact physical activity (as medically tolerated) and have the benefits of such engagement explained (SORT B).
8. Further high-quality longitudinal research is needed to inform a risk-benefit model to facilitate informed, individualised decisions about retirement or discontinuation from contact or collision sport.
9. Input and counsel from healthcare professionals knowledgeable regarding brain injury can be helpful in making decisions about retirement. The ultimate choice is one of an informed decision made by the athlete and/or their guardian or parent (SORT C).

CONCLUSION

There was no evidence identified by this systematic review to support the inclusion of any patient-specific, injury-specific or other factors (e.g., imaging findings) as absolute indications for retirement or discontinued participation in contact or collision sports.

REFERENCES

1. Russell ER, Mackay DF, Stewart K, MacLean JA, Pell JP, Stewart W. Association of Field Position and Career Length With Risk of Neurodegenerative Disease in Male Former Professional Soccer Players. *JAMA neurology* 2021;78(9):1057-63. doi: <https://dx.doi.org/10.1001/jamaneurol.2021.2403>
2. Katz DI, Bernick C, Dodick DW, Mez J, Mariani ML, Adler CH, Alosco ML, Balcer LJ, Banks SJ, Barr WB, Brody DL, Cantu RC, Dams-O'Connor K, Geda YE, Jordan BD, McAllister TW, Peskind ER, Petersen RC, Wethe JV, Zafonte RD, Foley EM, Babcock DJ, Koroshetz WJ, Tripodis Y, McKee AC, Shenton ME, Cummings JL, Reiman EM, Stern RA. National Institute of Neurological Disorders and Stroke Consensus Diagnostic Criteria for Traumatic Encephalopathy Syndrome. *Neurology* 2021;96(18):848-63. doi: 10.1212/wnl.00000000000011850
3. Montenegro PH, Alosco ML, Martin BM, Daneshvar DH, Mez J, Chaisson CE, Nowinski CJ, Au R, McKee AC, Cantu RC, McClean MD, Stern RA, Tripodis Y. Cumulative Head Impact Exposure Predicts Later-Life Depression, Apathy, Executive Dysfunction, and Cognitive Impairment in Former High School and College Football Players. *Journal of neurotrauma* 2017;34(2):328-40. doi: <https://dx.doi.org/10.1089/neu.2016.4413>
4. Caron JG, Bloom GA, Johnston KM, Sabiston CM. Effects of multiple concussions on retired national hockey league players. *Journal of sport & exercise psychology* 2013;35(2):168-79.
5. Monaco MR, Brewer BW, Van Raalte JL, May CN. How Many Concussions Would It Take for Athletes to Choose to Discontinue Participation in Their Primary Sport? *International journal of environmental research and public health* 2021;18(4) doi: <https://dx.doi.org/10.3390/ijerph18041582>
6. Reardon CL, Hainline B, Aron CM, Baron D, Baum AL, Bindra A, Budgett R, Campriani N, Castaldelli-Maia JM, Currie A, Derevensky JL, Glick ID, Gorczynski P, Gouttebauge V, Grandner MA, Han DH, McDuff D, Mountjoy M, Polat A, Purcell R, Putukian M, Rice S, Sills A, Stull T, Swartz L, Zhu LJ, Engebretsen L. Mental health in elite athletes: International Olympic Committee consensus statement (2019). *British Journal of Sports Medicine* 2019;53(11):667-99. doi: 10.1136/bjsports-2019-100715

7. Gouttebarga V, Castaldelli-Maia JM, Gorczynski P, Hainline B, Hitchcock ME, Kerkhoffs GM, Rice SM, Reardon CL. Occurrence of mental health symptoms and disorders in current and former elite athletes: a systematic review and meta-analysis. *British Journal of Sports Medicine* 2019;53(11):700-06. doi: 10.1136/bjsports-2019-100671
8. Rice SM, Purcell R, De Silva S, Mawren D, McGorry PD, Parker AG. The Mental Health of Elite Athletes: A Narrative Systematic Review. *Sports Medicine* 2016;46(9):1333-53. doi: 10.1007/s40279-016-0492-2
9. Baggish AL, Ackerman MJ, Putukian M, Lampert R. Shared Decision Making for Athletes with Cardiovascular Disease: Practical Considerations. *Current sports medicine reports* 2019;18(3):76-81. doi: 10.1249/jsr.0000000000000575
10. Barry MJ. Shared decision making: informing and involving patients to do the right thing in health care. *J Ambul Care Manage* 2012;35(2):90-8. doi: 10.1097/JAC.0b013e318249482f [published Online First: 2012/03/15]
11. Barry MJ, Edgman-Levitan S. Shared Decision Making — The Pinnacle of Patient-Centered Care. *New England Journal of Medicine* 2012;366(9):780-81. doi: 10.1056/NEJMp1109283
12. Creighton DW, Shrier I, Shultz R, Meeuwisse WH, Matheson GO. Return-to-play in sport: a decision-based model. *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine* 2010;20(5):379-85. doi: 10.1097/jsm.0b013e3181f3c0fe
13. Herring SA, Kibler WB, Putukian M. The team physician and the return-to-play decision: a consensus statement-2012 update. *Med Sci Sports Exerc* 2012;44(12):2446-8. doi: 10.1249/MSS.0b013e3182750534 [published Online First: 2012/11/20]
14. Cantu RC, Register-Mihalik JK. Considerations for return-to-play and retirement decisions after concussion. *PM & R : the journal of injury, function, and rehabilitation* 2011;3(10 Suppl 2):S440-4. doi: <https://dx.doi.org/10.1016/j.pmrj.2011.07.013>
15. Concannon LG, Kaufman MS, Herring SA. The million dollar question: When should an athlete retire after concussion? *Current sports medicine reports* 2014;13(6):365-9. doi: <https://dx.doi.org/10.1249/JSR.0000000000000098>
16. Davis-Hayes C, Baker DR, Bottiglieri TS, Levine WN, Desai N, Gossett JD, Noble JM. Medical retirement from sport after concussions: A practical guide for a difficult

discussion. *Neurology Clinical practice* 2018;8(1):40-47. doi:

<https://dx.doi.org/10.1212/CPJ.0000000000000424>

17. Rivara FP, Tennyson R, Mills B, Browd SR, Emery CA, Gioia G, Giza CC, Herring S, Janz KF, Labella C, Valovich McLeod T, Meehan W, Patricios J. Consensus Statement on Sports-Related Concussions in Youth Sports Using a Modified Delphi Approach. *JAMA Pediatrics* 2020;174(1):79-85. doi: <http://dx.doi.org/10.1001/jamapediatrics.2019.4006>
18. Northey JM, Cherbuin N, Pumpa KL, Smee DJ, Rattray B. Exercise interventions for cognitive function in adults older than 50: a systematic review with meta-analysis. *Br J Sports Med* 2018;52(3):154-60. doi: 10.1136/bjsports-2016-096587 [published Online First: 2017/04/26]
19. Rasmussen M, Laumann K. The academic and psychological benefits of exercise in healthy children and adolescents. *European Journal of Psychology of Education* 2013;28(3):945-62. doi: 10.1007/s10212-012-0148-z
20. Bull FC, Al-Ansari SS, Biddle S, Borodulin K, Buman MP, Cardon G, Carty C, Chaput J-P, Chastin S, Chou R, Dempsey PC, DiPietro L, Ekelund U, Firth J, Friedenreich CM, Garcia L, Gichu M, Jago R, Katzmarzyk PT, Lambert E, Leitzmann M, Milton K, Ortega FB, Ranasinghe C, Stamatakis E, Tiedemann A, Troiano RP, van der Ploeg HP, Wari V, Willumsen JF. World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *British Journal of Sports Medicine* 2020;54(24):1451-62. doi: 10.1136/bjsports-2020-102955
21. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R, Glanville J, Grimshaw JM, Hróbjartsson A, Lalu MM, Li T, Loder EW, Mayo-Wilson E, McDonald S, McGuinness LA, Stewart LA, Thomas J, Tricco AC, Welch VA, Whiting P, Moher D. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Systematic Reviews* 2021;10(1):89. doi: 10.1186/s13643-021-01626-4
22. Schneider KJ, Patricios J, Meeuwisse W, al. e. Amsterdam 2022 process: a summary of the methodology for the International Consensus on Concussion in Sport. *British Journal of Sports Medicine* 2023 doi: 10.1136/bjsports-2022-106663
23. Cooper C, Dawson S, Peters J, Varley-Campbell J, Cockcroft E, Hendon J, Churchill R. Revisiting the need for a literature search narrative: A brief methodological note. *Res*

- Synth Methods* 2018;9(3):361-65. doi: 10.1002/jrsm.1315 [published Online First: 2018/07/22]
24. Ebell MH, Siwek J, Weiss BD, Woolf SH, Susman J, Ewigman B, Bowman M. Strength of recommendation taxonomy (SORT): a patient-centered approach to grading evidence in the medical literature. *Am Fam Physician* 2004;69(3):548-56. [published Online First: 2004/02/20]
25. Arnold JT, Franklin EV, Baker ZG, Abowd M, Santana JA. Association Between Fear of Pain and Sports-Related Concussion Recovery in a Pediatric Population. *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine* 2021(b9t, 9103300) doi: <https://dx.doi.org/10.1097/JSM.0000000000000951>
26. Kontos AP, Elbin RJ, Sufrinko A, Marchetti G, Holland CL, Collins MW. Recovery Following Sport-Related Concussion: Integrating Pre- and Postinjury Factors Into Multidisciplinary Care. *The Journal of head trauma rehabilitation* 2019(8702552, c3t) doi: <https://dx.doi.org/10.1097/HTR.0000000000000536>
27. Kriz PK, Stein C, Kent J, Ruggieri D, Dolan E, O'Brien M, Meehan WP, 3rd. Physical Maturity and Concussion Symptom Duration among Adolescent Ice Hockey Players. *The Journal of pediatrics* 2016;171(jlz, 0375410):234-2. doi: <https://dx.doi.org/10.1016/j.jpeds.2015.12.006>
28. Aggarwal SS, Ott SD, Padhye NS, Meininger JC, Armstrong TS. Clinical and demographic predictors of concussion resolution in adolescents: A retrospective study. *Applied Neuropsychology: Child* 2017:1-11. doi: 10.1080/21622965.2017.1381099
29. Asplund CA, McKeag DB, Olsen CH. Sport-related concussion: factors associated with prolonged return to play. *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine* 2004;14(6):339-43.
30. Bramley H, Henson A, Lewis MM, Kong L, Stetter C, Silvis M. Sleep Disturbance Following Concussion Is a Risk Factor for a Prolonged Recovery. *Clinical pediatrics* 2017;56(14):1280-85. doi: <https://dx.doi.org/10.1177/0009922816681603>
31. Bressan S, Clarke CJ, Anderson V, Takagi M, Hearps SJC, Rausa V, Anderson N, Doyle M, Dunne K, Oakley E, Davis GA, Babl FE, Hearps SJC. Use of the sport concussion assessment tools in the emergency department to predict persistent post-concussive

- symptoms in children. *Journal of Paediatrics & Child Health* 2020;56(8):1249-56. doi: 10.1111/jpc.14910
32. Kara S, Crosswell H, Forch K, Cavadino A, McGeown J, Fulcher M. Less Than Half of Patients Recover Within 2 Weeks of Injury After a Sports-Related Mild Traumatic Brain Injury: A 2-Year Prospective Study. *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine* 2020;30(2):96-101. doi: <https://dx.doi.org/10.1097/JSM.0000000000000811>
33. Lau BC, Collins MW, Lovell MR. Cutoff scores in neurocognitive testing and symptom clusters that predict protracted recovery from concussions in high school athletes. *Neurosurgery* 2012;70(2):371-79. doi: <https://dx.doi.org/10.1227/NEU.0b013e31823150f0>
34. McCrea M, Guskiewicz K, Randolph C, Barr WB, Hammeke TA, Marshall SW, Powell MR, Woo Ahn K, Wang Y, Kelly JP. Incidence, clinical course, and predictors of prolonged recovery time following sport-related concussion in high school and college athletes. *Journal of the International Neuropsychological Society : JINS* 2013;19(1):22-33. doi: <https://dx.doi.org/10.1017/S1355617712000872>
35. Meehan WP, 3rd, Mannix RC, Stracciolini A, Elbin RJ, Collins MW. Symptom severity predicts prolonged recovery after sport-related concussion, but age and amnesia do not. *The Journal of pediatrics* 2013;163(3):721-5. doi: <https://dx.doi.org/10.1016/j.jpeds.2013.03.012>
36. Morgan CD, Zuckerman SL, Lee YM, King L, Beaird S, Sills AK, Solomon GS. Predictors of postconcussion syndrome after sports-related concussion in young athletes: a matched case-control study. *Journal of neurosurgery Pediatrics* 2015;15(6):589-98. doi: <https://dx.doi.org/10.3171/2014.10.PEDS14356>
37. Bock S, Grim R, Barron TF, Wagenheim A, Hu YE, Hendell M, Deitch J, Deibert E. Factors associated with delayed recovery in athletes with concussion treated at a pediatric neurology concussion clinic. *Child's Nervous System* 2015;31(11):2111-16. doi: <http://dx.doi.org/10.1007/s00381-015-2846-8>
38. Desai N, Wiebe DJ, Corwin DJ, Lockyer JE, Grady MF, Master CL. Factors Affecting Recovery Trajectories in Pediatric Female Concussion. *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine* 2019;29(5):361-67. doi: <https://dx.doi.org/10.1097/JSM.0000000000000646>

39. Elbin RJ, Sufrinko A, Schatz P, French J, Henry L, Burkhart S, Collins MW, Kontos AP. Removal From Play After Concussion and Recovery Time. *Pediatrics* 2016;138(3) doi: <https://dx.doi.org/10.1542/peds.2016-0910>
40. Fehr SD, Nelson LD, Scharer KR, Traudt EA, Veenstra JM, Tarima SS, Liu X-C, Walter KD. Risk Factors for Prolonged Symptoms of Mild Traumatic Brain Injury: A Pediatric Sports Concussion Clinic Cohort. *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine* 2019;29(1):11-17. doi: <https://dx.doi.org/10.1097/JSM.0000000000000494>
41. Heyer GL, Schaffer CE, Rose SC, Young JA, McNally KA, Fischer AN. Specific Factors Influence Postconcussion Symptom Duration among Youth Referred to a Sports Concussion Clinic. *The Journal of pediatrics* 2016;174(jlz, 0375410):33-38.e2. doi: <https://dx.doi.org/10.1016/j.jpeds.2016.03.014>
42. McGeown JP, Kara S, Fulcher M, Crosswell H, Borotkanics R, Hume PA, Quarrie KL, Theadom A. Predicting Sport-related mTBI Symptom Resolution Trajectory Using Initial Clinical Assessment Findings: A Retrospective Cohort Study. *Sports Medicine* 2020;50(6):1191-202.
43. Miller JH, Gill C, Kuhn EN, Rocque BG, Menendez JY, O'Neill JA, Agee BS, Brown ST, Crowther M, Davis RD, Ferguson D, Johnston JM. Predictors of delayed recovery following pediatric sports-related concussion: a case-control study. *Journal of neurosurgery Pediatrics* 2016;17(4):491-6. doi: <https://dx.doi.org/10.3171/2015.8.PEDS14332>
44. Thomas DJ, Coxe K, Li H, Pommering TL, Young JA, Smith GA, Yang J. Length of Recovery From Sports-Related Concussions in Pediatric Patients Treated at Concussion Clinics. *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine* 2018;28(1):56-63. doi: <https://dx.doi.org/10.1097/JSM.0000000000000413>
45. Vedung F, Hanni S, Tegner Y, Johansson J, Marklund N. Concussion incidence and recovery in Swedish elite soccer - Prolonged recovery in female players. *Scandinavian journal of medicine & science in sports* 2020;30(5):947-57. doi: <https://dx.doi.org/10.1111/sms.13644>
46. Asken BM, McCrea MA, Clugston JR, Snyder AR, Houck ZM, Bauer RM. "Playing Through It": Delayed Reporting and Removal From Athletic Activity After Concussion Predicts

- Prolonged Recovery. *Journal of athletic training* 2016;51(4):329-35. doi: <https://dx.doi.org/10.4085/1062-6050-51.5.02>
47. Berz K, Divine J, Foss KB, Heyl R, Ford KR, Myer GD. Sex-specific differences in the severity of symptoms and recovery rate following sports-related concussion in young athletes. *The Physician and sportsmedicine* 2013;41(2):58-63. doi: <https://dx.doi.org/10.3810/psm.2013.05.2015>
48. Corbin-Berrigan L-A, Gagnon I. Postconcussion Symptoms as a Marker of Delayed Recovery in Children and Youth Who Recently Sustained a Concussion: A Brief Report. *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine* 2017;27(3):325-27. doi: <https://dx.doi.org/10.1097/JSM.0000000000000355>
49. Howell DR, O'Brien MJ, Beasley MA, Mannix RC, Meehan WP, 3rd. Initial somatic symptoms are associated with prolonged symptom duration following concussion in adolescents. *Acta paediatrica (Oslo, Norway : 1992)* 2016;105(9):e426-32. doi: <https://dx.doi.org/10.1111/apa.13486>
50. Ono KE, Burns TG, Bearden DJ, McManus SM, King H, Reisner A. Sex-Based Differences as a Predictor of Recovery Trajectories in Young Athletes After a Sports-Related Concussion. *The American journal of sports medicine* 2016;44(3):748-52. doi: <https://dx.doi.org/10.1177/0363546515617746>
51. Schmidt JD, Rawlins MLW, Lynall RC, D'Lauro C, Clugston JR, McAllister TW, McCrea M, Broglio SP, Investigators CC. Medical Disqualification Following Concussion in Collegiate Student-Athletes: Findings from the CARE Consortium. *Sports medicine (Auckland, NZ)* 2020;50(10):1843-55. doi: <https://dx.doi.org/10.1007/s40279-020-01302-y>
52. Wilmoth K, Curcio N, Tarkenton T, Meredith-Duliba T, Tan A, Didehbani N, Hynan LS, Miller SM, Bell KR, Cullum CM. Utility of Brief Psychological Measures for Prediction of Prolonged Symptom Clearance in Concussed Student Athletes. *Archives of clinical neuropsychology : the official journal of the National Academy of Neuropsychologists* 2021;36(3):430-36. doi: <https://dx.doi.org/10.1093/arclin/acz061>
53. Barker T, Russo SA, Barker G, Rice MA, Jr., Jeffrey MG, Broderick G, Craddock TJA. A case matched study examining the reliability of using ImpACT to assess effects of multiple

- concussions. *BMC psychology* 2017;5(1):14. doi: <https://dx.doi.org/10.1186/s40359-017-0184-1>
54. Broglio SP, McAllister T, Katz BP, LaPradd M, Zhou W, McCrea MA, Investigators CC. The Natural History of Sport-Related Concussion in Collegiate Athletes: Findings from the NCAA-DoD CARE Consortium. *Sports medicine (Auckland, NZ)* 2022;52(2):403-15. doi: <https://dx.doi.org/10.1007/s40279-021-01541-7>
55. Covassin T, Stearne D, Elbin R. Concussion history and postconcussion neurocognitive performance and symptoms in collegiate athletes. *Journal of athletic training* 2008;43(2):119-24. doi: <https://dx.doi.org/10.4085/1062-6050-43.2.119>
56. Guskiewicz KM, McCrea M, Marshall SW, Cantu RC, Randolph C, Barr W, Onate JA, Kelly JP. Cumulative effects associated with recurrent concussion in collegiate football players: the NCAA Concussion Study. *JAMA* 2003;290(19):2549-55.
57. Guskiewicz KM, Marshall SW, Bailes J, McCrea M, Cantu RC, Randolph C, Jordan BD. Association between recurrent concussion and late-life cognitive impairment in retired professional football players. *Neurosurgery* 2005;57(4):719-26.
58. Haider MN, Cunningham A, Darling S, Suffoletto HN, Freitas MS, Jain RK, Willer B, Leddy JJ. Derivation of the Buffalo Concussion Physical Examination risk of delayed recovery (RDR) score to identify children at risk for persistent postconcussive symptoms. *British journal of sports medicine* 2021;55(24):1427-33. doi: <https://dx.doi.org/10.1136/bjsports-2020-103690>
59. Wang EX, Hwang CE, Nguyen JN, Segovia NA, Abrams GD, Kussman A. Factors Associated With a Prolonged Time to Return to Play After a Concussion. *The American journal of sports medicine* 2022(7609541, 3yr):3635465221083646. doi: <https://dx.doi.org/10.1177/03635465221083646>
60. Cook NE, Iverson GL, Maxwell B, Zafonte R, Berkner PD. Adolescents With ADHD Do Not Take Longer to Recover From Concussion. *Frontiers in pediatrics* 2020;8(101615492):606879. doi: <https://dx.doi.org/10.3389/fped.2020.606879>
61. Currie DW, Comstock RD, Fields SK, Cantu RC. A Paired Comparison of Initial and Recurrent Concussions Sustained by US High School Athletes Within a Single Athletic Season. *The Journal of head trauma rehabilitation* 2017;32(2):90-97. doi: <https://dx.doi.org/10.1097/HTR.0000000000000240>

62. Hannah TC, Li AY, Spiera Z, Kuohn L, Dai J, McAuley F, Ali M, Durbin JR, Dreher N, Marayati NF, Gometz A, Lovell M, Choudhri T. Sex-Related Differences in the Incidence, Severity, and Recovery of Concussion in Adolescent Student-Athletes Between 2009 and 2019. *American Journal of Sports Medicine* 2021;49(7):1929-37.
63. Kinney AR, Anderson D, Stearns-Yoder KA, Brenner LA, Forster JE. Risk and recovery among high school athletes who sustained at least one sports-related concussion. *Concussion (London, England)* 2020;5(2):CNC72. doi: <https://dx.doi.org/10.2217/cnc-2020-0002>
64. Lau BC, Kontos AP, Collins MW, Mucha A, Lovell MR. Which on-field signs/symptoms predict protracted recovery from sport-related concussion among high school football players? *The American journal of sports medicine* 2011;39(11):2311-8. doi: <https://dx.doi.org/10.1177/0363546511410655>
65. Mautner K, Sussman WI, Axtman M, Al-Farsi Y, Al-Adawi S. Relationship of Attention Deficit Hyperactivity Disorder and Postconcussion Recovery in Youth Athletes. *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine* 2015;25(4):355-60. doi: <https://dx.doi.org/10.1097/JSM.0000000000000151>
66. Teel EF, Marshall SW, Shankar V, McCrea M, Guskiewicz KM. Predicting Recovery Patterns After Sport-Related Concussion. *Journal of Athletic Training (Allen Press)* 2017;52(3):288-98.
67. Terry DP, Huebschmann NA, Maxwell BA, Cook NE, Mannix R, Zafonte R, Seifert T, Berkner PD, Iverson GL. Preinjury Migraine History as a Risk Factor for Prolonged Return to School and Sports following Concussion. *Journal of Neurotrauma* 2019;36(1):142-51. doi: <http://dx.doi.org/10.1089/neu.2017.5443>
68. Emery CA, Warriyar KV V, Black AM, Palacios-Derflingher L, Sick S, Debert C, Brooks BL, Yeates KO, Mrazik M, Lebrun C, Hagel BE, Dukelow S, Schneider KJ. Factors Associated With Clinical Recovery After Concussion in Youth Ice Hockey Players. *Orthopaedic Journal of Sports Medicine* 2021;9(5):23259671211013370. doi: 10.1177/23259671211013370
69. Ali M, Asghar N, Li A, Hannah T, Spiera Z, Marayati NF, Dreher N, Durbin J, Gometz A, Lovell M, Choudhri T. Incidence of concussion and recovery of neurocognitive dysfunction on ImpACT assessment among youth athletes with premorbid

- depression or anxiety taking antidepressants. *Journal of neurosurgery Pediatrics* 2021(101463759):1-7. doi: <https://dx.doi.org/10.3171/2020.11.PEDS20821>
70. Corwin DJ, Zonfrillo MR, Master CL, Arbogast KB, Grady MF, Robinson RL, Goodman AM, Wiebe DJ. Characteristics of prolonged concussion recovery in a pediatric subspecialty referral population. *The Journal of pediatrics* 2014;165(6):1207-15. doi: <https://dx.doi.org/10.1016/j.jpeds.2014.08.034>
71. Legarreta AD, Brett BL, Solomon GS, Zuckerman SL. The role of family and personal psychiatric history in postconcussion syndrome following sport-related concussion: a story of compounding risk. *Journal of neurosurgery Pediatrics* 2018;22(3):238-43. doi: <https://dx.doi.org/10.3171/2018.3.PEDS1850>
72. Martin AK, Petersen AJ, Sesma HW, Koolmo MB, Ingram KM, Slifko KB, Nguyen VN, Doss RC, Linabery AM. Concussion symptomology and recovery in children and adolescents with pre-existing anxiety. *Journal of neurology, neurosurgery, and psychiatry* 2020;91(10):1060-66. doi: <https://dx.doi.org/10.1136/jnnp-2020-323137>
73. Kent M, Brilliant A, Erickson K, Meehan W, Howell D. Symptom Presentation After Concussion and Pre-existing Anxiety Among Youth Athletes. *International journal of sports medicine* 2020;41(10):682-87. doi: <https://dx.doi.org/10.1055/a-1107-3025>
74. Teel EF, Caron JG, Gagnon IJ. Higher parental stress is significantly related to longer clinical recovery times in concussed children: A mixed-methods study. *Journal of Science and Medicine in Sport* 2022;25(2):108-14. doi: <https://dx.doi.org/10.1016/j.jsams.2021.08.014>
75. Charek DB, Elbin RJ, Sufrinko A, Schatz P, D'Amico NR, Collins MW, Kontos AP. Preliminary Evidence of a Dose-Response for Continuing to Play on Recovery Time After Concussion. *The Journal of head trauma rehabilitation* 2020;35(2):85-91. doi: <https://dx.doi.org/10.1097/HTR.0000000000000476>
76. Zynda AJ, Worrall HM, Sabatino MJ, Ellis HB, Chung JS, Cullum CM, Miller SM. Continued play following adolescent sport-related concussion: Prospective data from the North Texas Concussion Registry (ConTex). *Applied neuropsychology Child* 2021(101584990):1-12. doi: <https://dx.doi.org/10.1080/21622965.2021.1957677>
77. Kontos AP, Jorgensen-Wagers K, Trbovich AM, Ernst N, Emami K, Gillie B, French J, Holland C, Elbin RJ, Collins MW. Association of Time since Injury to the First Clinic

- Visit with Recovery Following Concussion. *JAMA Neurology* 2020;77(4):435-40. doi: <http://dx.doi.org/10.1001/jamaneurol.2019.4552>
78. Davis-Hayes C, Gossett JD, Levine WN, Shams T, Harada J, Mitnick J, Noble J. Sex-specific Outcomes and Predictors of Concussion Recovery. *The Journal of the American Academy of Orthopaedic Surgeons* 2017;25(12):818-28. doi: <https://dx.doi.org/10.5435/JAAOS-D-17-00276>
79. Hiploylee C, Dufort PA, Davis HS, Wennberg RA, Tartaglia MC, Mikulis D, Hazrati L-N, Tator CH. Longitudinal Study of Postconcussion Syndrome: Not Everyone Recovers. *Journal of neurotrauma* 2017;34(8):1511-23. doi: <https://dx.doi.org/10.1089/neu.2016.4677>
80. Putukian M, Riegler K, Amalfe S, Bruce J, Echemendia R. Preinjury and Postinjury Factors That Predict Sports-Related Concussion and Clinical Recovery Time. *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine* 2018(b9t, 9103300) doi: <https://dx.doi.org/10.1097/JSM.0000000000000705>
81. Worts PR, Haider MN, Mason JR, Schatz P. Norm-Based Cutoffs as Predictors of Prolonged Recovery After Adolescent Sport-Related Concussion. *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine* 2021(b9t, 9103300) doi: <https://dx.doi.org/10.1097/JSM.0000000000000952>
82. Kontos AP, Elbin RJ, Lau B, Simensky S, Freund B, French J, Collins MW. Posttraumatic migraine as a predictor of recovery and cognitive impairment after sport-related concussion. *The American journal of sports medicine* 2013;41(7):1497-504. doi: <https://dx.doi.org/10.1177/0363546513488751>
83. Eagle SR, Womble MN, Elbin RJ, Pan R, Collins MW, Kontos AP. Concussion Symptom Cutoffs for Identification and Prognosis of Sports-Related Concussion: Role of Time Since Injury. *The American journal of sports medicine* 2020;48(10):2544-51. doi: <https://dx.doi.org/10.1177/0363546520937291>
84. Ernst N, Eagle S, Trbovich A, Kissinger-Knox A, Bitzer H, Kontos AP. Lower post-injury psychological resilience is associated with increased recovery time and symptom burden following sport-related concussion. *Applied neuropsychology Child* 2021(101584990):1-8. doi: <https://dx.doi.org/10.1080/21622965.2021.1964966>
85. Chung JS, Zynda AJ, Didehbani N, Hicks C, Hynan LS, Miller SM, Bell KR, Cullum CM. Association Between Sleep Quality and Recovery Following Sport-Related Concussion

- in Pediatrics. *Journal of child neurology* 2019;34(11):639-45. doi:
<https://dx.doi.org/10.1177/0883073819849741>
86. Murdaugh DL, Ono KE, Reisner A, Burns TG. Assessment of Sleep Quantity and Sleep Disturbances During Recovery From Sports-Related Concussion in Youth Athletes. *Archives of physical medicine and rehabilitation* 2018;99(5):960-66. doi:
<https://dx.doi.org/10.1016/j.apmr.2018.01.005>
87. Anzalone AJ, Blueitt D, Case T, McGuffin T, Pollard K, Garrison JC, Jones MT, Pavur R, Turner S, Oliver JM. A Positive Vestibular/Ocular Motor Screening (VOMS) Is Associated With Increased Recovery Time After Sports-Related Concussion in Youth and Adolescent Athletes. *The American journal of sports medicine* 2017;45(2):474-79. doi: <https://dx.doi.org/10.1177/0363546516668624>
88. Corwin DJ, Wiebe DJ, Zonfrillo MR, Grady MF, Robinson RL, Goodman AM, Master CL. Vestibular deficits following youth concussion. *Journal of Pediatrics* 2015;166(5):1221-25. doi: <http://dx.doi.org/10.1016/j.jpeds.2015.01.039>
89. Ellis MJ, Cordingley DM, Vis S, Reimer KM, Leiter J, Russell K. Clinical predictors of vestibulo-ocular dysfunction in pediatric sports-related concussion. *Journal of neurosurgery Pediatrics* 2017;19(1):38-45. doi:
<https://dx.doi.org/10.3171/2016.7.PEDS16310>
90. Glendon K, Blenkinsop G, Belli A, Pain M. Does Vestibular-Ocular-Motor (VOM) Impairment Affect Time to Return to Play, Symptom Severity, Neurocognition and Academic Ability in Student-Athletes following acute Concussion? *Brain Injury* 2021;35(7):788-97. doi: 10.1080/02699052.2021.1911001
91. Knell G, Caze T, Burkhart SO. Evaluation of the vestibular and ocular motor screening (VOMS) as a prognostic tool for protracted recovery following paediatric sports-related concussion. *BMJ open sport & exercise medicine* 2021;7(1):e000970. doi:
<https://dx.doi.org/10.1136/bmjsem-2020-000970>
92. Martinez C, Christopherson Z, Lake A, Myers H, Bytomski JR, Butler RJ, Cook CE. Clinical examination factors that predict delayed recovery in individuals with concussion. *Archives of physiotherapy* 2020;10(101688712):10. doi:
<https://dx.doi.org/10.1186/s40945-020-00081-z>
93. Price AM, Knell G, Caze TJ, 2nd, Abt JP, Loveland D, Burkhart SO. Exploring Vestibular/Ocular and Cognitive Dysfunction as Prognostic Factors for Protracted

- Recovery in Sports-Related Concussion Patients Aged 8 to 12 Years. *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine* 2021;(b9t, 9103300) doi: <https://dx.doi.org/10.1097/JSM.0000000000000975>
94. Sinnott AM, Elbin RJ, Collins MW, Reeves VL, Holland CL, Kontos AP. Persistent vestibular-ocular impairment following concussion in adolescents. *Journal of Science & Medicine in Sport* 2019;22(12):1292-97. doi: 10.1016/j.jsams.2019.08.004
95. Whitney SL, Eagle SR, Marchetti G, Mucha A, Collins MW, Kontos AP, Investigators CC. Association of acute vestibular/ocular motor screening scores to prolonged recovery in collegiate athletes following sport-related concussion. *Brain injury* 2020;34(6):840-45. doi: <https://dx.doi.org/10.1080/02699052.2020.1755055>
96. DuPrey KM, Webner D, Lyons A, Kucuk CH, Ellis JT, Cronholm PF. Convergence Insufficiency Identifies Athletes at Risk of Prolonged Recovery From Sport-Related Concussion. *The American journal of sports medicine* 2017;45(10):2388-93. doi: <https://dx.doi.org/10.1177/0363546517705640>
97. Howell DR, Oldham J, Lanois C, Koerte I, Lin AP, Berkstresser B, Wang F, Meehan WP. Dual-Task Gait Recovery after Concussion among Female and Male Collegiate Athletes. *Medicine and science in sports and exercise* 2020;52(5):1015-21. doi: <https://dx.doi.org/10.1249/MSS.0000000000002225>
98. Iverson G. Predicting slow recovery from sport-related concussion: The new simple-complex distinction. *Clinical Journal of Sport Medicine* 2007;17(1):31-37. doi: <http://dx.doi.org/10.1097/JSM.0b013e3180305e4d>
99. Orr R, Bogg T, Fyffe A, Lam LT, Browne GJ. Graded Exercise Testing Predicts Recovery Trajectory of Concussion in Children and Adolescents. *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine* 2021;31(1):23-30. doi: <https://dx.doi.org/10.1097/JSM.0000000000000683>
100. Howell DR, Oldham J, Lanois C, Koerte I, Lin AP, Berkstresser B, Wang F, Meehan Iii WP. Dual-Task Gait Recovery after Concussion among Female and Male Collegiate Athletes. *Medicine & Science in Sports & Exercise* 2020;52(5):1015-21.
101. Howell DR, Zemek R, Brilliant AN, Mannix RC, Master CL, Meehan WP, 3rd. Identifying Persistent Postconcussion Symptom Risk in a Pediatric Sports Medicine Clinic. *The American journal of sports medicine* 2018;46(13):3254-61. doi: <https://dx.doi.org/10.1177/0363546518796830>

102. McDevitt J, Rubin LH, De Simone FI, Phillips J, Langford D. Association between (GT)n Promoter Polymorphism and Recovery from Concussion: A Pilot Study. *Journal of neurotrauma* 2020;37(10):1204-10. doi: <https://dx.doi.org/10.1089/neu.2019.6590>
103. Mondello S, Guedes VA, Lai C, Jeromin A, Bazarian JJ, Gill JM. Sex Differences in Circulating T-Tau Trajectories After Sports-Concussion and Correlation With Outcome. *Frontiers in neurology* 2020;11(101546899):651. doi: <https://dx.doi.org/10.3389/fneur.2020.00651>
104. Fedorchak G, Rangnekar A, Onks C, Loeffert AC, Loeffert J, Olympia RP, DeVita S, Leddy J, Haider MN, Roberts A, Rieger J, Uhlig T, Monteith C, Middleton F, Zuckerman SL, Lee T, Yeates KO, Mannix R, Hicks S. Saliva RNA biomarkers predict concussion duration and detect symptom recovery: a comparison with balance and cognitive testing. *Journal of neurology* 2021;268(11):4349-61. doi: <https://dx.doi.org/10.1007/s00415-021-10566-x>
105. Shahim P, Tegner Y, Gustafsson B, Gren M, Arlig J, Olsson M, Lehto N, Engstrom A, Hoglund K, Portelius E, Zetterberg H, Blennow K. Neurochemical Aftermath of Repetitive Mild Traumatic Brain Injury. *JAMA neurology* 2016;73(11):1308-15. doi: <https://dx.doi.org/10.1001/jamaneurol.2016.2038>
106. Brett BL, Bobholz SA, Espana LY, Huber DL, Mayer AR, Harezlak J, Broglio SP, McAllister TW, McCrea MA, Meier TB, Investigators CC. Cumulative Effects of Prior Concussion and Primary Sport Participation on Brain Morphometry in Collegiate Athletes: A Study From the NCAA-DoD CARE Consortium. *Frontiers in neurology* 2020;11(101546899):673. doi: <https://dx.doi.org/10.3389/fneur.2020.00673>
107. Panwar J, Hsu CC-T, Tator CH, Mikulis D. Magnetic Resonance Imaging Criteria for Post-Concussion Syndrome: A Study of 127 Post-Concussion Syndrome Patients. *Journal of neurotrauma* 2020;37(10):1190-96. doi: <https://dx.doi.org/10.1089/neu.2019.6809>
108. Bonow RH, Friedman SD, Perez FA, Ellenbogen RG, Browd SR, Mac Donald CL, Vavilala MS, Rivara FP. Prevalence of Abnormal Magnetic Resonance Imaging Findings in Children with Persistent Symptoms after Pediatric Sports-Related Concussion. *Journal of neurotrauma* 2017;34(19):2706-12. doi: <https://dx.doi.org/10.1089/neu.2017.4970>

109. Barlow KM, Iyer K, Yan T, Scurfield A, Carlson H, Wang Y. Cerebral Blood Flow Predicts Recovery in Children with Persistent Post-Concussion Symptoms after Mild Traumatic Brain Injury. *Journal of Neurotrauma* 2021;38(16):2275-83. doi: <http://dx.doi.org/10.1089/neu.2020.7566>
110. Bartnik-Olson BL, Holshouser B, Wang H, Grube M, Tong K, Wong V, Ashwal S. Impaired neurovascular unit function contributes to persistent symptoms after concussion: a pilot study. *Journal of neurotrauma* 2014;31(17):1497-506. doi: <https://dx.doi.org/10.1089/neu.2013.3213>
111. Churchill NW, Hutchison MG, Graham SJ, Schweizer TA. Cerebrovascular Reactivity After Sport Concussion: From Acute Injury to 1 Year After Medical Clearance. *Frontiers in neurology* 2020;11(101546899):558. doi: <https://dx.doi.org/10.3389/fneur.2020.00558>
112. Churchill NW, Hutchison MG, Graham SJ, Schweizer TA. Long-term changes in the small-world organization of brain networks after concussion. *Scientific reports* 2021;11(1):6862. doi: <https://dx.doi.org/10.1038/s41598-021-85811-4>
113. Cubon VA, Putukian M, Boyer C, Dettwiler A. A diffusion tensor imaging study on the white matter skeleton in individuals with sports-related concussion. *Journal of neurotrauma* 2011;28(2):189-201. doi: <https://dx.doi.org/10.1089/neu.2010.1430>
114. Lovell MR, Pardini JE, Welling J, Collins MW, Bakal J, Lazar N, Roush R, Eddy WF, Becker JT. Functional brain abnormalities are related to clinical recovery and time to return-to-play in athletes. *Neurosurgery* 2007;61(2):352-60.
115. Wu Y-C, Harezlak J, Elsaid NMH, Lin Z, Wen Q, Mustafi SM, Rikken LD, Koch KM, Nencka AS, Meier TB, Mayer AR, Wang Y, Giza CC, DiFiori JP, Guskiewicz KM, Mihalik JP, LaConte SM, Duma SM, Broglio SP, Saykin AJ, McCrea MA, McAllister TW. Longitudinal white-matter abnormalities in sports-related concussion: A diffusion MRI study. *Neurology* 2020;95(7):e781-e92. doi: <https://dx.doi.org/10.1212/WNL.00000000000009930>
116. Sharma A, Hind K, Hume P, Singh J, Neary JP. Neurovascular Coupling by Functional Near Infra-Red Spectroscopy and Sport-Related Concussion in Retired Rugby Players: The UK Rugby Health Project. *Frontiers in human neuroscience* 2020;14(101477954):42. doi: <https://dx.doi.org/10.3389/fnhum.2020.00042>

117. Abrahams S, Mc Fie S, Patricios J, Suter J, September AV, Posthumus M. Toxic tau: The TAU gene polymorphisms associate with concussion history in rugby union players. *Journal of science and medicine in sport* 2019;22(1):22-28. doi: <https://dx.doi.org/10.1016/j.jsams.2018.06.012>
118. Abrahams S, McFie S, Lacerda M, Patricios J, Suter J, September AV, Posthumus M. Unravelling the interaction between the DRD2 and DRD4 genes, personality traits and concussion risk. *BMJ open sport & exercise medicine* 2019;5(1):e000465. doi: <https://dx.doi.org/10.1136/bmjsem-2018-000465>
119. Shahim P, Politis A, van der Merwe A, Moore B, Chou Y-Y, Pham DL, Butman JA, Diaz-Arrastia R, Gill JM, Brody DL, Zetterberg H, Blennow K, Chan L. Neurofilament light as a biomarker in traumatic brain injury. *Neurology* 2020;95(6):e610-e22. doi: <https://dx.doi.org/10.1212/WNL.0000000000009983>
120. Zemek R, Osmond MH, Barrowman N, Pediatric Emergency Research Canada Concussion T. Predicting and preventing postconcussive problems in paediatrics (5P) study: protocol for a prospective multicentre clinical prediction rule derivation study in children with concussion. *BMJ open* 2013;3(8) doi: <https://dx.doi.org/10.1136/bmjopen-2013-003550>
121. Schneider KJ, Critchley M, Anderson V, al e. Targeted interventions and their effect on recovery in children, adolescents and adults who have suffered a sport-related concussion: a systematic review. *British Journal of Sports Medicine* 2023 doi: 10.1136/bjsports-2022-106685
122. Yeates KO, Räisänen AM, Premji Z, al E. What tests and measures accurately diagnose persisting post-concussive symptoms in children, adolescents and adults following sport-related concussion? A systematic review. *British Journal of Sports Medicine* 2023 doi: 10.1136/bjsports-2022-106657
123. Lassman ME, Rathwell S, Black AM, Caron JG. Exploring student-athletes' perceptions of their psychological readiness to return to sport following a concussion. *Sport, Exercise, and Performance Psychology* 2022;11:444-58. doi: 10.1037/spy0000309
124. van Ierssel J, Pennock KF, Sampson M, Zemek R, Caron JG. Which psychosocial factors are associated with return to sport following concussion? A systematic review. *Journal of Sport and Health Science* 2022;11(4):438-49. doi: <https://doi.org/10.1016/j.jshs.2022.01.001>

125. Saadi A, Himmelstein DU, Woolhandler S, Mejia NI. Racial disparities in neurologic health care access and utilization in the United States. *Neurology* 2017;88(24):2268-75. doi: 10.1212/wnl.0000000000004025 [published Online First: 2017/05/19]
126. Iverson G. Possible Long-Term Effects of Participation in Contact and Collision Sports 1 on Brain Health: A Systematic Review. *British Journal of Sports Medicine* 2023
127. Lee IM, Shiroma EJ, Lobelo F, Puska P, Blair SN, Katzmarzyk PT. Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *The Lancet* 2012;380(9838):219-29. doi: 10.1016/S0140-6736(12)61031-9
128. Lobelo F, Rohm Young D, Sallis R, Garber MD, Billinger SA, Duperly J, Hutber A, Pate RR, Thomas RJ, Widlansky ME, McConnell MV, Joy EA. Routine Assessment and Promotion of Physical Activity in Healthcare Settings: A Scientific Statement From the American Heart Association. *Circulation* 2018;137(18):e495-e522. doi: 10.1161/cir.0000000000000559 [published Online First: 2018/04/06]
129. Thompson WR, Sallis R, Joy E, Jaworski CA, Stuhr RM, Trilk JL. Exercise Is Medicine. *Am J Lifestyle Med* 2020;14(5):511-23. doi: 10.1177/1559827620912192 [published Online First: 2020/09/15]
130. Turner M, Maddocks D, Hassan M, Anderson A, McCrory P. Consent, capacity and compliance in concussion management: cave ergo medicus (let the doctor beware). *British Journal of Sports Medicine* 2021;55(10):539-44. doi: 10.1136/bjsports-2020-102108
131. Chang C, Putukian M, Aerni G, Diamond A, Hong G, Ingram Y, Reardon CL, Wolanin A. Mental health issues and psychological factors in athletes: detection, management, effect on performance and prevention: American Medical Society for Sports Medicine Position Statement—Executive Summary. *British Journal of Sports Medicine* 2020;54(4):216-20. doi: 10.1136/bjsports-2019-101583
132. Stambulova NB, Ryba TV, Henriksen K. Career development and transitions of athletes: the International Society of Sport Psychology Position Stand Revisited. *International Journal of Sport and Exercise Psychology* 2021;19(4):524-50. doi: 10.1080/1612197X.2020.1737836

133. Eliason P, Galarneau J, Emery CA. Prevention strategies and modifiable risk factors for sport-related concussions and head impacts: A systematic review and meta-analysis. *British Journal of Sports Medicine* 2023