

1 **Trends in Socioeconomic Inequalities in incidence of Severe Mental Illness – A**
2 **Population-Based Linkage Study Using Primary and Secondary Care Routinely**
3 **Collected Data between 2000 and 2017**

4 Sze Chim Lee^{a,b}, Marcos DelPozo-Banos^{a,b}, Keith Lloyd^{a,b}, Ian Jones^{b,c}, James T.R.
5 Walters^{b,c}, Ann John^{a,b*}

6 ^a HDRUK, Swansea University Medical School, Singleton Park, Swansea, UK, SA2
7 8PP.

8 ^b National Centre for Mental Health. Cardiff University, Hadyn Ellis Building, Maindy
9 Road, Cardiff, UK, CF24 4HQ.

10 ^c MRC Centre for Neuropsychiatric Genetics and Genomics. School of Medicine,
11 Cardiff University, Hadyn Ellis Building, Maindy Road, Cathays, Cardiff, UK, CF24
12 4HQ.

13 *Corresponding author

14 Professor Ann John, Professor of Public Health and Psychiatry, Swansea University Medical
15 School, 3/F Data Science Building, Swansea University, Swansea, UK, SA2 8PP

16 E-mail: a.john@swansea.ac.uk

17 Telephone: +44 (0)1792 602568

18 **Abstract**

19 **Objective:** In 2008, the UK entered a period of economic recession followed by sustained
20 austerity measures. We investigate changes in inequalities by area deprivation and urbanicity
21 in incidence of severe mental illness (SMI, including schizophrenia-related disorders and
22 bipolar disorder) between 2000 and 2017.

23 **Methods:** We analysed 4.4 million individuals from primary and secondary care routinely
24 collected datasets (2000-2017) in Wales and estimated the incidence of SMI by deprivation
25 and urbanicity measured by the Welsh Index of Multiple Deprivation (WIMD) and urban/rural
26 indicator respectively. Using linear modelling and joinpoint regression approaches, we
27 examined time trends of the incidence and incidence rate ratios (IRR) of SMI by the WIMD
28 and urban/rural indicator adjusted for available confounders.

29 **Results:** We observed a turning point of time trends of incidence of SMI at 2008/2009 where
30 slope changes of time trends were significantly increasing. IRRs by deprivation/urbanicity
31 remained stable or significantly decreased over the study period except for those with bipolar
32 disorder sourced from secondary care settings, with increasing trend of IRRs (increase in IRR
33 by deprivation after 2010: 1.6% per year, 95% CI: 1.0%-2.2%; increase in IRR by urbanicity
34 1.0% per year, 95% CI: 0.6%-1.3%).

35 **Conclusions:** There was an association between recession/austerity and an increase in the
36 incidence of SMI over time. There were variations in the effects of deprivation/urbanicity on
37 incidence of SMI associated with short- and long-term socioeconomic change. These findings
38 may support targeted interventions and social protection systems to reduce incidence of SMI.

39 **Keyword**

40 austerity, deprivation, inequality, recession, severe mental illness, urbanicity

41 **Abbreviations**

42 APC –Annual Percentage Change

43 CI – Confidence Interval

44 CPRD – Clinical Practice Research Datalink

45 GPD – General Practice Database

46 ICD – International Classification of Diseases

47 IGRP – Information Governance Review Panel

48 IRR – Incidence Rate Ratio

49 LSOA – Lower-Layer Super-Output Area

50 ONS – Office for National Statistics

51 PEDW – Patient Episode Database for Wales

52 pyar – person-years at risk

53 SAIL – Secure Anonymised Information Linkage

54 SMI – Severe Mental Illness

55 WIMD – Welsh Index of Multiple Deprivation

56 **1. Introduction**

57 The complex aetiology of severe mental illness (SMI), including schizophrenia-related
58 disorders and bipolar disorder has been well documented (Radua et al., 2018). Although the
59 direction of causality remains a source of debate (Collip et al., 2008; Dunham, 1965; Goldberg
60 and Morrison, 1963; Hudson, 2012, 2005; Sariaslan et al., 2016; Selten et al., 2013; Van Os
61 et al., 2008), SMI is more prevalent in urban and socially deprived areas (Allardyce and
62 Boydell, 2006; Faris and Dunham, 1939; Heinz et al., 2013; Kaymaz et al., 2006; Lee et al.,
63 2020; March et al., 2008). This socioeconomic inequality, however, may vary due to person-
64 level factors (age and sex), different pace of urbanisation, population growth between areas
65 and socioeconomic changes (Haukka et al., 2001; Marcelis et al., 1998; Pedersen, 2006).

66 The global financial crisis of 2007/8 triggered an economic recession in the UK characterised
67 by a sharp rise in unmanageable household debt, house repossessions, falls in financial
68 markets, rises in unemployment and under-employment, and falling wages (Carrera and
69 Beaumont, 2011; Coope et al., 2014; Gamble, 2009; ONS, 2018). Following the recession,
70 the UK government implemented a stringent policy of austerity where public spending was
71 pervasively cut to reduce expenditure and restore economic growth (Crawford and Phillips,
72 2012; Karanikolos et al., 2013; Kitson et al., 2011; Reeves et al., 2013). These measures
73 potentially reduced funding to healthcare and social welfare services in real terms and
74 introduced reforms in social security and welfare benefits (Akhter et al., 2018). The Welfare
75 Reform Act 2012 (The UK government, 2012) was enacted to cap and restrict social security
76 and housing benefits (e.g., 'bedroom tax'). Besides, a controversial reassessment on the
77 eligibility of approximately 1.5 million existing claimants was performed via the Work Capability
78 Assessment (The UK government, 2013) and those who were reassessed fit for work were
79 moved from the benefit (Barr et al., 2016; Harrington, 2010; Litchfield, 2014).

80 Substantial research has suggested that social and geographical inequalities were widened
81 in already vulnerable individuals during the period of the 2007/8 recession and austerity
82 measures, particularly those from disadvantaged social groups, were impacted
83 disproportionately (Akhter et al., 2018; Beatty and Fothergill, 2014; Browne and Levell, 2010;
84 Crawford and Phillips, 2012; De Agostini et al., 2014; Taylor-Robinson et al., 2014). There
85 have been quantitative studies evaluating the effects of the recession and austerity on social
86 inequalities in mental health (Akhter et al., 2018; Barr et al., 2015, 2016; Katikireddi et al.,
87 2012). However, few targeted those with SMIs specifically, nor were they free from
88 methodological limitations, including non-representative/small samples, reliance on self-
89 reported surveys often with un-validated measures on mental health, and limited data
90 coverage after the onset of recession through to austerity (from 2008, up to 2015). Using the

91 UK clinical practice research datalink (CPRD) gathered between 2003 and 2013 from primary
92 care settings, Kendrick and colleagues (Kendrick et al., 2015) reported a significant increase
93 in prevalence of depression only for the most socioeconomically deprived quintile of practices
94 after the second quarter of 2008. However, this widening gap following recession was not
95 replicated in another study comparing prevalence of SMIs from 2000 to 2012 by area
96 deprivation using the same CPRD dataset (Reilly et al., 2015). In this study, we explored the
97 time trends of socioeconomic inequalities in incidence of SMIs, including the period of
98 recession and austerity, using both population-based primary and secondary care routinely
99 collected data in Wales, UK. We estimated both incidence and change in incidence of SMIs
100 by level of area deprivation and urbanicity between 01/01/2000 and 31/12/2017.

101 **2. Materials and methods**

102 2.1 Study design

103 This is a retrospective population-based observational electronic cohort study.

104 2.2 Study population and setting

105 The eligible study population was all individuals (~4.4 million) aged 15 years or above and
106 who continuously lived in Wales, UK for one or more years between 01/01/2000 and
107 31/12/2017 (Suppl. Figure 1A). Inclusion and exclusion criteria for calculation of annual
108 incidences are described below (section 2.5.3).

109 2.3 Data Source

110 We interrogated the Secure Anonymised Information Linkage (SAIL) databank
111 (www.saildatabank.com), a databank that contains anonymised privacy protecting person-
112 based linkable data from healthcare and public settings (Ford et al., 2009; Lyons et al., 2009).
113 In this study, we used data extracted from the General Practice Database (GPD) that includes
114 contacts to general practices, the Patient Episode Database for Wales (PEDW) that contains
115 inpatient hospital admissions, Welsh Demographic Service and the Office for National
116 Statistics (ONS) deaths register datasets. Description of each dataset is outlined in the Suppl.
117 Methods.

118 2.4 Ethical approval

119 Ethical approval was granted by the Information Governance Review Panel (IGRP) with the
120 approval number 0466. The IGRP is an independent body including representatives with a
121 variety of expertise from different organisations to oversee study approvals in line with

122 permissions already granted to the analysis of data in the SAIL databank (Ford et al., 2009;
123 Lyons et al., 2009).

124 2.5 Measures

125 2.5.1 Outcomes (SMI diagnoses)

126 We classified SMI diagnoses into schizophrenia-related disorders (schizophrenia, schizotypal,
127 delusional and schizoaffective disorders) and bipolar disorder as described in our previous
128 study (Lee et al., 2020). We used Read Code version 2 (5-byte) for the GPD (primary care
129 cohort) and the International Classification of Diseases (ICD) version 10 for the PEDW
130 (secondary care cohort) to identify both disorders. The use and validation of diagnostic codes
131 for SMI have been described elsewhere (Economou et al., 2012; Ford et al., 2009; John et al.,
132 2018; Lee et al., 2020; Lloyd et al., 2015), with codes being cross-mapped between Read
133 and ICD-10 classifications by the NHS used in our recent study (Lee et al., 2020).

134 2.5.2 Exposure (Area deprivation and urbanicity)

135 We adopted the previously used measures for area deprivation and urbanicity (Lee et al.,
136 2020). For area deprivation, the Welsh Index of Multiple Deprivation (WIMD) 2011 was used
137 (Welsh Government, 2017) throughout the study period (01/01/2000-31/12/2017). The WIMD
138 measures deprivation for each Lower-layer super-output area (LSOA), geographic unit used
139 in reporting small area statistics comprised of approximately 1,500 individuals, by a weighted
140 score that combines eight domains of deprivation. These domains were income, housing,
141 employment, geographical access to services, education, health, community safety and
142 physical environment. We used WIMD quintiles where the WIMD scores were ranked and
143 grouped from the least (Q1) to most (Q5) deprived areas. For urbanicity, we used the
144 urban/rural indicator for England and Wales to categorise LSOAs as either urban or rural
145 (Barham and Begum, 2006). Rural areas contained town and fringe, villages, hamlets and
146 isolated dwellings while urban areas contained all urban settlement types with a population of
147 10,000 or more. To capture moving of individuals, both WIMD quintile and urban/rural indicator
148 were extracted at the beginning of each year (1st January) until the date of the respective first
149 SMI diagnosis in the respective cohorts. However, these exposures were time-fixed with
150 respect to LSOAs (e.g., an urban LSOA remained urban throughout the study period).

151 2.5.3 Incidence of SMIs

152 We calculated first recorded incidence of SMIs between 01/01/2000 and 31/12/2017.
153 Incidence was defined as the number of first diagnoses (with no previous recorded
154 schizophrenia-related or bipolar disorders) over the whole 18-year period divided by the

155 number of person years at risk (pyar) within each WIMD quintile or urban/rural group.
156 Incidence rates by calendar year were calculated by grouping incident cases and pyar for each
157 year. Individuals and time at risk were included whenever they resided in Wales and were 15
158 years or above. Person-time were excluded whenever they moved out Wales and they were
159 excluded (permanently) after the date of death (if present within the study period). Individuals
160 and time at risk were not included in the incidence calculations if information of LSOA was not
161 available on 1st January each year or at the date of incident diagnosis of SMIs.

162 Incidence was calculated separately for the primary and secondary care cohorts because
163 these two cohorts did not share the same denominators for their differences in data coverage.
164 However, individuals with both schizophrenia-related disorders and bipolar disorder were
165 considered in the calculation of incidence of each of the conditions (i.e., the incident population
166 of the two conditions were not mutually exclusive). For the primary care cohort, we adopted
167 an algorithm (Davies et al., 2018) to identify periods of valid GP data coverage within the study
168 period to avoid biased estimation of incidence due to non-complete data coverage. For the
169 denominator of incidence within the GP population, the pyar of an individual was the
170 summation of all valid periods bounded by the GP registration start and end dates within the
171 study period. For the secondary care cohort, the whole population in Wales and the
172 corresponding pyar within the study period contributed to the incidence calculation. Incidences
173 (and pyars) were stratified by year throughout the study period.

174 2.6 Analysis and statistical methods

175 We extracted linked data in SAIL via structured query language (SQL DB2). We summarised
176 sociodemographic characteristics of the study population as descriptive statistics with 95%
177 confidence intervals (CIs). We adopted two-tailed mid-*p* exact CIs (assuming Poisson
178 distribution) for incidence rates (Rothman and Boice, 1979), Wilson score with continuity
179 correction for estimating CIs for proportions (Newcombe, 1998). Incidence were expressed as
180 number of individuals per 100,000 pyar. Unless otherwise stated, all statistical analyses were
181 performed using Stata version 16.1 (StataCorp, 2019) and the level of statistical significance
182 was set at $p = 0.05$.

183 2.6.1 Gradient of incidence by deprivation and urbanicity

184 We estimated gradients of incidence by deprivation and urbanicity across time using Poisson
185 regression. Four regressions were run separately for deprivation and urbanicity, as well as for
186 data sourced from primary and secondary care cohort. Details of model specifications are
187 described in the Suppl. Methods. The parameter of interest was the incidence rate ratio (IRR)
188 by WIMD quintile and urban/rural indicator by year, which quantifies the direction and

189 magnitude of socioeconomic inequality at a particular year. An IRR greater than one indicated
190 higher incidence in more deprived/urban areas. Widening of inequalities over the study period
191 would be evident if an increase in IRRs over time was observed.

192 Stratified analyses were conducted for sex and age groups (15-24, 25-34, 35-44, 45-54; 55-
193 64, 65-74, and ≥ 75 years). Due to sample size issues, stratification by age could not be
194 performed. We alternatively restricted individuals aged between 25 and 64 years only as a
195 sensitivity analysis (see below).

196 2.6.2 Trends of gradient of incidence by deprivation and urbanicity

197 We performed joinpoint regressions (Kim et al., 2000; National Cancer Institute, 2015; Wagner
198 et al., 2002) using the Joinpoint software (Version 4.7.0.0) on the incidences rates and by
199 deprivation and urbanicity by year to identify years where change in trends is evident (see
200 Suppl. Methods for details). For IRRs, increasing (decreasing) trends indicated widening
201 (narrowing) of social inequalities over the respective period. We reported annual percentage
202 change (APC), i.e., exponentiating the slope (β coefficient) in incidence rates/IRRs within a
203 line segment, years where the break points occurred, change in slope (change in β) at break
204 points with corresponding 95% CIs.

205 2.6.3 Sensitivity Analysis

206 We carried out three sensitivity analyses to examine robustness of our results. The first one
207 was to test the sensitivity of age inclusion criteria. The second and the third addressed the
208 potential effects of temporal change in health service provision on our trends. The last two
209 sensitivity analyses utilised the annual counts of available diagnoses to measure overall
210 service provision. Detailed analytical strategies for the sensitivity analyses are outlined in
211 Suppl. Methods.

212 3. Results

213 3.1 Study population

214 We identified 3,771,811 eligible individuals (68.9%) who were 15 years or older and
215 continuously registered in Wales between 01/01/2000 and 31/12/2017 (Suppl. Figure 1A).
216 There were 3,054,737 individuals (81.0% of 3,771,811) in the primary care cohort.
217 Demographic characteristics, including population distributions over deprivation/urbanicity
218 between primary and secondary care cohorts were similar (Suppl. Table 1A). Among the
219 35,394 individuals identified with schizophrenia-related disorders or bipolar disorder (Suppl.
220 Figure 1A), 20,558 individuals (58.1%) identified with schizophrenia-related disorders only,

221 12,154 (34.3%) with bipolar disorder only and 2,682 (7.6%) with both disorders (Suppl. Figure
222 1B). Respectively 29.8% (10,563 out of 35,394) and 44.9% (15,894) of the individuals with
223 SMIs were sourced from primary and secondary care cohorts only, and the remaining 25.3%
224 (8,397) were from both cohorts (Suppl. Figure 1C). For the primary care cohort, 12,907 and
225 7,400 individuals were identified having schizophrenia-related disorders and bipolar disorder
226 respectively whereas 15,469 for schizophrenia-related disorders and 11,028 for bipolar
227 disorder individuals were identified from the secondary care cohort (Suppl. Figure 1A). During
228 the 18-year study period, incidence of schizophrenia-related disorders were 39.2 and 33.8 per
229 100,000 pyar from primary and secondary care cohort respectively whereas for bipolar
230 disorder the corresponding incidences were 22.5 and 24.0 per 100,000 pyar respectively
231 (Suppl. Table 2).

232 3.2 Time trends of incidence of SMIs

233 Incidence of schizophrenia-related disorders and bipolar disorder by year from primary and
234 secondary care cohort and the respective joinpoint regression estimates are summarised in
235 Figure 1. In general, incidences of schizophrenia-related disorders (26.4-61.1 per 100,000
236 pyar) were higher than Bipolar disorder (15.6-36.3 per 100,000 pyar). Before 2004, we found
237 incidences were increasing only in primary care cohort (Figure 1A and C). Incidences of
238 schizophrenia-related disorders were significantly decreasing between 2004 and 2009 but
239 then significantly increased over 2009 and 2014, with a significant positive change of slope at
240 2009 for both cohorts (change in $\beta = 0.215$, 95% CI: 0.154-0.265, $p < 0.001$ for primary and
241 change in $\beta = 0.120$, 95% CI: 0.061-0.178, $p = 0.005$ for secondary). While for the primary
242 care cohort incidence of schizophrenia-related disorders decreased again from 2014, the
243 increase started from 2009 continued to the end of study period for the secondary care cohort.
244 For bipolar disorder, incidence trend sourced from the primary care cohort significantly
245 changed from decreasing between 2003 and 2008 to increasing between 2008 and 2014
246 (change in $\beta = 0.096$, 95% CI: 0.046-0.146, $p = 0.007$). A similar break point with a significant
247 increase in slope at 2008 was found for the secondary care cohort (change in $\beta = 0.062$, 95%
248 CI: 0.015-0.110, $p = 0.037$). From 2014 onwards, incidence of bipolar disorder sourced from
249 the primary but not the secondary care cohort resumed decreasing.

250 3.3 Time trends of gradients of incidence of SMIs by deprivation and urbanicity

251 Incidence of schizophrenia-related disorders and bipolar disorder stratified by WIMD
252 quintile/urbanicity per calendar year from primary and secondary care cohort are depicted in
253 panel A, B, E and F of Figure 2-3 respectively (see also Suppl. Table 3-4). In general, we
254 observed associations of higher incidence of SMIs in more deprived and urban areas. For
255 bipolar disorder, these gaps were still evident but smaller.

256 Annual IRRs by deprivation/urbanicity also reflected the presence of inequalities (panel C, D,
257 G and H of Figure 2-3). IRRs were significantly > 1 for all years for schizophrenia-related
258 disorders (IRRs from 1.2 to 1.4) in the deprivation model (Suppl. Table 5). For bipolar disorder,
259 the corresponding IRRs ranged from 1.0 to 1.3 over the study period. In the urbanicity models
260 (Suppl. Table 6), statistical significance of IRRs > 1 was much less prevalent compared to
261 deprivation models (IRRs from 0.9 to 1.5 and 0.8-1.4 for schizophrenia-related disorders and
262 bipolar disorder respectively) due to their wider CIs.

263 Results from the joinpoint regression modelling (Table 1, Figure 2-3) for the IRRs shows that
264 all but two time trends of IRRs by deprivation and urbanicity within the study period consisted
265 of only one linear segment. One exception was the IRRs by deprivation for schizophrenia-
266 related disorders sourced from the primary care cohort, consisting of three linear segments
267 with joinpoints at 2004 and 2014 (Figure 2C). The other exception was the IRR by deprivation
268 for bipolar disorder sourced from the secondary care cohort; containing two linear segments
269 with a joinpoint at 2010 (Figure 3H). In the secondary care cohort, we found significant
270 increasing trends of IRRs by deprivation after 2010 (APC = 1.6%, 95% CI: 1.0% to 2.2%, $p <$
271 0.001) and by urbanicity throughout the whole study period (APC = 1.0%, 95% CI: 0.6% to
272 1.3%, $p < 0.001$) for bipolar disorder. IRR by deprivation for schizophrenia-related disorders
273 sourced from the primary care cohort increased from 2000 to 2004 and then decreased until
274 2014. Starting from 2014, the increasing trend resumed (APC = 2.6%, 95% CI: 0.7% to 4.5%,
275 $p < 0.022$). All other IRRs decreased or remained stable with slopes not significantly different
276 from zero.

277 3.4 Sex-specific time trends of gradients of incidence of SMIs by deprivation and urbanicity

278 Annual IRRs by deprivation and urbanicity stratified by sex (Suppl. Figure 2-3 and Suppl. Table
279 7-11) showed slightly larger IRRs by deprivation in males for schizophrenia-related disorders
280 (average IRR for males vs. females: 1.3 vs. 1.2). Time trends of IRRs were not identical
281 between sexes but consistent with the overall trends (Figure 2-3). For the primary care cohort,
282 the increasing IRRs by deprivation after 2014 (Suppl. Figure 2A) and the overall decreasing
283 IRRs by urbanicity (Suppl. Figure 2B) for schizophrenia-related disorders were contributed by
284 females. The monotonic decrease in IRRs by urbanicity for bipolar disorder consisted of a
285 decreasing trend from males before 2007 and another from females after 2013 (Suppl. Figure
286 2D). For the secondary care cohort, trends in IRRs by deprivation for schizophrenia-related
287 disorders was significantly increasing for males but not females (Suppl. Figure 3B) while the
288 increasing trends of IRRs for bipolar disorder was mainly observed in females (Suppl. Figure
289 3D).

290 3.5 Sensitivity analysis

291 Table 2 schematically summarises the post-2008 trends in incidences, proportions, gradients
292 of incidences and proportions by deprivation and urbanicity from the main analysis and the
293 three sensitivity analyses. The first sensitivity analysis by including individuals aged between
294 25 and 64 years only revealed similar results to the main analysis (Suppl. Figure 4 and Suppl.
295 Table 12-14). The significant increasing trend of gradient of incidence by deprivation for bipolar
296 disorder sourced from secondary care cohort were still robust after 2008. The second
297 sensitivity analysis (including an additional term of overall available diagnosis to the models)
298 yielded nearly identical results with the main analysis (Suppl. Figure 5 and Suppl. Table 15-
299 17). Our third sensitivity analysis using proportion outcomes generally tallied with the main
300 analysis (Suppl. Figure 6-8 and Suppl. Table 18-20). Proportion trends (Suppl. Figure 6) were
301 similar to the incidence trends from the main analysis (Figure 1) with the exception in
302 schizophrenia-related disorders. Besides, annual relative risks (RRs) by urbanicity appeared
303 slightly higher than the IRRs from the main analysis, particularly for schizophrenia-related
304 disorders. Comparisons of trends of annual RRs and IRRs by deprivation/urbanicity showed
305 robust the increasing trends of gradients by deprivation for bipolar disorders sourced
306 secondary care cohort (Figure 3E and 3G and Suppl. Figure 8E and 8G).

307 **4. Discussion**

308 4.1 Main findings

309 We found changes in time trends of incidence of SMIs, including schizophrenia-related
310 disorders and bipolar disorder, and associated variations in socioeconomic inequalities in
311 terms of area deprivation and urbanicity from 2000 to 2017. Our study period covered the
312 2007/8 economic recession and the subsequent implementation of austerity measures by the
313 UK government. Regardless of the cohorts and conditions we examined, we found the period
314 2008/2009 was a time associated with significant increases in the slopes of time trends of
315 incidences of SMIs. These changes in slopes occurred roughly at the time of the recession.
316 For trends of socioeconomic inequalities of incidence of SMIs (quantified by IRRs), although
317 we did not observe significant widening of inequalities at the time of recession/austerity,
318 inequality was significantly increasing for those with bipolar disorder for data sourced from the
319 secondary care cohort (apparently irrespective of the recession/austerity), which has not been
320 previously reported in literature to our knowledge. Results from our sensitivity analyses were
321 generally consistent with the main analysis with few exceptions. Our analyses were able to
322 mutually adjust between area deprivation and urbanicity, to capture the sex-specific effects on
323 socioeconomic inequalities in incidence of SMIs. To our knowledge, these analyses are not
324 commonly reported in the available literature.

325 Our overall incidence rates of SMIs are comparable to other studies (Hardoon et al., 2013;
326 Jongasma et al., 2019, 2018; Kirkbride et al., 2012; Lee et al., 2020; Saha et al., 2005; Simeone
327 et al., 2015). Differences in demographic characteristics between primary and secondary care
328 cohort were minimal and outcomes between cohorts were highly comparable. Our time trend
329 of incidence of SMIs and proportion of SMIs diagnoses to all available diagnoses were
330 generally decreasing or did not change before 2008/2009 and then began increasing. This
331 pattern was previously demonstrated in mental health outcomes such as depression
332 (Katikireddi et al., 2012; Kendrick et al., 2015), population mental health (Barr et al., 2015),
333 and suicide (Coope et al., 2014), indicating the robust impact of recession on mental health.

334 Possible reasons for the increases in incidences in SMIs include chronic stressors such as
335 financial adversity associated with the economic recession on mental health, including debt,
336 reduction of income, house repossession, and (anticipated) unemployment/job insecurity at
337 personal/household level (Coope et al., 2014). Such adversities during economic downturn
338 may also place strains on relationships further deteriorating individuals' mental health. It has
339 also been proposed that changes in public spending or reallocation of resources particularly
340 on healthcare and social welfare services during times of economic uncertainty may curtail
341 the usual social and financial support for prevention, personal resources to help seek and
342 access to treatments for mental health problems to the individuals at-risk and the community
343 (Akhter et al., 2018; Barr et al., 2015, 2016; Katikireddi et al., 2012). Additionally greater and
344 widening inequalities within societies are associated with increased levels of mental disorders
345 (Wilkinson and Pickett, 2009). It is possible that causal relationships in the development and
346 expression of SMIs are mediated by complex gene-environment (socio-economic)
347 interactions.

348 We believe the increase in incidences from 2000 and 2004 in the primary care cohort only are
349 associated with the Quality and Outcomes Framework (QOF) introduced in 2004, aiming to
350 improve quality of primary care provision by providing financial incentive for family doctors in
351 managing and documenting certain health indicators/outcomes (Fichera et al., 2016). Brief
352 increases of diagnoses on physical and mental health were reported elsewhere reflecting
353 improved case finding associated with QOF (Fichera et al., 2016; John et al., 2016; McLintock
354 et al., 2014; O'Donoghue, 2009).

355 We also replicated robust associations between incidence of SMIs and deprivation/urbanicity
356 as demonstrated by numerous research studies (Hardoon et al., 2013; Kaymaz et al., 2006;
357 Lee et al., 2020; O'Donoghue et al., 2016; Radua et al., 2018; Vassos et al., 2012). Our IRRs
358 of schizophrenia-related disorders between the most and least deprived and between urban
359 and rural areas are consistent with others (Hardoon et al., 2013; Kelly et al., 2010; Lee et al.,

360 2020; O'Donoghue et al., 2016). In keeping with others, we reported smaller socioeconomic
361 inequalities for bipolar disorder compared with schizophrenia-related disorders (Gruebner et
362 al., 2017; Hardoon et al., 2013; Heinz et al., 2013; Kaymaz et al., 2006; Laursen et al., 2007;
363 Lee et al., 2020; March et al., 2008).

364 We discovered that changes in socioeconomic inequalities in incidence of SMIs over time
365 might not be consistent between primary and secondary (individuals with higher severity or
366 less social support) care cohorts. For schizophrenia-related disorders, socioeconomic
367 inequalities significantly decreased over the majority of the study period from the primary but
368 remained stable in the secondary care cohort. Interestingly, we observed signs of increasing
369 inequality by deprivation for the primary care cohort from 2014. For bipolar disorder, we found
370 significant widening of inequality from the secondary but not in primary care cohort. The
371 increasing gap in incidence of bipolar disorder by deprivation found from hospitalisation data
372 over the 18-year period is novel, alarming and requires further investigation. We also argue
373 for further research to disentangle the relationships between severity of illness, contacts with
374 types of health services and the socioeconomic inequalities of SMIs. Apart from cohort
375 settings, we observed discordance of time trends of inequality between deprivation and
376 urbanicity. This could be explained by the joint adjustments between deprivation and urbanicity
377 in our models to reduce confounding between level of deprivation and urbanicity as previously
378 reported (Lee et al., 2020).

379 Overall, we did not identify significant increase in socioeconomic gap of incidence of SMIs
380 associated with recession and austerity. Quantitative evidence supporting widening
381 socioeconomic inequalities in population mental health following recession/austerity was
382 inconclusive using self-reported population survey data (Akhter et al., 2018; Barr et al., 2016,
383 2015; Coope et al., 2014; Katikireddi et al., 2012). Results were also mixed from population-
384 based routinely collected data. From a UK analysis on primary care data using the CPRD,
385 prevalence of depression was found statistically increased after the second quarter of 2008
386 for GPs located at the most deprived quintile only (Kendrick et al., 2015). In a recent study
387 using the same data source, Reilly and others (Reilly et al., 2015) compared prevalences of
388 SMIs including schizophrenia and bipolar disorders across four nations in the UK and
389 deprivation from 2000 to 2012. For the UK data, they found that the gap by deprivation in SMIs
390 did not increase corresponding to the recession (Reilly et al., 2015).

391 Our findings of relatively stable/decreasing trends in urban-rural inequality of schizophrenia-
392 related disorders over the 18-year study period is not consistent with earlier European studies
393 showing long-term increase in the effects of urban birth to schizophrenia/psychoses
394 throughout the 20th century (Haukka et al., 2001; Marcelis et al., 1998; Pedersen, 2006). In

395 contrast, a more recent study from the ONS using the same urban/rural classification as in
396 this study showed that population growth in rural- and urban-Wales were very similar between
397 2001 and 2009 (Pateman, 2011). We believe that this similar expansion of urban and rural
398 areas in Wales implies a stable urbanisation phase in recent years and could further explain
399 our relatively stable trends compared with earlier studies where rapid urbanisation was still
400 underway (Haukka et al., 2001; Marcelis et al., 1998; Pedersen, 2006).

401 This study also identify differences in socioeconomic inequalities between sexes. Our
402 comparison of gradients of incidence between males and females did not indicate wider
403 inequalities for either sex in most studied conditions, except that wider inequalities existed in
404 males for schizophrenia-related disorders. This tallies with the findings that the effect of urban
405 birth on the elevated risk of schizophrenia and other psychoses is stronger in males (Haukka
406 et al., 2001; Kelly et al., 2010; Marcelis et al., 1998; Pedersen, 2006).

407 4.2 Strength and limitations

408 This study revealed changes in inequalities in incidence of schizophrenia-related disorders
409 and bipolar disorder by deprivation/urbanicity over time for data sourced from primary and
410 secondary care settings. Besides the main analysis with sex-specific trends of socioeconomic
411 inequalities, we replicated our findings by using different age inclusion criteria, controlling for
412 overall use of health services and using alternative outcome (proportion of diagnoses). Such
413 comprehensive analyses are not commonly performed using population-based datasets.
414 Similar to our previous study (Lee et al., 2020), our complete data coverage of the Wales
415 population in the hospital admission dataset and >70% coverage in the GP dataset is
416 advantageous in terms of data coverage and representation. To circumvent the issue of
417 incomplete data coverage in the primary care cohort, we reported outcomes sourced from
418 both primary and secondary care cohorts to obtain results representative to the general
419 population. We identified individuals with SMIs using diagnostic codes adopted and validated
420 in previous studies to ensure validity and reliability (Economou et al., 2012; John et al., 2018;
421 Lee et al., 2020; Lloyd et al., 2015). We also believe our study period between 2000 and 2017
422 was long enough to capture possible effects of recession/austerity on socioeconomic
423 inequalities compared with other studies (Akhter et al., 2018; Barr et al., 2016, 2015;
424 Katikireddi et al., 2012).

425 One major limitation of our study is the lack of single agreed measures on area deprivation
426 nor urbanicity (Lee et al., 2020). We used WIMD and urban/rural indicator because of their
427 popularity of use (Barham and Begum, 2006; Welsh Government, 2017) and data availability.
428 Nonetheless, our results on the gradients of incidence of SMIs were comparable to others
429 using alternative measures of deprivation and urbanicity. Using WIMD and urban/rural

430 indicator as area variable for individual characteristics may lead to ecological bias (March et
431 al., 2008). We acknowledge that socioeconomic characteristics in an area may change over
432 time and indeed the WIMD is recalculated from time to time. However, comparing the WIMD
433 from different versions does not inform absolute change of deprivation over time (Welsh
434 Government, 2017). We need further efforts to establish absolute level of deprivation and
435 urbanicity.

436 We might underestimate change in socioeconomic inequalities associated with
437 recession/austerity. The widening inequality associated with recession/austerity could be
438 masked by floor effects e.g., individuals already resided in the most deprived areas before
439 recession and further deterioration of socioeconomic circumstance/environment could not be
440 captured (Akhter et al., 2018). Our 18-year study period might not capture longer-term
441 mechanisms that change mental health outcomes in response to change in neighbourhood
442 characteristics, Mechanisms underlying the presence of the inequalities of SMIs could be
443 lifelong or inter-generational processes (Faris and Dunham, 1939; Heinz et al., 2013; March
444 et al., 2008; Norman, 2018; Paksarian et al., 2018; Plomin and Deary, 2015). These processes
445 not only involve socioeconomic factors, but also familial and genetic factors (Colodro-Conde
446 et al., 2018; Paksarian et al., 2018; Sariaslan et al., 2016, 2015) that could remain stable over
447 generations.

448 We used annual total number of available diagnoses as a measure of change in health
449 services provision in two of our sensitivity analyses. However, number of available diagnoses
450 is not a direct measure of health services provision (e.g., number of hospital available). Neither
451 could we assess the effect of policy change in health care provision (e.g., Home Treatment
452 teams) on identifying SMIs using GPD and hospitalisation datasets only. We used annual
453 proportion of SMIs to overall diagnose as alternative outcome. An important caveat of
454 interpreting results from proportion is that it measures socioeconomic inequality of SMIs
455 relative to socioeconomic inequality in health, instead of absolute inequality based on
456 incidences.

457 Other limitations include those shared by other studies using routinely collected data for
458 research. Information bias can occur on underestimating rates and misclassify SMIs for
459 individuals who do not contact to health services or have non-detectable symptoms. Selection
460 bias due to loss of follow-up and missing data may reduce validity. In keeping with our previous
461 study (Lee et al., 2020), we only included individuals with continuous registration in Wales
462 within the follow-up period so that both WIMD and urban/rural indicator were available for
463 analyses. However, this may bias incidence calculation by excluding e.g., the homeless
464 populations and individuals moving in/out Wales intermittently. We analysed first recorded

465 incidence instead of prevalence because of the difficulty in defining remission and relapse of
466 SMIs from routinely collected data (Lee et al., 2020). Given the incidence gradients of SMIs
467 by deprivation/urbanicity are similar as previously shown (Lee et al., 2020), we would obtain
468 similar results if prevalence was used. Other limitations include the presence of unmeasured
469 confounding due to data unavailability (e.g., ethnicity, employment, genetic liability and familial
470 background, status of substance use). Age-specific trends of gradients were not analysed due
471 to sample size issues. Due to sample size issues, we did not perform analyses separately for
472 individuals with both schizophrenia-related disorders and bipolar disorder.

473 4.3 Implications for policy and practice

474 This study makes an integral contribution to policy makers concerning the effects of economic
475 downturn and subsequent policies such as austerity on socioeconomic gap in individuals with
476 SMIs. Our findings not only identify an increase in risk (incidences) of SMIs associated with
477 the economic recession/austerity, but also groups (individuals with bipolar disorder requiring
478 hospitalisation) exhibiting widening of inequalities over the 18-year period. Taking our results
479 together, we highlighted that the impact of recession/austerity on SMIs was similar among
480 socioeconomic levels and these findings were crucial on formulating intervention policies in
481 similar situations. Although socioeconomic gap was not shown as widening in times of
482 recession, our findings do not argue against the potential benefits of providing social protection
483 systems and programmes to fend off risks associated with economic downturn. Recent
484 research suggested that disadvantaged social groups indeed benefit most from increases in
485 spending on social protection following recession (Niedzwiedz et al., 2016). While the debate
486 on how recession/austerity affects inequalities in mental health remains controversial and
487 unsettled, analysis similar to the current study could help identify subtle changes in
488 socioeconomic inequalities in mental illnesses linked to both short- and long-term social and
489 economic changes and help facilitate timely planning of relevant infrastructure.

490 **5. Competing Interest**

491 None declared.

492 **6. Funding**

493 The study was funded by the Welsh Government through Health and Care Research Wales:
494 grant awarded to The National Centre for Mental Health' with Grant No.: CA04, as well as by
495 Health Data Research UK which receives its funding from HDR UK Ltd (NIWA1) funded by
496 the UK Medical Research Council, Engineering and Physical Sciences Research Council,
497 Economic and Social Research Council, Department of Health and Social Care (England),
498 Chief Scientist Office of the Scottish Government Health and Social Care Directorates, Health
499 and Social Care Research and Development Division (Welsh Government), Public Health
500 Agency (Northern Ireland), British Heart Foundation (BHF) and the Wellcome Trust.

501 **7. Contributors**

502 All authors conceived the study; AJ and SCL designed the study; AJ supervised the study;
503 SCL conduct the analysis; SCL and AJ wrote the initial draft and all authors commented on
504 the manuscript.

505 **References**

- 506 Akhter, N., Bamba, C., Mattheys, K., Warren, J., Kasim, A., 2018. Inequalities in mental
507 health and well-being in a time of austerity: Follow-up findings from the Stockton-on-
508 Tees cohort study. *SSM - Popul. Heal.* 6, 75–84.
509 <https://doi.org/10.1016/J.SSMPH.2018.08.004>
- 510 Allardyce, J., Boydell, J., 2006. Environment and schizophrenia: Review: The wider social
511 environment and schizophrenia. *Schizophr. Bull.* 32, 592–598.
512 <https://doi.org/10.1093/schbul/sbl008>
- 513 Barham, C., Begum, N., 2006. The new urban/rural indicator in the Labour Force Survey.
514 *Labour Mark. Trends* 114, 409–418.
- 515 Barr, B., Kinderman, P., Whitehead, M., 2015. Trends in mental health inequalities in
516 England during a period of recession, austerity and welfare reform 2004 to 2013. *Soc.*
517 *Sci. Med.* 147, 324–331. <https://doi.org/10.1016/J.SOCSCIMED.2015.11.009>
- 518 Barr, B., Taylor-Robinson, D., Stuckler, D., Loopstra, R., Reeves, A., Whitehead, M., 2016.
519 ‘First, do no harm’: are disability assessments associated with adverse trends in mental
520 health? A longitudinal ecological study. *J. Epidemiol. Community Health* 70, 339 LP –
521 345. <https://doi.org/10.1136/jech-2015-206209>
- 522 Beatty, C., Fothergill, S., 2014. The local and regional impact of the UK’s welfare reforms.
523 *Cambridge J. Reg. Econ. Soc.* 7, 63–79. <https://doi.org/10.1093/cjres/rst035>
- 524 Browne, J., Levell, P., 2010. The distributional effect of tax and benefit reforms to be
525 introduced between June 2010 and April 2014: a revised assessment. London.
- 526 Carrera, S., Beaumont, J., 2011. Income and wealth. *Soc. Trends* 41, 237–259.
527 <https://doi.org/10.1057/st.2011.10>
- 528 Collip, D., Myin-Germeys, I., Van Os, J., 2008. Does the concept of “sensitization” provide a
529 plausible mechanism for the putative link between the environment and schizophrenia?
530 *Schizophr. Bull.* 34, 220–225. <https://doi.org/10.1093/schbul/sbm163>
- 531 Colodro-Conde, L., Couvy-Duchesne, B., Whitfield, J.B., Streit, F., Gordon, S., Kemper,
532 K.E., Yengo, L., Zheng, Z., Trzaskowski, M., De Zeeuw, E.L., Nivard, M.G., Das, M.,
533 Neale, R.E., MacGregor, S., Olsen, C.M., Whiteman, D.C., Boomsma, D.I., Yang, J.,
534 Rietschel, M., McGrath, J.J., Medland, S.E., Martin, N.G., 2018. Association between
535 population density and genetic risk for schizophrenia. *JAMA Psychiatry* 75, 901–910.
536 <https://doi.org/10.1001/jamapsychiatry.2018.1581>
- 537 Coope, C., Gunnell, D., Hollingworth, W., Hawton, K., Kapur, N., Fearn, V., Wells, C.,
538 Metcalfe, C., 2014. Suicide and the 2008 economic recession: Who is most at risk?
539 Trends in suicide rates in England and Wales 2001–2011. *Soc. Sci. Med.* 117, 76–85.
540 <https://doi.org/10.1016/J.SOCSCIMED.2014.07.024>
- 541 Crawford, R., Phillips, D., 2012. Local government spending: where is the axe falling?
- 542 Davies, G., Jordan, S., Brooks, C.J., Thayer, D., Storey, M., Morgan, G., Allen, S., Garaiova,
543 I., Plummer, S., Gravenor, M., 2018. Long term extension of a randomised controlled
544 trial of probiotics using electronic health records. *Sci. Rep.* 8, 7668.
- 545 De Agostini, P., Hills, J., Sutherland, H., 2014. Were we really all in it together? The
546 distributional effects of the UK Coalition government’s tax-benefit policy changes,
547 CASE - Social Policy in a Cold Climate Working Paper. Centre for Analysis of Social

- 548 Exclusion, LSE. <https://doi.org/DOI:>
- 549 Dunham, H.W., 1965. *Community and schizophrenia: An epidemiological analysis*. Wayne
550 State University Press, Oxford, England.
- 551 Economou, A., Grey, M., McGregor, J., Craddock, N., Lyons, R.A., Owen, M.J., Price, V.,
552 Thomson, S., Walters, J.T.R., Lloyd, K., 2012. The Health Informatics Cohort
553 Enhancement project (HICE): using routinely collected primary care data to identify
554 people with a lifetime diagnosis of psychotic disorder. *BMC Res. Notes* 5, e95.
555 <https://doi.org/10.1186/1756-0500-5-95>
- 556 Faris, R.E.L., Dunham, H.W., 1939. *Mental Disorders in Urban Areas: An Ecological Study
557 of Schizophrenia and Other Psychoses*. University of Chicago Press, Oxford, England.
- 558 Fichera, E., Gray, E., Sutton, M., 2016. How do individuals' health behaviours respond to an
559 increase in the supply of health care? Evidence from a natural experiment. *Soc. Sci.
560 Med.* 159, 170–179. <https://doi.org/https://doi.org/10.1016/j.socscimed.2016.05.005>
- 561 Ford, D. V., Jones, K.H., Verplancke, J., Lyons, R.A., John, G., Brown, G., Brooks, C.J.,
562 Thompson, S., Bodger, O., Couch, T., Leake, K., 2009. The SAIL Databank: building a
563 national architecture for e-health research and evaluation. *BMC Health Serv. Res.* 9,
564 157. <https://doi.org/10.1186/1472-6963-9-157>
- 565 Gamble, A., 2009. *The spectre at the feast: Capitalist crisis and the politics of recession*.
566 Macmillan International Higher Education, London.
- 567 Goldberg, E.M., Morrison, S.L., 1963. Schizophrenia and social class. *Br. J. Psychiatry* 109,
568 785–802. <https://doi.org/10.1192/bjp.109.463.785>
- 569 Gruebner, O., Rapp, M.A., Adli, M., Kluge, U., Galea, S., Heinz, A., 2017. Cities and Mental
570 Health. *Dtsch. Arztebl. Int.* 114, 121. <https://doi.org/10.3238/arztebl.2017.0121>
- 571 Hardoon, S., Hayes, J.F., Blackburn, R., Petersen, I., Walters, K., Nazareth, I., Osborn,
572 D.P.J., 2013. Recording of severe mental illness in United Kingdom primary care, 2000-
573 2010. *PLoS One* 8, e82365. <https://doi.org/10.1371/journal.pone.0082365>
- 574 Harrington, M., 2010. *An Independent Review of the Work Capability Assessment*. London.
- 575 Haukka, J., Suvisaari, J., Varilo, T., Lnnqvist, J., 2001. Regional variation in the incidence of
576 schizophrenia in Finland: A study of birth cohorts born from 1950 to 1969. *Psychol.
577 Med.* 31, 1046–1053. <https://doi.org/10.1017/S0033291701004299>
- 578 Heinz, A., Deserno, L., Reininghaus, U., 2013. Urbanicity, social adversity and psychosis.
579 *World Psychiatry* 12, 187–197. <https://doi.org/10.1002/wps.20056>
- 580 Hudson, C.G., 2012. Patterns of residential mobility of people with schizophrenia: Multi-level
581 tests of downward geographic drift. *J. Sociol. Soc. Welf.* 39, 149.
- 582 Hudson, C.G., 2005. Socioeconomic status and mental illness: Tests of the social causation
583 and selection hypotheses. *Am. J. Orthopsychiatry* 75, 3–18.
584 <https://doi.org/10.1037/0002-9432.75.1.3>
- 585 John, A., McGregor, J., Fone, D., Dunstan, F., Cornish, R., Lyons, R.A., Lloyd, K.R., 2016.
586 Case-finding for common mental disorders of anxiety and depression in primary care:
587 An external validation of routinely collected data. *BMC Med. Inform. Decis. Mak.* 16, 35.
588 <https://doi.org/10.1186/s12911-016-0274-7>
- 589 John, A., McGregor, J., Jones, I., Lee, S.C., Walters, J.T.R., Owen, M.J., O'Donovan, M.,

- 590 DelPozo-Banos, M., Berridge, D., Lloyd, K., 2018. Premature mortality among people
591 with severe mental illness — New evidence from linked primary care data. *Schizophr.*
592 *Res.* <https://doi.org/10.1016/j.schres.2018.04.009>
- 593 Jongsma, H.E., Gayer-Anderson, C., Lasalvia, A., Quattrone, D., Mulè, A., Szöke, A., Selten,
594 J.P., Turner, C., Arango, C., Tarricone, I., Berardi, D., Tortelli, A., Llorca, P.M., De
595 Haan, L., Bobes, J., Bernardo, M., Sanjuán, J., Santos, J.L., Arrojo, M., Del-Ben, C.M.,
596 Menezes, P.R., Murray, R.M., Rutten, B.P., Jones, P.B., Van Os, J., Morgan, C.,
597 Kirkbride, J.B., Reininghaus, U., Di Forti, M., Hubbard, K., Beards, S., Stilo, S.A.,
598 Tripoli, G., Parellada, M., Cuadrado, P., Solano, J.J.R., Carracedo, A., Bernardo, E.G.,
599 Roldán, L., López, G., Cabrera, B., Lorente-Rovira, E., Garcia-Portilla, P., Costas, J.,
600 Jiménez-López, E., Matteis, M., Rapado, M., González, E., Martínez, C., Sánchez, E.,
601 Olmeda, M.S., Franke, N., Velthorst, E., Termorshuizen, F., Van Dam, D., Van Der
602 Ven, E., Messchaert, E., Leboyer, M., Schürhoff, F., Jamain, S., Frijda, F., Baudin, G.,
603 Ferchiou, A., Pignon, B., Richard, J.R., Charpeaud, T., Tronche, A.M., La Barbera, D.,
604 La Cascia, C., Marrazzo, G., Sideli, L., Sartorio, C., Ferraro, L., Seminerio, F., Loureiro,
605 C.M., Shuhama, R., Ruggeri, M., Tosato, S., Bonetto, C., Cristofalo, D., 2018. Treated
606 incidence of psychotic disorders in the multinational EU-GEI study. *JAMA Psychiatry*
607 *75*, 36–46. <https://doi.org/10.1001/jamapsychiatry.2017.3554>
- 608 Jongsma, H.E., Turner, C., Kirkbride, J.B., Jones, P.B., 2019. International incidence of
609 psychotic disorders, 2002–17: a systematic review and meta-analysis. *Lancet Public*
610 *Heal.* *4*, E229–E244. [https://doi.org/https://doi.org/10.1016/S2468-2667\(19\)30056-8](https://doi.org/https://doi.org/10.1016/S2468-2667(19)30056-8)
- 611 Karanikolos, M., Mladovsky, P., Cylus, J., Thomson, S., Basu, S., Stuckler, D., Mackenbach,
612 J.P., McKee, M., 2013. Financial crisis, austerity, and health in Europe. *Lancet* *381*,
613 1323–1331. [https://doi.org/10.1016/S0140-6736\(13\)60102-6](https://doi.org/10.1016/S0140-6736(13)60102-6)
- 614 Katikireddi, S.V., Niedzwiedz, C.L., Popham, F., 2012. Trends in population mental health
615 before and after the 2008 recession: a repeat cross-sectional analysis of the 1991–
616 2010 Health Surveys of England. *BMJ Open* *2*, e001790.
617 <https://doi.org/10.1136/bmjopen-2012-001790>
- 618 Kaymaz, N., Krabbendam, L., Graaf, R., Nolen, W., Have, M., van Os, J., 2006. Evidence
619 that the urban environment specifically impacts on the psychotic but not the affective
620 dimension of bipolar disorder. *Soc. Psychiatry Psychiatr. Epidemiol.* *41*, 679–685.
621 <https://doi.org/10.1007/s00127-006-0086-7>
- 622 Kelly, B.D., O’Callaghan, E., Waddington, J.L., Feeney, L., Browne, S., Scully, P.J., Clarke,
623 M., Quinn, J.F., McTigue, O., Morgan, M.G., Kinsella, A., Larkin, C., 2010.
624 Schizophrenia and the city: A review of literature and prospective study of psychosis
625 and urbanicity in Ireland. *Schizophr. Res.* *116*, 75–89.
626 <https://doi.org/10.1016/j.schres.2009.10.015>
- 627 Kendrick, T., Stuart, B., Newell, C., Geraghty, A.W.A., Moore, M., 2015. Changes in rates of
628 recorded depression in English primary care 2003-2013: Time trend analyses of effects
629 of the economic recession, and the GP contract quality outcomes framework (QOF). *J.*
630 *Affect. Disord.* *180*, 68–78. <https://doi.org/10.1016/j.jad.2015.03.040>
- 631 Kim, H.J., Fay, M.P., Feuer, E.J., Midthune, D.N., 2000. Permutation tests for joinpoint
632 regression with applications to cancer rates. *Stat. Med.* *19*, 335–351.
633 [https://doi.org/10.1002/\(SICI\)1097-0258\(20000215\)19:3<335::AID-SIM336>3.0.CO;2-Z](https://doi.org/10.1002/(SICI)1097-0258(20000215)19:3<335::AID-SIM336>3.0.CO;2-Z)
- 634 Kirkbride, J.B., Errazuriz, A., Croudace, T.J., Morgan, C., Jackson, D., Boydell, J., Murray,
635 R.M., Jones, P.B., 2012. Incidence of schizophrenia and other psychoses in England,
636 1950-2009: A systematic review and meta-analyses. *PLoS One* *7*, e31660.
637 <https://doi.org/10.1371/journal.pone.0031660>

- 638 Kitson, M., Martin, R., Tyler, P., 2011. The geographies of austerity. *Cambridge J. Reg.*
639 *Econ. Soc.* 4, 289–302. <https://doi.org/10.1093/cjres/rsr030>
- 640 Laursen, T.M., Munk-Olsen, T., Nordentoft, M., Bo Mortensen, P., 2007. A comparison of
641 selected risk factors for unipolar depressive disorder, bipolar affective disorder,
642 schizoaffective disorder, and schizophrenia from a danish population-based cohort. *J.*
643 *Clin. Psychiatry* 68, 1673–1681. <https://doi.org/10.4088/JCP.v68n1106>
- 644 Lee, S.C., DelPozo-Banos, M., Lloyd, K., Jones, I., Walters, J.T.R., Owen, M.J., O'Donovan,
645 M., John, A., 2020. Area deprivation, urbanicity, severe mental illness and social drift —
646 A population-based linkage study using routinely collected primary and secondary care
647 data. *Schizophr. Res.* <https://doi.org/10.1016/J.SCHRES.2020.03.044>
- 648 Litchfield, P., 2014. *An Independent Review of the Work Capability Assessment – year five.*
649 London.
- 650 Lloyd, K., McGregor, J., John, A., Craddock, N., Walters, J.T., Linden, D., Jones, I., Bentall,
651 R., Lyons, R.A., Ford, D. V., Owen, M.J., 2015. A national population-based e-cohort of
652 people with psychosis (PsyCymru) linking prospectively ascertained phenotypically rich
653 and genetic data to routinely collected records: Overview, recruitment and linkage.
654 *Schizophr. Res.* 166, 131–136. <https://doi.org/10.1016/j.schres.2015.05.036>
- 655 Lyons, R. a, Jones, K.H., John, G., Brooks, C.J., Verplancke, J.-P., Ford, D. V, Brown, G.,
656 Leake, K., 2009. The SAIL databank: linking multiple health and social care datasets.
657 *BMC Med. Inform. Decis. Mak.* 9, 3. <https://doi.org/10.1186/1472-6947-9-3>
- 658 Marcelis, M., Navarro-Mateu, F., Murray, R., Selten, J.P., Van Os, J., 1998. Urbanization
659 and psychosis: A study of 1942-1978 birth cohorts in The Netherlands. *Psychol. Med.*
660 28, 871–879. <https://doi.org/10.1017/S0033291798006898>
- 661 March, D., Hatch, S.L., Morgan, C., Kirkbride, J.B., Bresnahan, M., Fearon, P., Susser, E.,
662 2008. Psychosis and place. *Epidemiol. Rev.* 30, 84–100.
663 <https://doi.org/10.1093/epirev/mxn006>
- 664 McLintock, K., Russell, A.M., Alderson, S.L., West, R., House, A., Westerman, K., Foy, R.,
665 2014. The effects of financial incentives for case finding for depression in patients with
666 diabetes and coronary heart disease: interrupted time series analysis. *BMJ Open* 4,
667 e005178. <https://doi.org/10.1136/bmjopen-2014-005178>
- 668 National Cancer Institute, 2015. *Joinpoint Regression Program - Surveillance Research*
669 *Program.* June.
- 670 Newcombe, R.G., 1998. Two-sided confidence intervals for the single proposition:
671 comparison of seven methods. *Stat. Med.* 17, 857–872.
672 [https://doi.org/10.1002/\(SICI\)1097-0258\(19980430\)17](https://doi.org/10.1002/(SICI)1097-0258(19980430)17)
- 673 Niedzwiedz, C.L., Mitchell, R.J., Shortt, N.K., Pearce, J.R., 2016. Social protection spending
674 and inequalities in depressive symptoms across Europe. *Soc. Psychiatry Psychiatr.*
675 *Epidemiol.* 51, 1005–1014. <https://doi.org/10.1007/s00127-016-1223-6>
- 676 Norman, P.D., 2018. Clarity in research frameworks for studying 'health selective migration.'
677 *J. Epidemiol. Community Health* 72, 449 LP – 450. [https://doi.org/10.1136/jech-2018-](https://doi.org/10.1136/jech-2018-210678)
678 [210678](https://doi.org/10.1136/jech-2018-210678)
- 679 O'Donoghue, B., Roche, E., Lane, A., 2016. Neighbourhood level social deprivation and the
680 risk of psychotic disorders: a systematic review. *Soc. Psychiatry Psychiatr. Epidemiol.*
681 51, 941–950. <https://doi.org/10.1007/s00127-016-1233-4>

- 682 O'Donoghue, D.J., 2009. GOING UPSTREAM: THE IMPLICATION AND OPPORTUNITIES
683 OF EARLY DETECTION. *J. Ren. Care* 35, 3–7.
684 <https://doi.org/https://doi.org/10.1111/j.1755-6686.2009.00126.x>
- 685 ONS, 2018. The 2008 recession 10 years on [WWW Document]. URL
686 <https://www.ons.gov.uk/economy/grossdomesticproductgdp/articles/the2008recession10yearson/2018-04-30> (accessed 4.22.20).
687
- 688 Paksarian, D., Trabjerg, B.B., Merikangas, K.R., Mors, O., Borglum, A.D., Hougaard, D.M.,
689 McGrath, J.J., Pedersen, C.B., Mortensen, P.B., Agerbo, E., 2018. The role of genetic
690 liability in the association of urbanicity at birth and during upbringing with schizophrenia
691 in Denmark. *Psychol. Med.* 48, 305–314. <https://doi.org/10.1017/S0033291717001696>
- 692 Pateman, T., 2011. Rural and urban areas : comparing lives using rural / urban
693 classifications. *Reg. trends* 43, 11–86. <https://doi.org/10.1057/rt.2011.2>
- 694 Pedersen, C.B., 2006. No evidence of time trends in the urban-rural differences in
695 schizophrenia risk among five million people born in Denmark from 1910 to 1986.
696 *Psychol. Med.* 36, 211–219. <https://doi.org/10.1017/S003329170500663X>
- 697 Plomin, R., Deary, I.J., 2015. Genetics and intelligence differences: Five special findings.
698 *Mol. Psychiatry* 20, 98. <https://doi.org/10.1038/mp.2014.105>
- 699 Radua, J., Ramella-Cravaro, V., Ioannidis, J.P.A., Reichenberg, A., Phiphophthasanee, N.,
700 Amir, T., Yenn Thoo, H., Oliver, D., Davies, C., Morgan, C., McGuire, P., Murray, R.M.,
701 Fusar-Poli, P., 2018. What causes psychosis? An umbrella review of risk and protective
702 factors. *World Psychiatry* 17, 49–66. <https://doi.org/10.1002/wps.20490>
- 703 Reeves, A., Basu, S., McKee, M., Marmot, M., Stuckler, D., 2013. Austerity's health effects:
704 a comparative analysis of European budgetary changes: Aaron Reeves. *Eur. J. Public*
705 *Health* 23. <https://doi.org/10.1093/eurpub/ckt126.148>
- 706 Reilly, S., Olier, I., Planner, C., Doran, T., Reeves, D., Ashcroft, D.M., Gask, L.,
707 Kontopantelis, E., 2015. Inequalities in physical comorbidity: A longitudinal comparative
708 cohort study of people with severe mental illness in the UK. *BMJ Open* 5, e009010.
709 <https://doi.org/10.1136/bmjopen-2015-009010>
- 710 Rothman, K., Boice, J.J., 1979. Epidemiologic analysis with a programmable calculator, NIH
711 Publication 79–1649. Washington, DC.
- 712 Saha, S., Chant, D., Welham, J., McGrath, J., 2005. A systematic review of the prevalence
713 of schizophrenia. *PLoS Med.* 2, e141. <https://doi.org/10.1371/journal.pmed.0020141>
- 714 Sariaslan, A., Fazel, S., D'Onofrio, B.M., Långström, N., Larsson, H., Bergen, S.E., Kuja-
715 Halkola, R., Lichtenstein, P., 2016. Schizophrenia and subsequent neighborhood
716 deprivation: Revisiting the social drift hypothesis using population, twin and molecular
717 genetic data. *Transl. Psychiatry* 6, e796. <https://doi.org/10.1038/tp.2016.62>
- 718 Sariaslan, A., Larsson, H., D'Onofrio, B., Långström, N., Fazel, S., Lichtenstein, P., 2015.
719 Does Population Density and Neighborhood Deprivation Predict Schizophrenia? A
720 Nationwide Swedish Family-Based Study of 2.4 Million Individuals. *Schizophr. Bull.* 41,
721 494–502. <https://doi.org/10.1093/schbul/sbu105>
- 722 Selten, J.P., Van Der Ven, E., Rutten, B.P.F., Cantor-Graae, E., 2013. The social defeat
723 hypothesis of schizophrenia: An update. *Schizophr. Bull.* 39, 1180–1186.
724 <https://doi.org/10.1093/schbul/sbt134>
- 725 Simeone, J.C., Ward, A.J., Rotella, P., Collins, J., Windisch, R., 2015. An evaluation of

726 variation in published estimates of schizophrenia prevalence from 1990-2013: A
727 systematic literature review. *BMC Psychiatry* 15, 193. [https://doi.org/10.1186/s12888-](https://doi.org/10.1186/s12888-015-0578-7)
728 015-0578-7

729 StataCorp, 2019. *Stata Statistical Software: Release 16*. StataCorp LLC, College Station,
730 TX.

731 Taylor-Robinson, D., Whitehead, M., Barr, B., 2014. Great leap backwards. *BMJ Br. Med. J.*
732 349, g7350. <https://doi.org/10.1136/bmj.g7350>

733 The UK government, 2013. *Employment and Support Allowance: outcomes of Work*
734 *Capability Assessments [WWW Document]*. URL
735 [https://www.gov.uk/government/collections/employment-and-support-allowance-](https://www.gov.uk/government/collections/employment-and-support-allowance-outcomes-of-work-capability-assessment)
736 [outcomes-of-work-capability-assessment](https://www.gov.uk/government/collections/employment-and-support-allowance-outcomes-of-work-capability-assessment) (accessed 4.22.20).

737 The UK government, 2012. *Welfare Reform Act 2012 [WWW Document]*. URL
738 <http://www.legislation.gov.uk/ukpga/2012/5/contents/enacted> (accessed 4.22.20).

739 Van Os, J., Rutten, B.P.F., Poulton, R., 2008. Gene-environment interactions in
740 schizophrenia: Review of epidemiological findings and future directions. *Schizophr.*
741 *Bull.* 34, 1066–1082. <https://doi.org/10.1093/schbul/sbn117>

742 Vassos, E., Pedersen, C.B., Murray, R.M., Collier, D.A., Lewis, C.M., 2012. Meta-analysis of
743 the association of urbanicity with schizophrenia. *Schizophr. Bull.* 38, 1118–1123.
744 <https://doi.org/10.1093/schbul/sbs096>

745 Wagner, A.K., Soumerai, S.B., Zhang, F., Ross-Degnan, D., 2002. Segmented regression
746 analysis of interrupted time series studies in medication use research. *J. Clin. Pharm.*
747 *Ther.* 27, 299–309. <https://doi.org/10.1046/j.1365-2710.2002.00430.x>

748 Welsh Government, 2017. *Welsh Index of Multiple Deprivation (WIMD) 2014 Revised*.
749 Cardiff, UK.

750 Wilkinson R, Pickett K (2009) *The Spirit level. Why equality is better for everyone*. Penguin Books,
751 London

752