

# **The importance of tropical tree-ring chronologies for global change research**

Peter GROENENDIJK<sup>1</sup>, Flurin BABST<sup>2,3</sup>, Valerie TROUET<sup>3,4</sup>, Ze-Xin FAN<sup>5</sup>, Daniela GRANATO-SOUZA<sup>6</sup>, Giuliano Maselli LOCOSSELLI<sup>7,8</sup>, Mulugeta MOKRIA<sup>9,10</sup>, Shankar PANTHI<sup>5</sup>, Nathsuda PUMIJUMNONG<sup>11</sup>, Abraham ABIYU<sup>12</sup>, Rodolfo ACUÑA-SOTO<sup>13,14</sup>, Eduardo ADENESKY-FILHO<sup>15</sup>, Raquel ALFARO-SÁNCHEZ<sup>16,17</sup>, Claudio Roberto ANHOLETTO JUNIOR<sup>18</sup>, José Roberto Vieira ARAGÃO<sup>19,1</sup>, Gabriel ASSIS-PEREIRA<sup>20,21</sup>, Claudia C. ASTUDILLO-SÁNCHEZ<sup>22</sup>, Ana Carolina BARBOSA<sup>23</sup>, Nathan de Oliveira BARRETO<sup>20</sup>, Giovanna BATTIPAGLIA<sup>24</sup>, Hans BEECKMAN<sup>25</sup>, Paulo Cesar BOTOSSO<sup>26</sup>, Nils BOURLAND<sup>25</sup>, Achim BRÄUNING<sup>9</sup>, Roel BRIENEN<sup>27</sup>, Matthew BROOKHOUSE<sup>28</sup>, Supaporn BUAJAN<sup>11</sup>, Brendan M. BUCKLEY<sup>29</sup>, J. Julio CAMARERO<sup>30</sup>, Artemio CARRILLO-PARRA<sup>31</sup>, Gregório CECCANTINI<sup>7</sup>, Librado R. CENTENO-ERGUERA<sup>32</sup>, Julián CERANO-PAREDES<sup>32</sup>, Rosalinda CERVANTES-MARTÍNEZ<sup>33</sup>, Wirong CHANTHORN<sup>34</sup>, Ya-Jun CHEN<sup>5</sup>, Bruno Barçante Ladvoat CINTRA<sup>35,36</sup>, Eladio Heriberto CORNEJO-OVIEDO<sup>33</sup>, Otoniel CORTÉS-CORTÉS<sup>33</sup>, Clayane Matos COSTA<sup>37</sup>, Camille COURALET<sup>25</sup>, Doris Bianca CRISPÍN-DE-LA-CRUZ<sup>1,38</sup>, Rosanne D'ARRIGO<sup>29</sup>, Diego A. DAVID<sup>39</sup>, Maaike DE RIDDER<sup>25</sup>, Jorge Ignacio DEL VALLE<sup>39</sup>, Oscar A. DÍAZ-CARRILLO<sup>31</sup>, Mário DOBNER JR<sup>40</sup>, Jean-Louis DOUCET<sup>41</sup>, Oliver DÜNISCH<sup>42</sup>, Brian J. ENQUIST<sup>43,44</sup>, Karin ESEMANN-QUADROS<sup>45,46</sup>, Gerardo ESQUIVEL-ARRIAGA<sup>32</sup>, Adeline FAYOLLE<sup>47,48</sup>, Tatiele Anete Bergamo FENILLI<sup>15</sup>, M. Eugenia FERRERO<sup>49,38</sup>, Esther FICHTLER<sup>50</sup>, Patrick M. FINNEGAN<sup>51</sup>, Claudia FONTANA<sup>20</sup>, Kainana S. FRANCISCO<sup>52</sup>, Pei-Li FU<sup>5</sup>, Franklin GALVÃO<sup>53</sup>, Aster GEBREKIRSTOS<sup>10</sup>, Jorge A. GIRALDO<sup>54</sup>, Emanuel GLOOR<sup>27</sup>, Milena GODOY-VEIGA<sup>7</sup>, Anthony GUERRA<sup>55</sup>, Kristof HANECA<sup>56</sup>, Grant Logan HARLEY<sup>57</sup>, Ingo HEINRICH<sup>58,59</sup>, Gerhard HELLE<sup>60</sup>, José Ciro HERNÁNDEZ-DÍAZ<sup>31</sup>, Bruna HORNINK<sup>1,20</sup>, Wannes HUBAU<sup>61</sup>, Janet G. INGA<sup>38</sup>, Mahmuda ISLAM<sup>62,63</sup>, Yumei JIANG<sup>64</sup>, Mark KAIB<sup>65</sup>, Zakia Hassan KHAMISI<sup>3</sup>, Marcin KOPROWSKI<sup>66,67</sup>, Eva LAYME<sup>68</sup>, A. Joshua LEFFLER<sup>69</sup>, Gauthier LIGOT<sup>41</sup>, Claudio Sergio LISI<sup>37</sup>, Neil J. LOADER<sup>70</sup>, Francisco de Almeida LOBO<sup>71</sup>, Tomaz LONGHI-SANTOS<sup>72</sup>, Lidio LOPEZ<sup>49</sup>, María I. LÓPEZ-HERNÁNDEZ<sup>33</sup>, José Luís Penetra Cerveira LOUSADA<sup>73</sup>, Rubén D. MANZANEDO<sup>74,75</sup>, Amanda K. MARCON<sup>53</sup>, Justin T. MAXWELL<sup>76</sup>, Hooz A. MENDIVELSO<sup>77</sup>, Omar N. MENDOZA-VILLA<sup>78</sup>, Ítallo Romany Nunes MENEZES<sup>37,79</sup>, Valdiney Ribeiro MONTÓIA<sup>80</sup>, Eddy MOORS<sup>81,82</sup>, Miyer MORENO<sup>83</sup>, Miguel Angel MUÑIZ-CASTRO<sup>84</sup>, Cristina NABAIS<sup>85</sup>, Anuttara NATHALANG<sup>86</sup>, Justine NGOMA<sup>87</sup>, Francisco de Carvalho NOGUEIRA JR.<sup>88</sup>, Juliano Morales OLIVEIRA<sup>89,90</sup>, Gabriela Morais OLMEDO<sup>91</sup>, Daigard Ricardo ORTEGA-RODRIGUEZ<sup>20</sup>, Carmen Eugenia Rodríguez ORTÍZ<sup>92</sup>, Mariana Alves PAGOTTO<sup>37</sup>, Kathelyn PAREDES-VILLANUEVA<sup>93,94</sup>, Gonzalo PÉREZ-DE-LIS<sup>95</sup>, Laura Patricia PONCE CALDERÓN<sup>96</sup>, Leif Armando PORTAL-CAHUANA<sup>97,20</sup>, Darwin Alexander PUCHA-COFREP<sup>98</sup>, Paulo QUADRI<sup>99</sup>, Mizanur RAHMAN<sup>62</sup>, Jorge Andrés RAMÍREZ<sup>100</sup>, Edilson Jimmy REQUENA-ROJAS<sup>38</sup>, Judith REYES-FLORES<sup>33</sup>, Adauto de Souza RIBEIRO<sup>37</sup>, Iain ROBERTSON<sup>70</sup>, Fidel Alejandro ROIG<sup>49,101</sup>, José Guilherme ROQUETTE<sup>102</sup>, Ernesto Alonso RUBIO-CAMACHO<sup>103</sup>, Raúl SÁNCHEZ-SALGUERO<sup>104</sup>, Ute SASS-KLAASSEN<sup>105,106</sup>, Jochen SCHÖNGART<sup>107</sup>, Marcelo Callegari SCIPIONI<sup>40</sup>, Paul R. SHEPPARD<sup>3</sup>, Lucas C.R. SILVA<sup>108</sup>, Franziska SLOTTA<sup>109</sup>, Leroy SORIA-DÍAZ<sup>110</sup>, Luciana K.V.S. SOUSA<sup>111</sup>, James H. SPEER<sup>112</sup>, Matthew D. THERRELL<sup>113</sup>, Ginette TICSE-OTAROLA<sup>38,114</sup>, Mario TOMAZELLO-FILHO<sup>20</sup>, Max C.A. TORBENSON<sup>115</sup>, Pantana TOR-NGERN<sup>116,117</sup>, Ramzi TOUCHAN<sup>3</sup>, Jan VAN DEN BULCKE<sup>61</sup>, Lorenzo VÁZQUEZ-SELEM<sup>118</sup>, Adín H. VELÁZQUEZ-PÉREZ<sup>119</sup>, Alejandro VENEGAS-GONZÁLEZ<sup>120</sup>, Ricardo VILLALBA<sup>49</sup>, Jose VILLANUEVA-DIAZ<sup>32</sup>, Mart VLAM<sup>106</sup>, George VOURLITIS<sup>121</sup>, Christian WEHENKEL<sup>31</sup>, Tommy WILS<sup>122,123</sup>, Erika S. ZAVALETA<sup>124</sup>, Eshetu Asfaw ZEWDU<sup>125</sup>, Yong-Jiang ZHANG<sup>126</sup>, Zhe-Kun ZHOU<sup>5</sup>, Pieter A. ZUIDEMA<sup>105</sup>

<sup>1</sup>Department of Plant Biology, Institute of Biology, P.O. Box: 6109, University of Campinas (UNICAMP), 13083-970, Campinas, SP, Brazil. <sup>2</sup>School of Natural Resources and the Environment, University of Arizona, Tucson, AZ, USA. <sup>3</sup>Laboratory of Tree-Ring Research, University of Arizona, Tucson, AZ, USA. <sup>4</sup>Belgian Climate Centre, Ringlaan 3, 1090 Uccle, Belgium. <sup>5</sup>CAS Key Laboratory of Tropical Forest Ecology,

47 Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Mengla, Yunnan, 666303, China.  
 48 <sup>6</sup>Natural Resources and Environmental Sciences Department, Forestry, Ecology and Wildlife Program, Alabama  
 49 A&M University, Alabama, US. <sup>7</sup>Department of Botany, Institute of Biosciences, University of São Paulo, R. do  
 50 Matão, 277, São Paulo, 05508-090, Brazil. <sup>8</sup>Center of Nuclear Energy in Agriculture, University of São Paulo,  
 51 Av. Centenário 303, 13416-000, Brazil. <sup>9</sup>Institute of Geography, Friedrich-Alexander-University Erlangen-  
 52 Nuremberg, Wetterkreuz 15, 91058, Erlangen, Germany. <sup>10</sup>World Agroforestry Centre (ICRAF), United Nations  
 53 Avenue, P.O. Box 30677-00100, Nairobi, Kenya. <sup>11</sup>Faculty of Environment and Resource studies, Mahidol  
 54 University, 999 Phutthamonthon Rd4, Salaya, Phutthamonthon, Nakhon Pathom 73170, Thailand. <sup>12</sup>World  
 55 Agroforestry Centre (ICRAF), C/O ILRI Campus, Gurd Shola, P.O. Box 5689, Addis Ababa, Ethiopia.  
 56 <sup>13</sup>Department of Microbiology and Parasitology, Universidad Nacional Autónoma de México, Mexico City,  
 57 Mexico. <sup>14</sup>Colegio de Geografía, Facultad de Filosofía y Letras, Universidad Nacional Autónoma de México,  
 58 Mexico City, Mexico. <sup>15</sup>Department of Forest Engineering, Universidade Regional de Blumenau, R. São Paulo  
 59 3366, Itoupava Seca, Santa Catarina, 89030-000, Brazil. <sup>16</sup>Northern Forestry Centre, Canadian Forest Service,  
 60 Natural Resources Canada, 5320 122 Street NW, Edmonton, AB, T6H 3S5, Canada. <sup>17</sup>Department of Biology,  
 61 Wilfrid Laurier University, 75 University Avenue W, Waterloo, ON, N2L 3C5, Canada. <sup>18</sup>Forest Management  
 62 Program, Mamirauá Institute for Sustainable Development, Tefé, Brazil. <sup>19</sup>Department of Plant Biology, Federal  
 63 University of Ceará (UFC), Av. Mister Hull S/N - PICI Campus, 60440-900, Fortaleza, CE, Brazil. <sup>20</sup>Department  
 64 of Forest Sciences, Luiz de Queiroz College of Agriculture, University of São Paulo, Av. Pádua Dias 11,  
 65 Piracicaba - SP, 13418-900, Brazil. <sup>21</sup>Instituto de Pesquisas Ambientais do Estado de São Paulo, Assis, São  
 66 Paulo, Brazil. <sup>22</sup>Facultad de Ingeniería y Ciencias, Universidad Autónoma de Tamaulipas, 87149, Ciudad  
 67 Victoria, Tamaulipas, Mexico. <sup>23</sup>Tree-Ring Laboratory, Forest Science Department, Federal University of  
 68 Lavras, 37200-900, Lavras, MG, Brazil. <sup>24</sup>University of Campania "L. Vanvitelli", Department of  
 69 Environmental, Biological and Pharmaceutical Sciences and Technologies, via Vivaldi 43, 81100, Caserta, Italy.  
 70 <sup>25</sup>Wood Biology Service, Royal Museum for Central Africa, Leuvense steenweg 13, 3080 Tervuren, Belgium.  
 71 <sup>26</sup>Brazilian Agricultural Research Corporation (Embrapa), Embrapa Forestry, Estrada da Ribeira - Km 111,  
 72 83411-000, Colombo, PR, Brazil. <sup>27</sup>School of Geography, University of Leeds, Leeds, LS2 9JT, UK. <sup>28</sup>Fenner  
 73 School of Environment and Society, Australian National University, Canberra, Australian Capital Territory,  
 74 Australia. <sup>29</sup>Lamont-Doherty Earth Observatory, Columbia University, Palisades, New York, USA. <sup>30</sup>Instituto  
 75 Pirenaico de Ecología (IPE-CSIC), E-50192 Zaragoza, Spain. <sup>31</sup>Instituto de Silvicultura e Industria de la Madera,  
 76 Universidad Juárez del Estado de Durango, Durango 34120, Mexico. <sup>32</sup>Instituto Nacional de Investigaciones  
 77 Forestales, Agrícolas y Pecuarias (INIFAP), Centro Nacional de Investigación Disciplinaria en Relación Agua-  
 78 Suelo-Planta-Atmósfera (CENID-RASPA), Margen Derecha del Canal del Sacramento Km 6.5, CP 35140,  
 79 Gómez Palacio, Durango, Mexico. <sup>33</sup>Departamento Forestal, Universidad Autónoma Agraria Antonio Narro,  
 80 Calzada Antonio Narro 1923, Buenavista CP 25315, Saltillo, Coahuila, Mexico. <sup>34</sup>Department of Environmental  
 81 Technology and Management, Faculty of Environment, Kasetsart University, 50 Ngamwongwan Road, Jatujak  
 82 District, Bangkok, 10900, Thailand. <sup>35</sup>School of Geography, Earth and Environmental Sciences, University of  
 83 Birmingham, UK. <sup>36</sup>Birmingham Institute of Forest Research, University of Birmingham, UK. <sup>37</sup>Laboratory of  
 84 Plant Anatomy and Dendrochronology, Department of Biology, Universidade Federal de Sergipe - UFS, Aracaju  
 85 - SE, 49100-000, Brazil. <sup>38</sup>Laboratorio de Dendrocronología, Universidad Continental, Av. San Carlos 1980,  
 86 Huancayo, Perú. <sup>39</sup>Department of Forest Sciences, Universidad Nacional de Colombia - Sede Medellín,  
 87 Colombia. <sup>40</sup>Departamento de Agricultura, Biodiversidade e Florestas, Universidade Federal de Santa Catarina –  
 88 UFSC, Curitiba, Santa Catarina, Brazil. <sup>41</sup>TERRA Teaching and Research Centre, Gembloux Agro-Bio Tech,  
 89 University of Liège, Gembloux, Belgium. <sup>42</sup>Master School for Carpentry and Cabinetmaking, Gleusdorfer Str.  
 90 14, D-96106 Ebern, Germany. <sup>43</sup>Department of Ecology and Evolutionary Biology, University of Arizona,  
 91 Tucson, AZ 85721, USA. <sup>44</sup>Santa Fe Institute, 1399 Hyde Park Rd., Santa Fe, NM 87501, USA. <sup>45</sup>Department of  
 92 Biological Sciences, University of Joinville Region - UNIVILLE, Joinville, SC, 89219-710, Brazil.  
 93 <sup>46</sup>Postgraduate Program in Forestry, Regional University of Blumenau - FURB, Blumenau, SC, 89030-000,  
 94 Brazil. <sup>47</sup>UR Forêts et Sociétés du CIRAD - Campus de Baillarguet TA C-22E, 34398 Montpellier cedex 5,

95 France. <sup>48</sup>Gestion des Ressources Forestières de Gembloux Agro-Bio Tech, ULiège, Passage des Déportés 2,  
96 5030 Gembloux, Belgium. <sup>49</sup>Laboratorio de Dendrocronología e Historia Ambiental, IANIGLA, CCT-  
97 CONICET-Mendoza, Av. Ruiz Leal s/n, Parque Gral. San Martín, CC 330, CP 5500, Mendoza, Argentina.  
98 <sup>50</sup>Department of Crop Sciences, Tropical Plant Production and Agricultural Systems Modelling, Göttingen  
99 University, Grisebachstrasse 6, Göttingen, 37077, Germany. <sup>51</sup>School of Biological Sciences, University of  
100 Western Australia, Perth WA 6009, Australia. <sup>52</sup>USDA Forest Service, Pacific Southwest Research Station,  
101 Institute of Pacific Islands Forestry, Hilo, HI, USA. <sup>53</sup>Department of Forest Sciences, Federal University of  
102 Paraná, Curitiba - PR, Brazil. <sup>54</sup>Facultad de Ingeniería, Tecnológico de Antioquia, Medellín, Colombia.  
103 <sup>55</sup>Programa de Pós-Graduação em Agronomia/Fisiologia Vegetal, Departamento de Biologia, Instituto de  
104 Ciências Naturais, Universidade Federal de Lavras, 7203-202 Lavras, MG, Brazil. <sup>56</sup>Flanders Heritage Agency,  
105 Havenlaan 88 box 5, 1000 Brussels, Belgium. <sup>57</sup>Department of Earth and Spatial Sciences, University of Idaho,  
106 Moscow, ID, USA. <sup>58</sup>German Archaeological Institute DAI, Department of Natural Sciences, 14195, Berlin,  
107 Germany. <sup>59</sup>Geography Department, Humboldt-University Berlin, 12489, Berlin, Germany. <sup>60</sup>GFZ German  
108 Research Centre for Geosciences, 14473, Potsdam, Germany. <sup>61</sup>Faculty of Bioscience Engineering, Department  
109 of Environment, Laboratory of Wood Technology (UGent-Woodlab), Ghent University, Ghent, Belgium.  
110 <sup>62</sup>Department of Forestry and Environmental Science, Shahjalal University of Science and Technology, Sylhet  
111 3114, Bangladesh. <sup>63</sup>School of Biological Sciences, Washington State University, PO Box 644236, Pullman, WA  
112 99164-4236, USA. <sup>64</sup>Faculty of Forestry and Wood Sciences, Czech University of Life Sciences Prague, Prague,  
113 Czech Republic. <sup>65</sup>U.S. Fish and Wildlife Service, Region 2, 500 Gold Ave. SW, PO BOX 1306, Albuquerque,  
114 NM 87103, USA. <sup>66</sup>Department of Ecology and Biogeography, Faculty of Biological and Veterinary Sciences,  
115 Nicolaus Copernicus University, Lwowska 1, 87-100, Toruń, Poland. <sup>67</sup>Centre for Climate Change Research,  
116 Nicolaus Copernicus University, ul. Lwowska 1, 87-100 Toruń, Poland. <sup>68</sup>Instituto Nacional de Innovación  
117 Agraria, Programa Nacional de Investigación Forestal, Hualahoyo km 8, Huancayo, Perú. <sup>69</sup>Department of  
118 Natural Resource Management, South Dakota State University, Brookings, SD, USA. <sup>70</sup>Department of  
119 Geography, Swansea University, Singleton Park, Swansea SA2 8PP, UK. <sup>71</sup>Department of Soils and Rural  
120 Engineering, Faculty of Agronomy and Animal Science, Federal University of Mato Grosso - UFMT, Cuiabá,  
121 Mato Grosso, Brazil. <sup>72</sup>Department of Animal Science, Federal University of Paraná, Curitiba - PR, 80035-050,  
122 Brazil. <sup>73</sup>CITAB - Department of Forestry Sciences and Landscape (CIFAP), University of Trás-os-Montes and  
123 Alto Douro, Quinta de Prados, 5000-801, Vila Real, Portugal. <sup>74</sup>Plant Ecology, Institute of Integrative Biology,  
124 D-USYS, ETH-Zürich, 8006 Zürich, Switzerland. <sup>75</sup>Institute for Plant Sciences (IPS), University of Bern, 3013  
125 Bern, Switzerland. <sup>76</sup>Department of Geography, Indiana University, Bloomington, IN, USA. <sup>77</sup>Escuela de  
126 Ciencias Biológicas, Universidad Pedagógica y Tecnológica de Colombia (UPTC), Avenida Central del Norte  
127 39-115, Tunja, Colombia. <sup>78</sup>Facultad de Ciencias Naturales, Universidad Autónoma de Querétaro, Av. de las  
128 Ciencias s/n Juriquilla, CP 76230, Querétaro, Querétaro, Mexico. <sup>79</sup>Graduate Program in Ecology and  
129 Conservation, Federal University of Sergipe – UFS, São Cristóvão, Sergipe, Brazil. <sup>80</sup>Brazilian Agricultural  
130 Research Corporation (Embrapa), Embrapa Amazônia Ocidental, 69010-970, Manaus, AM, Brazil. <sup>81</sup>IHE Delft,  
131 P.O. Box 3015, 2601 DA Delft, The Netherlands. <sup>82</sup>VU University Amsterdam, De Boelelaan 1085, 1081 HV  
132 Amsterdam, The Netherlands. <sup>83</sup>Posgrado en Bosques y Conservación Ambiental, Universidad Nacional de  
133 Colombia, Medellín, Colombia. <sup>84</sup>Centro Universitario de Ciencias Biológicas y Agropecuarias, Universidad de  
134 Guadalajara, 45200 Zapopan, Jalisco, Mexico. <sup>85</sup>Centre for Functional Ecology, Department of Life Sciences,  
135 Faculty of Sciences and Technology, University of Coimbra, Portugal. <sup>86</sup>National Biobank of Thailand, National  
136 Science and Technology Development Agency, Pathum Thani 12120, Thailand. <sup>87</sup>Department of Biomaterials  
137 Science and Technology, School of Natural Resources, The Copperbelt University, P.O. Box 21692, Kitwe,  
138 Zambia. <sup>88</sup>Laboratory of Ecology and Dendrology, Federal Institute of Sergipe IFS, São Cristóvão, Sergipe,  
139 49100-000, Brazil. <sup>89</sup>Laboratory of Plant Ecology, Universidade do Vale do Rio dos Sinos (UNISINOS), Av.  
140 Unisinos, 950, Cristo Rei, São Leopoldo, RS, 93022-750, Brazil. <sup>90</sup>Instituto Biodiversa, Porto Alegre, RS, Brazil.  
141 <sup>91</sup>Nuclear Energy in Agriculture, University of São Paulo, Avenida Centenário, 303 - São Dimas, Piracicaba -  
142 SP, 13416-000, Brazil. <sup>92</sup>Department of Botany and Ecology, Institute of Biosciences, Federal University of

Mato Grosso (UFMT), Cuiabá, Mato Grosso, Brazil. <sup>93</sup>Tree-Ring Laboratory, Lamont-Doherty Earth Observatory, Columbia University, 61 Route 9W, Palisades, NY 10964, USA. <sup>94</sup>Carrera de Ingeniería Forestal, Universidad Autónoma Gabriel René Moreno, Facultad de Ciencias Agrícolas, Km 9 al Norte, El Vallecito, Santa Cruz, Bolivia. <sup>95</sup>BIOAPLIC, Departamento de Botánica, Universidade de Santiago de Compostela, EPSE, Campus Terra, 27002 Lugo, Spain. <sup>96</sup>El Colegio de la Frontera Sur, Departamento de Sociedad y Cultura, San Cristóbal de Las Casas, Chiapas, Mexico. <sup>97</sup>Escuela Profesional de Ingeniería Forestal de la Universidad Nacional Toribio Rodríguez de Mendoza de Amazonas, Calle Higos Urco N° 342, Chachapoyas, Perú. <sup>98</sup>Laboratorio de Dendrocronología, Carrera de Ingeniería Forestal, Universidad Nacional de Loja, 110101 Loja, Ecuador. <sup>99</sup>Department of Environmental Studies, University of California Santa Cruz, Santa Cruz, California, USA. <sup>100</sup>Facultad de Ciencias Agrarias, Universidad del Cauca, Popayán, Colombia. <sup>101</sup>Héméra Centro de Observación de la Tierra, Escuela de Ingeniería Forestal, Facultad de Ciencias, Universidad Mayor, Santiago, Chile. <sup>102</sup>Centro de Apoio Técnico à Execução Ambiental, Ministério Público do Estado de Mato Grosso. <sup>103</sup>Instituto Nacional de Investigaciones Forestales, Agrícolas y Pecuarias (INIFAP), Centro de Investigación Regional Pacífico Centro - Campo Experimental, Centro Altos de Jalisco, Jalisco, Mexico. <sup>104</sup>DendrOlavide, Departamento de Sistemas Físicos, Químicos y Naturales, Universidad Pablo de Olavide, Crta. de Utrera km. 1, 41013, Sevilla, Spain. <sup>105</sup>Forest Ecology & Forest Management Group, Wageningen University, Wageningen, The Netherlands. <sup>106</sup>Forest and Nature Management, Van Hall Larenstein University of Applied Sciences, Velp, The Netherlands. <sup>107</sup>National Institute for Amazon Research, Av. André Araújo, 2.936, Petrópolis, CEP 69.067-375, Manaus, Amazonas, Brazil. <sup>108</sup>Environmental Studies, Department of Biology, Institute of Ecology and Evolution, University of Oregon, 97403-5223, Eugene, OR, USA. <sup>109</sup>Department of Earth Sciences, Freie Universität Berlin, Malteserstraße 74-100, 12249 Berlin, Germany. <sup>110</sup>Instituto de Ecología Aplicada, Universidad Autónoma de Tamaulipas, Av. División del Golfo No 356, Col. Libertad, CP 87020, Ciudad Victoria, Tamaulipas, Mexico. <sup>111</sup>Institute of Biodiversity and Forests, Federal University of Western Pará - UFOPA, Santarém - PA, 68040-255, Brazil. <sup>112</sup>Department of Earth and Environmental Systems, Indiana State University, Science 159E, Terre Haute, IN, 47809, USA. <sup>113</sup>Department of Geography, University of Alabama, Tuscaloosa, AL, USA. <sup>114</sup>Programa de Investigación de Ecología y Biodiversidad, Asociación ANDINUS, Calle Miguel Grau 370, Sicaya, Huancayo, Junín, Peru. <sup>115</sup>Department of Geography, Johannes Gutenberg University, Mainz, Germany. <sup>116</sup>Department of Environmental Science, Faculty of Science, Chulalongkorn University, Bangkok, 10330, Thailand. <sup>117</sup>Water Science and Technology for Sustainable Environment Research Unit, Chulalongkorn University, Bangkok, 10330, Thailand. <sup>118</sup>Instituto de Geografía, Universidad Nacional Autónoma de México, Ciudad Universitaria, Mexico City, 04510, Mexico. <sup>119</sup>Facultad de Ciencias Forestales, Colegio de Postgraduados, Montecillos, Texcoco, Mexico. <sup>120</sup>Instituto de Ciencias Agroalimentarias, Animales y Ambientales (ICA3), Universidad de O'Higgins, San Fernando, Chile. <sup>121</sup>Biological Sciences Department, California State University, San Marcos (CSUSM), CA, USA. <sup>122</sup>Faculty of Civil Engineering and Geosciences, Delft University of Technology, Delft, The Netherlands. <sup>123</sup>School of Teacher Training for Secondary Education Tilburg, Fontys University of Applied Sciences, Tilburg, The Netherlands. <sup>124</sup>Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, Santa Cruz, CA, USA. <sup>125</sup>School of Earth Sciences, Addis Ababa University, Addis Ababa, Ethiopia. <sup>126</sup>School of Biology and Ecology, University of Maine, Orono, ME 04469, USA.

## Abstract

Tropical forests and woodlands are key components of the global carbon and water cycles. Yet, how climate change affects these biogeochemical cycles is poorly understood because of scarce long-term observations of tropical tree growth. The recent rise in tropical tree-ring studies may help to fill this gap, but a large-scale quantitative analysis of their potential in global change research is missing.

We compiled a list of all tropical tree species known to form annual tree rings and built a network encompassing 492 tropical ring-width chronologies to evaluate the potential to generate insights on climate sensitivity of woody productivity and to build centuries-long reconstructions of climate variability. We assess chronology quality, length, and climatic representativeness and explore how these change along climatic gradients. Finally, we applied species-distribution modeling to identify regions with potential for tree-ring studies in ecological and climatic studies.

The number of tropical chronologies has rapidly increased, with ~400 added over the past two decades. Yet, tree-ring studies are biased towards high-elevation locations, with gaps in warmer and wetter climates, on the African continent, and for angiosperm species. The longest chronologies with strongest climate signals (i.e., synchronous growth variations among trees) are from cool regions. In wet regions, climate signals and precipitation sensitivity decrease. Most tropical regions harbor 5–15 (and up to 80) species with proven potential to generate chronologies. The potential for long climate reconstructions is particularly high in drier high elevation sites. Our findings support strategies to effectively expand tree-ring research in the tropics, by targeting specific species and regions. Tropical dendrochronology can importantly contribute to global change research by generating historical context of climate extremes, quantifying climate sensitivity of woody productivity and benchmarking vegetation models.

**Keywords:** Climate sensitivity, growth synchrony, pantropical tree growth, dendrochronology

## Introduction

Tropical forested ecosystems are key components of the global carbon and water cycles. Forests and woodlands in the tropics and subtropics (herein defined based on latitude between 30 °S and 30 °N; Corlett, 2013) contribute substantially to carbon sequestration (Pan et al., 2024), drive the interannual variability in the land carbon sink (Fan et al., 2019; Friedlingstein et al., 2020), and generate rainfall through high transpiration rates at regional and sub-continental levels (Staal et al., 2018). These contributions depend critically on the productivity of tropical vegetation (Poulter et al., 2014), and is modulated by geographic location and climate variability (Wang et al., 2016; Humphrey et al., 2018).

Insights on this climate sensitivity of tropical vegetation are limited by the paucity of ecological field data and long-term climate data in the tropics (Menne et al., 2012). Compared to temperate and boreal zones, the density of meteorological stations, flux towers, permanent sampling plots, climate manipulation experiments, and tree-ring studies is much lower in the tropics (Babst et al., 2021; Crowther et al., 2015; Villarreal and Vargas, 2021; Zhao et al., 2019). In addition, the duration of tropical studies on climate-productivity relations is often much shorter compared to that of studies in other climate zones (Pastorello et al., 2020). This data scarcity limits options to calibrate and benchmark Dynamic Global Vegetation Models (DGVMs) for tropical ecosystems (Zuidema et al., 2018; Xu et al., 2024). In addition, the low density of meteorological stations and short duration of instrumental climate across much of the tropics impairs the accuracy of gridded climate data products and prevents climatologists to put current climate extremes into a long-term perspective (Menne et al., 2012).

Tree-ring analyses can contribute to filling these data gaps by providing long-term, annually resolved datasets on species-level woody productivity and by allowing climate reconstructions. Extra-tropical tree-ring studies have been used to reveal shifts in climate-growth relationships (Babst et al., 2019), to benchmark DGVMs (Barichivich et al., 2021), and to perform distribution-wide analyses of climate sensitivity (Klesse et al., 2020). At tropical and subtropical latitudes, the annual formation of tree rings has now been proven for approximately 500 tree species (Brienen et al., 2016; Locosselli et al., 2020; Schöngart et al., 2017), and almost 500 tree-ring chronologies – time series of common tree-

ring width patterns within a population – have been published (Zuidema et al., 2022). These recent advances exemplify the large potential of tropical dendrochronology to reconstruct past climate variability, to improve our understanding of the effects of climate fluctuation on tropical woody productivity, and offer opportunities to benchmark and calibrate remote sensing products and DGVM output (Babst et al., 2014; Jeong et al., 2020). Yet, a pantropical assessment of the potential of tropical tree-ring data for global change studies is missing to date.

Here we leverage quantitatively review the quality, length, and climatic representativeness of tropical tree-ring chronologies, and by assessing the magnitude of climate sensitivity of tropical tree growth. We also evaluate the potential of current tropical dendrochronology to contribute to our understanding of climate sensitivity of woody productivity and to reconstruct climate variability over the past centuries. We conclude by identifying opportunities and difficulties for tropical tree-ring studies in hitherto underrepresented regions.

We address the following questions: (1) To what extent are tropical tree-ring chronologies geographically and climatically representative of tropical wooded ecosystems? (2) What is the strength of the climate sensitivity of tropical tree growth and does this sensitivity depend on mean climate and on the quality of meteorological data? (3) Which climatic conditions offer the best opportunities for climate reconstruction based on tropical tree-ring chronologies? (4) What is the pantropical distribution of the potential for tree-ring studies and climate reconstruction?

## Methods

### *Assembling tree-ring chronologies*

Our review is based on chronologies included in a recently established tropical tree-ring network ([www.tropicaltreeringnetwork.org](http://www.tropicaltreeringnetwork.org)). We compiled published ring-width chronologies from naturally regenerating tree populations in tropical and subtropical latitudes (30°N to 30°S). Thus, we included chronologies based on geographical limits of the tropics in a broader sense and we did not select sites exclusively within the climatological definition of the tropics or exclusively in tropical biomes. This implies that several of the genera we include are commonly present in temperate regions (*Abies*, *Picea*, *Larix*, etc.).

We used raw ring-width data from two sources: (1) 242 chronologies from the International Tree-Ring Data Bank (ITRDB, <https://www.ncdc.noaa.gov/data-access/paleoclimatology-data/datasets/tree-ring>) and (2) 250 chronologies from individual contributors. We thus included a total of 492 chronologies in the quantitative analyses, based on measurements of 10,936 individual trees and 20,915 radial series from 139 species (Appendix 1). Short or statistically “unsafe” chronologies are usually removed from studies that combine datasets. Yet, as our intent was to provide the widest possible perspective on tropical tree-ring studies, we included short chronologies (here, the minimum was 16 years) and those based on only a few individuals (minimum here = 4 trees). In our final dataset, 82.7% of the chronologies covered at least 35 years and included a minimum of 10 individuals.

Additionally, to explore the broader potential of using tree rings in the tropics, we also compiled a species list from studies that use tropical tree rings for other purposes than building chronologies. Several studies exist that have applied tree-ring data without building chronologies to answer questions related to forest ecology (Van der Sleen et al., 2015a), dynamics (Godoy-Veiga et al., 2018), isotope-based rainfall reconstructions (Woodborne et al., 2015), tree physiology (Garcia et al., 2022; Loader et al., 2011), anthropogenic disturbances (Caetano-Andrade et al., 2020), or forest management (Groenendijk et al., 2017). Many of these studies did not focus on climate-growth related questions *per se*, and hence did not seek to build chronologies. A large part of these studies therefore do provide proof of annuality of ring formation using radiocarbon dating, phenology observations, dendrometers,



cambium wounds or ring counts in plantations of known age (Brienen et al., 2016). Although not producing chronologies, these studies reflect the potential of a broader set of species for tree-ring studies. This species list was created by complementing species lists from review papers (e.g., Brienen et al., 2016; Locosselli et al., 2020; Quesada-Román et al., 2022; Portal-Cahuana et al., 2023) with a literature search using the search terms “tropical tree rings” and “tropical dendrochronology”. This was not intended to be an exhaustive search and we acknowledge that we have missed non-English publications (e.g., in Spanish; see Portal-Cahuana et al., 2023), but we believe it does provide a good representation of tropical tree species with potential to be used in global change studies.

### *Chronology construction*

To allow for the comparison of results across sites, we redeveloped chronologies from the raw ring-width data. We applied a flexible 30-year cubic-spline detrending method (with a 50% frequency cut-off) to all individual raw ring-width series to remove low-frequency signals (i.e., trends in the growth series at frequencies of decades or centuries) and emphasizes the interannual variation in ring widths that was our main interest (Hughes et al., 2010). We developed mean site chronologies of a dimensionless ring-width index (RWI) from the detrended series using a bi-weight robust mean. We then used the most recent 50 years of each chronology (or less in case of short chronologies) for further analyses (e.g., of inter-series correlations ( $R_{bar}$ ) or monthly climate correlations, etc.). We followed the Schulman convention in the development of all tree-ring series from the Southern Hemisphere (Schulman, 1956), except for chronologies in the Brazilian Caatinga biome, where the rainy season coincides with the Northern Hemisphere’s growing season (Zuidema et al., 2022). We conducted detrending and chronology building using the dplR package (Bunn, 2008; Bunn et al., 2023) in R (R Core Team, 2023). We note that chronologies and further analyses presented here may slightly differ from those of the published chronologies, because of differences in detrending procedure and period covered in the study.

### *Climate data and distance to climate stations*

We used two types of gridded climate data: (1) Worldclim version 2 (worldclim.org) to obtain mean annual precipitation (MAP) and temperature (MAT) between 1970-2000 at 1-km spatial resolution, and (2) CRU TS4.02 to obtain monthly maximum temperature ( $T_{\max}$ ) and monthly precipitation data for the most recent 50 years of each chronology, at a coarser spatial resolution of 0.5°. The quality of gridded climate data is a function of the local density of meteorological stations. A low density likely weakens the accuracy of the interpolation and, with that, also weakens the correlation between climate variability and ring width. To evaluate the impact of meteorological station density on the magnitude of climate-growth relationships we calculated the mean distance between each chronology location and its five nearest meteorological stations. We do not account for elevation differences between the climate stations and the study location as we assume that these changes will affect the absolute values of temperature and precipitation, but not so much the interannual variation in the data. We obtained locations of all stations within the (sub-)tropics from the Global Historical Climatology Network (Menne et al., 2012) and selected stations with at least 25 years of data. We then calculated the distance of each study location to the nearest stations per corresponding continent using 'Vincenty' (ellipsoid) great-circle distance estimations with the *distm* function of the *geosphere* R package (Hijmans, 2021). We then identified the five stations closest to each of the tree-ring sites and calculated their mean distance to the site. Finally, we checked for a statistical relationship between this mean distance and the strength of the climate-growth correlations obtained from the tree-ring data.

### *Climatic representativeness*

We evaluated the climatic representativeness of our network (research question 1) by comparing the MAP and MAT distribution of the chronologies from our tree-ring study sites against MAP and MAT distributions from the entire tropical land areas supporting woody vegetation (i.e., with >10% tree cover). We used tree-cover data from the MODIS-derived MOD44B product (version 6; <https://lpdaac.usgs.gov/products/mod44bv006/>) to mask out areas with <10% tree cover from the Worldclim 2 data. We then calculated continent-level relative MAP and MAT distributions (i.e., the kernel density estimates) of the tropical land area with woody vegetation. To obtain corresponding

density estimations for precipitation and temperature of the tree-ring network, we calculated these density estimates for the sites in the network using the same maximum, minimum and bin-size values as for the entire forested area. By dividing the scaled distribution values of the network by those of the forested land area per continent, we obtained a measure of biases in how the network represents the climatic envelope: values above 1 indicate “overrepresentation”, those below 1 “underrepresentation”. Thus, if a certain MAT bin contains 5% of the values of the tree-ring network but represents only 2% of the values of all tropical land area, this ratio would be 2.5 and an indication that the network is biased towards overrepresenting this MAT range. Note that this measure of representativeness tells how well a part of the climatic envelope is covered in relative terms but does not provide information on the absolute density of tree-rings sites (per unit area).

#### *Climate correlations and growth synchronicity*

To estimate the strength of climate signals embedded in tropical tree-ring chronologies (research question 2), we calculated simple Pearson’s correlation coefficients between Ring Width Index (RWI) and monthly  $T_{\max}$  and precipitation data for a 24-month period that includes the year of growth (e.g., the peak of the growing season) and the previous year. For Northern Hemisphere and *Caatinga* sites, this period includes the full calendar year prior to the year of ring formation, plus the full calendar year during which the ring was formed. For Southern Hemisphere sites, it contains the 12-months July-June period preceding the onset of ring formation, and the following 12 months during which the ring was formed. For each climatic variable we then selected the highest monthly correlation coefficient of the 24 correlations. We do not present a more exhaustive analysis of climate responses, because our aim is to provide an indication of the maximum climate sensitivity of growth, which is relevant for climate reconstructions and studies quantifying climate effects on tree growth.

The quality of tree-ring chronologies is commonly indicated by a measure of growth synchronicity. As a measure of this synchronicity, we used the dendrochronological statistic 'Rbar': the mean correlation between all the ring-width series within a population (Hughes et al., 2010). Low Rbar-values may indicate poor dating quality, weak effects of common environmental drivers on

growth, or both. We did not set a minimum Rbar criterion for this review, because we were interested in presenting the full scope of characteristics of tropical tree-ring chronologies.

#### *Statistical analyses: climate correlations, Rbar, chronology length*

To evaluate factors driving the degree of climate sensitivity (question 2), we performed a multiple regression of the strongest monthly climate correlation, which we expected to increase with site aridity (i.e., higher MAT and lower MAP) and to be reduced when meteorological stations are scarce (i.e., higher distance to the five nearest stations). A second regression assessed factors influencing the growth synchronicity (Rbar) of the chronologies, which needs to be high for climate reconstructions (question 3). Rbar is expected to be higher at low MAT (i.e., in colder, mountainous climates), at low MAP (i.e., where precipitation exerts a strong common limitation to the growth of co-occurring trees) and to increase with the strongest monthly correlation (i.e., a strong synchronicity is likely driven by a common climatic limitation). The third multiple regression analysis evaluated how the potential for climate reconstruction (question 3) in terms of chronology length (tree longevity) is associated with mean site climate. We expected longer chronologies at lower MAT and lower MAP, thus at sites at cold and/or dry limits of the species where its growth is restricted by climate.

For all multiple regressions, we used the “leaps” algorithm for model selection, an all-subset model comparison that is more robust than stepwise methods (Furnival and Wilson, 1974). We checked the Variance Inflation Factors (VIFs) of all models and found these to be lower than 2. Analyses were conducted in R using packages *leaps* (Lumley, 2020) and *bestglm* (McLeod et al., 2020).

#### *Species distribution models*

To identify geographic regions that harbor tree-ring forming species, and areas that are particularly suitable for building tropical tree-ring chronologies, we ran species distribution models (Maxent) based on occurrence data from Global Biodiversity Information Facility – GBIF – extracted using the *rgbif* R package (Chamberlain et al., 2023) – and Worldclim2 climate data. We ran these distribution models for the larger set of ring-forming species (i.e., the 513 species extracted from the literature) and

the set of species for which chronologies have been built (i.e., the 139 species of which chronologies are included in the network). For each species, we ran distribution models only for the continent where the species occurs naturally. To reduce the effect of a high local density of occurrence data on model fits, we thinned observation data to a maximum of one observation per 0.5° grid cell. Additionally, we only conducted distribution models for species that had a minimum of 10 grid cells with observations, that is, for 450 (out of 513) species for the large set of ring-forming species, and for 122 (out of 139) species for the chronology-bearing species list. We summed the probability of occurrence of all species per cell to produce a map with estimates of the total number of ring-forming species and that of species with dendrochronological potential per grid cell.

We used a similar approach to quantify the potential length of the chronologies that can be built in each cell, using the species list for which chronologies exist. For grid cells with a probable species occurrence ( $P > 0.5$ ), we calculated the maximum length (90% of the maximum chronology length) of the chronologies for that species and averaged this across all species occurring in that cell. This yielded a map with estimates of the maximum chronology-length per grid cell, a proxy for the potential for climate reconstruction.


We note that values in the resulting maps should be interpreted cautiously because (1) the 0.5 probability cut-off for the species distribution model is rather arbitrary and probably does not represent well marginal sites where the most growth-limiting conditions for a species occur and thus where the longest and most climate-sensitive chronologies for that species can be built, (2) species occurrence in a certain grid cell does not necessarily imply its suitability for chronology construction at that local climate (Baker et al., 2017), (3) the quality and density of species-occurrence data may vary between continents (Meyer et al., 2016), which affects distribution modelling (Beck et al., 2014), (4) the maximum lifespan of a species changes along environmental gradients (Locosselli et al., 2020), which was not accounted for, and (5) distribution modelling based on climate data does not account for changes in species occurrence due to soil conditions (Zuquim et al., 2020) and effects of forest degradation on tree age (Feeley et al., 2012). Thus, the resulting maps provide a first representation of the geographic distribution of the potential for chronology building and climate constructions.

## Results and Discussion

### Geographic and taxonomic distribution of tropical tree-ring chronologies

The geographic distribution of tropical tree-ring chronologies is uneven (Fig 1A). Among the 492 chronologies included in this review, less than 10% are located on the African continent, whereas more than half are from the Americas. Dendrochronology studies are also biased towards the Northern hemisphere: most studies were performed in the Northern Hemisphere (~1.5 times more), while a larger proportion of land between 30°S and 30°N is in the Southern hemisphere (134% more land than the Northern Hemisphere). These geographic biases coincide with those of the global tree-ring databank (ITRDB), in which extra-tropical and Northern Hemisphere chronologies dominate (Zhao et al., 2019). These biases possibly arise from the nature of research funding (concentrated in the Global North) or the role of site selection close to home institutions. The number of tropical chronologies has increased rapidly over the past decades. Since the year 2000, close to 400 chronologies have been added at a rather steady rate of around 20 per year (Fig 1B); equivalent to an annual addition 8.3% for tropical chronologies to the ITRDB. About two thirds of the new additions (217 out of 378) are from lowland regions (<1500m a.s.l.) and the growth in the number of chronologies has been particularly steep in the Americas.

In total, our literature search identified 513 species belonging to 287 genera and 72 plant families. Our network encompasses chronologies for 139 tree species (Fig 1B-D), belonging to 88 genera and 35 plant families (Appendix 2). Our network thus includes about one quarter of the (sub)tropical tree species with known annual ring formation (139 out of a total of 513). The lower number of species covered in our network partly reflects chronologies that have not (yet) been included, but a much larger share represents species for which tree-ring studies did not include the construction of chronologies. The difference in species number also demonstrate the high potential to increase the number of species and chronologies for tropical trees. On the other hand, the rather high proportion of tree-ring studies without chronology building also likely reflects difficulties of constructing ring-width chronologies in the tropics. Ring boundaries are diverse and can be difficult to identify in tropical species (Fichtler and Worbes, 2010; Brien et al., 2016).



**Figure 1. Pantropical distribution and recent increase in the number of tropical tree-ring chronologies.** (A) Geographical distribution of 492 tropical tree-ring chronologies from angiosperms (black squares) and gymnosperms (triangles) included in this review. The map background is MODIS-based tree cover percentage. (B-C) Temporal change in the number of chronologies per continent (B) and the number of species for which chronologies have been constructed (C), grouped by gymnosperms and angiosperms and separated in columns per continent: Americas, Africa, Asia.

Tropical trees can also show irregular growth (e.g., fluted stems, buttress roots), form false rings, and partial cambial dormancy (Boninsegna et al., 2009). These difficulties increase the chances of accumulated measurement errors hampering crossdating further into the past (Black et al., 2016) and it is recommended to work with a large number of individuals, with cross-sectional discs or multiple radii collected per tree (Brienen & Zuidema, 2005; Groenendijk et al., 2014) and sampling above parts of the stem with irregular growth (Granato-Souza et al., 2019). Another difficulty in constructing chronologies may be the lack of a strong limiting climatic factor on growth (e.g., a

pronounced dry season) (Groenendijk et al., 2014). Yet, this does not apply everywhere, because chronologies have been successfully built in a-seasonal, hyper-wet tropical forests (Giraldo et al., 2023). Additional barriers for chronology building include the difficulty to assess remote field areas for collection of additional samples, the low number of tree-ring researchers and laboratories, and difficulties to publish in English (Portal-Cahuana et al., 2023). National and international training programs can strongly boost tree-rings studies in a country (Portal-Cahuana et al., 2023) and are needed, especially in Africa where tree-ring analyses has a large but unfulfilled potential (Gebrekirstos et al., 2014). International projects should also foresee building laboratory and analytical infrastructure (increment borers, microtomes, polishing machines, measuring tables, scanners, software, etc.) and establish long-term collaborations and training to overcome these barriers in tropical dendrochronology and ensure a lasting legacy.

The taxonomic distribution of tree species with existing chronologies is strongly skewed, with 44% of chronologies stemming from pines (Pinaceae) and with gymnosperm taxa representing 55% of all chronologies. Yet, the dominance of gymnosperms in tropical chronologies is less pronounced than in the global ITRDB, where they represent >80% chronologies (Zhao et al., 2019). Among chronologies from angiosperm species, taxonomic diversity is particularly high, with 96 species represented belonging to 76 genera. The top-10 angiosperm families include Fabaceae (55 chronologies) and the typical tropical (lowland) families Meliaceae (53) and Bignoniaceae (9). During the past two decades, the number of species for which chronologies have been constructed increased by about 100 (Fig 1C). This growth has been particularly rapid in the Americas since 2010, where the number of species represented by chronologies has reached 100. The recent increase in the number of studied tree species producing rings and the number of chronologies built demonstrates the large and underexploited potential of dendrochronology in the tropics. This potential may even be larger, as many studies in the tropics are not published in peer-reviewed journals (remaining in grey literature, such as theses and reports; Portal-Cahuana et al., 2023), many studies are published in languages other than English (e.g., initial exploratory works in the 1930s and 1970s; Worbes, 2002), and many tropical chronologies are not added to global tree-ring networks. The rapid research advances in the tropical




Americas show no signs of leveling off. A likely factor responsible for this faster development of American dendrochronology is a higher number of tree-ring labs and trained professionals compared to Asia and Africa (Gebrekirstos et al., 2014; Pumijumnong, 2013). Yet, such potential is also likely to exist in Asia and Africa, given the abundance of ring-forming tree taxa and the availability of long-lived tree species (Gebrekirstos et al., 2014; Groenendijk et al., 2014; Pumijumnong, 2013), but the high diversity and abundance of dipterocarps that do not form rings may limit opportunities in the wet Asian tropics (especially in the ‘Indo-Malayan Realm’).

### **Climatic representativeness of tropical tree-ring chronologies: new frontiers**

Tropical tree-ring chronologies have been constructed across a wide MAP range, from <200 to >5000mm and these chronologies thus represent precipitation regimes of the tropical woody land area well. Arid climates (with MAP <500 mm), that cover >15% of the tropical woody land area, are also somewhat underrepresented with ~5% of the chronologies. Sites with more semi-arid and seasonal climates (500-1500 mm MAP) are better represented than wetter regions (>2000 mm MAP), with the latter covering only 8% of all chronologies (n=39 sites). The chronologies at wetter sites and recent studies in extremely wet regions (Giraldo et al., 2023) illustrate the prospects to conduct tree-ring studies at the wettest extremes of the tropical rainfall distribution.

In contrast to the fairly good representation of precipitation regimes in tree-ring chronologies, their distribution rather poorly follows that in temperature. Half of the tropical tree-ring studies have been conducted in montane ecosystems (>1500 m a.s.l.), particularly studies with a focus on (long-lived) gymnosperm species in Asia and the Americas. These areas with low-temperature climates (<15°C MAT), which cover just 2% of the tropical land area with woody vegetation, are strongly overrepresented (Fig 2A-C) in our network. Conversely, warm tropical lowland ecosystems with a MAT >25°C, which cover 88% of wooded land in the tropics, are represented by only 15% of chronologies.



**Figure 2. Climatic distribution and skewness of tropical tree-ring chronologies.** (A-C) Continental-level relative kernel-density estimates of the mean annual precipitation (MAP) and temperature (MAT) envelope covered by the tree-ring chronologies ('Chronologies') and the tropical land area with woody vegetation (pixels with >10% tree cover; 'Continent all'). Spatial distribution of over- and underrepresentation of MAP (B) and MAT (C) of tropical tree-ring chronologies. Underrepresentation (values < 1) implies that – for a given climatic condition – the proportion of sites in the tree-ring network is smaller than that of the forested area. Values >3 were set to 3 to improve clarity.

We projected the climatic “representativeness” covered by tropical dendrochronological studies spatially, to identify regions that require more research attention or that are well represented climatologically (Fig 2D-E). In terms of precipitation, regions with seasonally dry climates (Central America, Northeastern Brazil, the Sahel, large areas in India) are more prominently represented in our network while the wet tropics (Amazon, Central Africa, Southeast Asia) are underrepresented. In

terms of temperature, patterns are similar but show the prevalence of colder climates in the network. Regions that are overrepresented in terms of temperature include high-elevation areas in parts of Mexico and Asia, and – to a lesser extent – low-elevation subtropical regions in Southeast Brazil and subtropical China. We note that, independent of the climate-envelope coverage, the overall low number of tropical dendrochronology studies implies that the representation of tropical climates in tree-ring records is considerably lower (by a factor 5) than that of temperate and Mediterranean climates (Babst et al., 2019; Zhao et al., 2019). The interpretation of these maps should thus be cautious, especially for Africa, where the number of chronologies is low, and representation is poor across the full climatic gradients.


What does this climatic skewness in tropical chronologies imply for their use in global change studies? First, climatic over- and underrepresentation needs to be accounted for in regional or pantropical analyses by checking its influence on results or explicitly accounting for its influence using weighted statistical analyses (e.g., Zuidema et al., 2022). Second, published tropical tree-ring chronologies should be readily uploaded in databases such as the ITRDB, because this increases their representation in pantropical or global analyses. This also applies to chronologies with weak climate responses as this helps drawing a more complete picture of tropical and global tree responses to climatic variation. Additionally, ring-width data need to be archived correctly (e.g., TRiDaS format; Jansma et al., 2010), and supported by appropriate metadata (Zhao et al., 2019) and preferably also with ancillary tree-level data that currently not supported by the ITRDB (Rayback et al., 2020).

Third, our analyses of climatic representativeness provide guidance to identify priority regions for tree-ring sampling. Evidently, the highest priority is in collecting samples and building chronologies for African tree species, which are poorly represented. In addition, increased sampling efforts are needed in the wet and warm tropics, which cover a vast area but are poorly represented. Chronology building in the wet and warm tropics can be challenging, because interannual climatic variation is limited and common climate responses across trees can be concealed by the impacts of canopy dynamics (Giraldo et al., 2023; Groenendijk et al., 2014). Nevertheless, quantifying such subtle and variable responses of tree growth to climate variability in the wet and warm tropics is

crucial to understanding and predicting climate-change effects on forest productivity. Broadly, the priority regions indicated here are consistent with those based on a global analysis of the ITRDB (Zhao et al., 2019), but the considerably larger number of tropical chronologies included here allowed us to provide more specific recommendations for tropical climates and biomes.

#### **Potential of tropical tree-ring chronologies to reveal climate sensitivity of tree growth.**

The potential of tree-ring chronologies to provide insight into the sensitivity of tropical tree growth to climate fluctuations depends on the strength of climate-growth correlations and the degree to which temporal variations in tree-ring width are similar across trees in a population (the interseries correlations, i.e.,  $R_{\text{bar}}$ ). We analyzed both factors for all chronologies. Tree-ring width may strongly respond to climatic conditions during one or more months in a year. Strong correlations between ring width and climate variables such as precipitation or maximum temperatures ( $T_{\text{max}}$ ) during a particular month provide a first indication of the climate sensitivity of tree growth. We found these maximum correlations with monthly precipitation to be mostly positive (in 73.6% of chronologies) with a mean Pearson correlation coefficient of 0.39 (Fig 3A). By contrast, monthly correlations for  $T_{\text{max}}$  were mostly negative (66.5 %), and slightly stronger in magnitude ( $r=-0.40$ , Fig 3B). This distribution of prevailing positive precipitation and negative  $T_{\text{max}}$  sensitivities is consistent with those obtained in global and pantropical analyses (Babst et al., 2019; Zuidema et al., 2022). Across continents, stronger correlations were found for chronologies from the Americas, for both  $T_{\text{max}}$  (negative) and precipitation (positive) (Fig 1A).



**Figure 3. Climate sensitivity of tropical tree growth and its relation to mean climate and the quality of climate data.** Maximum climate sensitivity is shown as the highest correlation coefficient between the tree-ring chronologies and monthly climate (precipitation or  $T_{\max}$ ) for 24 months including the year of ring formation and the preceding year. Multiple regression models were constructed, with explanatory variables MAP, MAT and distance to the nearest meteorological station (as a proxy for climate data quality). Lines represent predicted relations of the regression model (Table 1): black lines represent the results including all correlations, grey lines the results of modeling with only the prevailing correlations (positive correlations for Precipitation and negative for  $T_{\max}$ ), full lines represent significant and dashed lines non-significant correlations ( $p < 0.05$ ).

For a subset of the prevailing correlations (positive for precipitation, negative for  $T_{\max}$ ) we tested associations of correlation coefficients with mean climate and with distance to the nearest meteorological station (viewed as a proxy for the representativeness and quality of climate data). Only a small portion of the variation in precipitation and temperature sensitivity across chronologies was explained by these variables (Table 1). Climate sensitivity of precipitation was stronger at drier sites, for both precipitation and temperature, suggesting a stronger water limitation during hot or dry years in more arid conditions. Negative effects of  $T_{\max}$  increased with decreasing MAP, but, contrary to our expectations reduced at warmer sites (i.e., less negative with increasing temperatures).

**Table 1. Associations of climate sensitivity with mean climate and the quality of climate data.**


The proxy for climate sensitivity used is the maximum correlation between ring-width index (RWI) and monthly precipitation or  $T_{\max}$ . Only prevailing correlations are used, so: positive correlations for precipitation and negative for  $T_{\max}$ . Shown are estimates (Est, unscaled) of coefficients, significance levels and relative importance values (RI) per explanatory variable. N=number of chronologies. \*= $p<0.05$ , \*\*= $p<0.01$ , \*\*\*= $p<0.001$ .

Explanatory variables	Precipitation sensitivity		$T_{\max}$ sensitivity	
	Est	RI	Est	RI
Intercept	0.43***		-0.52***	
MAP (mm)	-0.000017*	0.23	0.000030**	0.24
MAT (°C)	NS	-	0.0034***	0.3
Distance to nearest meteorological station (km)	-0.00029***	0.77	0.00049***	0.46
$R^2$	0.07		0.19	
Degrees of freedom	361		326	

Climate sensitivity was weaker for chronologies situated at longer distances from meteorological stations. An increase in the distance to the nearest station of 100 km reduced the absolute correlation coefficient by 0.03 for precipitation and by 0.05 for  $T_{\max}$ . Distances to the nearest station were smaller in South America, reflecting the higher density of stations. As a large share of tropical chronologies is obtained from sites at >100 km from the nearest station (~24%), this finding suggests that dendrochronological analyses underestimate the climate sensitivity of tropical tree growth. In addition, in mountainous areas elevational differences between tree-ring sites and climate stations may add additional uncertainties in the position of chronologies in our climate space based on interpolated data. Yet, a more detailed analysis of the effect of distance to climate station conducted for a smaller set of tropical chronologies (n=347) revealed limited sensitivity of climate correlations to proximity of climate stations (Zuidema et al., 2022).

The extent to which tree growth is synchronized depends on the degree to which climate exerts a dominant limitation on tree growth, compared to other factors such as canopy dynamics, disturbances, pests, and diseases. The  $R_{\text{bar}}$  of all series within a chronology varied strongly between chronologies, from -0.04 to 0.85, thus from no common growth signal to very strong synchronization among trees. The strength of growth synchronicity differed between continents and was considerably

lower for African chronologies (mean=0.30) compared to American and Asian chronologies (0.50-0.52; Fig 4). We further assessed to what extent the variability in growth synchronicity can be explained by mean climate (MAT and MAP) and by information on the maximum climate sensitivity (i.e., the maximum climate correlation). As expected, we found a higher Rbar (i.e., stronger synchronicity) at more arid (low MAP) and cooler (low MAT) sites. At those sites, years with low water availability or low temperatures may impose strong limitations to tree growth. We also found growth synchronicity to be stronger for chronologies that exhibited stronger correlations with monthly climate (Fig. 4).



**Figure 4. Growth synchronicity of tropical tree species as a function of mean climate and maximum climate sensitivity.** Growth synchronicity is calculated as the mean of all correlation coefficients between individual tree-ring series within a chronology (Rbar). Lines represent predicted effects of Mean Annual Precipitation (MAP), Mean Annual Temperature (MAT) and maximum

climate sensitivity (i.e., highest maximum monthly correlation) based on a multiple regression model (Table 1).


Our review of climate sensitivity shows that tropical tree-ring chronologies can be an important tool to assess climatic impacts on radial stem growth. This sensitivity can be evaluated at annual resolution for many species and sites. Another important virtue of tropical tree-ring chronologies is the ability to obtain tree growth data retrospectively and thus fill gaps in growth measurements in both space and time. We also found that the quality of some tropical tree-ring chronologies is rather poor, as they are based on few radial series or from few individuals only, resulting in low growth synchronicity and weak climate correlations. Part of the variability in growth synchrony and climate-signal strength in the chronologies from these limitations and from decision of individual researchers building chronologies (e.g., synchronization effort, number of trees in the dataset to select only the most sensitive, etc.). Such data limitations can be overcome by teaching, applying, and publishing best practices of (tropical) chronology construction (Black et al., 2016), but also by reporting on the failure to build chronologies (Aragão et al., 2022; Groenendijk et al., 2014). Assessing the quality of tree-ring chronologies is not straightforward, because low growth synchronicity can result from small sample sizes, poor quality of ring measurements and cross-dating, or from the lack of a common signal and complacent growth. No analyses tools are currently available to disentangle these causes in tree-ring datasets. Long-term annual monitoring of large permanent plots (Feeley et al., 2007) and studies using (automatic) high-resolution dendrometer measurements on species with varying strategies (Wagner et al., 2016) may help elucidating these causes.

An important limitation of the current set of tropical tree-ring chronologies is poor species replication. Among the 164 species included in the Network, 134 are represented by just 1-3 chronologies, and only 8 species are represented by 10 or more chronologies, mostly in montane climates. Proper replication is needed to assess and predict how climate sensitivity shifts across climate gradients within the distributional ranges of species (Babst et al., 2018; Klesse et al., 2020), and thus to assess its vulnerability to shifting climates (Heilman et al., 2022; Perret et al., 2024). At present, this is possible for only very few tropical tree species and genera (Aragão et al., 2022; Baker et al., 2015; Zuidema et al., 2020), with limited spatial replication and across limited climate ranges.



## Potential of tropical tree-ring chronologies for climate reconstruction

An obvious prerequisite for tree-ring based climate reconstruction is the length of the chronology. Chronology length varied from 16 to 1237 years (Fig 5). Median chronology length was 203 years, but it was substantially higher for gymnosperms (median length of 288 years) than angiosperms (131 years). About half ( $n=249$ ) of the chronologies had a length of  $>200$  years. Given that instrumental climate data in tropical climate zones are sparse and short, these longer chronologies provide opportunities for climate reconstruction, provided that their climate signal is strong. Successful climate reconstructions based on tropical tree-ring data have been conducted on all tropical continents (Buckley et al., 2010; Granato-Souza et al., 2019; Heinrich et al., 2008; Stahle et al., 2011; Therrell et al., 2006) and have been used to establish continental-scale drought atlases, as well as analyses of major drought events (Cook et al., 2010; Morales et al., 2020; Stahle et al., 2016).



**Figure 5. Potential for tree-ring based climate reconstruction and its relation to mean climate.** Shown is the longevity of all reviewed chronologies – an important prerequisite for climate reconstruction – for gymnosperms and angiosperms separately. Lines represent predicted effects of MAP and MAT on chronology length based on a multiple regression model (Table 2).

The exploration of additional proxies of climate information (e.g., stable isotopes, wood anatomy) have also shown promising results (Van der Sleen et al., 2015b; Wils et al., 2010). We tested the extent to which mean climate is associated with chronology length and whether this relationship differs between gymnosperms and angiosperms. We found chronologies to be longer in cooler

climates and for gymnosperms compared to angiosperms (Table 2). The temperature effects were quite strong: chronology length extended by almost 8 years for every degree decrease in MAT. While high-elevation sites and gymnosperm species dominated the subset of long tropical chronologies, about 29.5% of angiosperm chronologies from warm ( $>20^{\circ}\text{C}$ ) and moist ( $>1200\text{ mm yr}^{-1}$ ) tropical climate sites extended to  $>200$  years ( $n=26$  of 88). Some of these angiosperm chronologies have been used for climate reconstruction in lowland tropical forests, adding essential, century-scale climate records to the instrumental record (D'Arrigo et al., 2011; Granato-Souza et al., 2019; Lopez et al., 2017).

**Table 2. Results of multiple regression models relating chronology characteristics with mean climate and climate sensitivity.** Shown are estimates (Est, unscaled) of coefficients, significance levels and relative importance values (RI) per explanatory variable. Dashes (-) indicate variables not initially included in the model. N=number of chronologies.  $*=p<0.05$ ,  $**=p<0.01$ ,  $***=p<0.001$ .


Explanatory variables	Growth synchronicity (Rbar)		Chronology length (y)	
	Est	RI	Est	RI
Intercept	0.616***		295***	
MAT( $^{\circ}\text{C}$ )	-0.00427***	0.26	-7.78***	0.43
MAP (mm)	-0.0000621***	0.36	0.0191*	0.02
Highest $T_{\text{max}}$ correlation	-0.0730*	0.22	-	-
Highest P correlation	0.0649***	0.16	-	-
Gymnosperms	-	-	134***	-
$R^2$	0.19		0.25	
Degrees of freedom	479		493	

The rapid rise of tree-ring studies and chronology construction in the Americas reveals the large potential for developing the field. But this rapid development likely also reflects recent increase in opportunities, funding, and interest in this field within the Americas, which may be slower in other continents. Main barriers to rapid development of tropical dendrochronology include limited laboratory infrastructure, experience, and funding for tree-ring analysis. In addition, factors such as low tree abundance in the forest, high wood density, poor taxonomic knowledge, and difficulties to

anatomically identify ring boundaries may limit practical application of tree-ring studies on many tropical (angiosperm) tree species (e.g., Groenendijk et al., 2014). High-resolution X-ray CT scanning (De Mil et al., 2016) and pith-to-bark histological sections (Quintilhan et al., 2021) can aid in ring-boundary identification. Opportunities to overcome these barriers include capacity-building programs, North-to-Tropical skill transfer and support with laboratory equipment, intensified cross-continental tropical collaborations (e.g., in networks) and open science (access to data, data standards, scripts for analyses and publications).

### **Pantropical and continental-level gaps and opportunities for tree-ring analysis**

To identify opportunities for tropical dendrochronology, we produced global maps of estimated number of species with proven annual tree-ring formation, of the number of species with published chronologies and of age estimates based on species distribution models (Fig 6). The maps show that across large parts of the woody tropical land area >20 tree species are expected to occur with known annual ring formation (Fig 6A), and >5 species with potential for chronology building (Fig 6B). Yet, large continental differences on these projections exist, with overall a greater apparent potential in regions in the Americas and Asia compared to Africa. In some areas – South-Eastern Brazil, Mesoamerica, and Southern China – more than 15 species with potential for chronology construction are expected to occur. The geographic distribution of species in these maps is a direct function of the total number of studied species per continent, because species distribution models are produced at continental level. As a result, the values for the African continent are low and should be interpreted with caution. Yet, this map does provide a first guidance to select target areas for future dendrochronological studies.



**Figure 6. Pantropical distribution of tree species with potential for tree-ring studies, chronology building and climate reconstruction.** Distribution of the estimated number of species for which (A) annual tree-ring formation has been proven and (B) ring-width chronologies have been constructed. Values were obtained by adding the probability of occurrence of all species distribution models per  $0.5^\circ \times 0.5^\circ$  pixel. Areas with  $<10\%$  tree cover are masked (white). (B) Distribution of maximum chronology length for species with known potential for chronology building. Values are obtained using the maximum chronology length for species with likely occurrence per  $0.5^\circ \times 0.5^\circ$  pixel and averaging this across all species in a pixel.

We also evaluated the opportunity for centuries-long climate reconstruction using tropical tree-ring chronologies. We used the results of species distribution modeling to identify areas with the longest chronologies, based on species-specific maximum chronology lengths (Fig 6C). The resulting map provides a first indication of regions that are likely more suitable for constructing long chronologies. For a large part of the woody tropics, maximum chronology length is expected to be  $>200$  years, whereas in some areas – northern Mexico, Central India, and southern China – chronology length can be 500 years or more. The importance of climate reconstructions is particularly large in regions where meteorological data are scarce or cover short periods. This is particularly the case for Africa, where a large proportion of the continent lacks long-term instrumental climate data.

## Conclusions

How can tree-ring chronologies contribute to global change research in tropical ecosystems? We identify three primary opportunities. First, tropical tree-ring chronologies can generate much-needed historical climate data to understand past climate dynamics and provide necessary context for climate extremes. Tree-ring based climate reconstructions have been crucial in identifying historical megadroughts (Cook et al., 2022; Morales et al., 2020; Stahle et al., 2011), and putting recent drought events into a long historical perspective (Belmecheri et al., 2016; Morales et al., 2020; Williams et al., 2022). In addition, they can be used to evaluate the possible attribution of meteorological extremes to ongoing climate change (Williams et al., 2022). To realize this potential, the replication and length of tropical tree-ring chronologies need to be substantially increased.

Second, tropical tree-ring chronologies can help assess the sensitivity of woody net primary productivity (NPP) to climate fluctuations, adding a century-long perspective and complementing other approaches such as eddy covariance flux towers, permanent sampling plots, and remote sensing data. Understanding this sensitivity helps to quantify the woody component of NPP and to improve our understanding of the mechanisms driving climate-C-sink dynamics of tropical vegetation (Zuidema et al., 2018). This approach could be of particular interest to assess El Niño Southern Oscillation (ENSO) effects and drought/heat effects on the tropical carbon balance (Rifai et al., 2018). Tree-ring derived climate sensitivity may also help estimate the potential and risks of failure of forest-based natural climate solutions (Anderegg et al., 2020). This is particularly true as long-term carbon sequestration critically depends on wood formation, which can be affected by climate extremes (Anderegg et al., 2020).

Third, tree-ring chronologies can help improve the simulation of woody productivity in Dynamic Global Vegetation Models (DGVMs). DGVMs increasingly represent wood formation processes (Friend et al., 2022) and tree-ring derived data can be used to simulate individuals or cohorts (Fisher et al., 2018), for model benchmarking (Jeong et al., 2020; Xu et al., 2024) and for improving models by data assimilation in DGVMs (Barichivich et al., 2021). So far, tropical tree-ring data have hardly been used for these purposes. Yet, the potential contribution of tropical tree-ring data is large,

because of the relatively low density and duration of tropical sampling plots, flux towers and global change experiments.

## **Acknowledgments:**

Support and assistance: Smithsonian Tropical Research Institute-Panama, Sebastian Bernal (RaAS). Supervision: Helene Muller-Landau, S. Joseph Wright (RaAS). Fieldwork support: COOMFLONA - FLONA TAPAJÓS, Universidade do Oeste do Pará (BH), Sutó Company and Angel Chavez at Consultora Forestal Bosques e Industria (KPV), Logging company AMATA (DROR). Lab support: Rebecca Franklin, Guillermo Guada, Quirine Hakkaart, Annemarijn Nijmeijer and Peter van der Sleen (KPV). This work was carried out with the support of CAPES - Financing Code 001. All authors have no conflict of interest to declare.

## **Data availability**

A large share of the raw tree-ring data (99 chronologies) have previously been uploaded to the International Tree-Ring Data Bank – ITRDB ([www.ncei.noaa.gov/products/paleoclimatology/tree-ring](http://www.ncei.noaa.gov/products/paleoclimatology/tree-ring)) following Zuidema et al. 2022. We are processing the remaining datasets to also be uploaded to the ITRDB. Analysis scripts will be made available via the main author's Github page: <https://github.com/groenendijk>.

## **Funding**

AECID grant 11-CAP2-1730 (to JJC & HM); Agencia Nacional de Promoción Científica y Tecnológica, Argentina grants PICT 2014-2797 and PICT 2019-01336 (MEF); BBVA Foundation (JJC & HM); Belspo BRAIN grant BR/143/A3/HERBAXYLAREDD (HB); CAPES - Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (TLS & EAF), grants 88887.199858/2018-00 (GAP), 88887.495294/2020-00 (BH), CAPES/PDSE 15011/13-5 (MAP); CGIAR (MM); Conselho Nacional de Desenvolvimento Científico e Tecnológico CNPq (CRA, FG, CSL, FAL, ASR), grants 140849/2015-7 (AKM), 1009/4785031-2 (GC), 140277/2024-2 (DBCC), 311247/2021-0 (JS), 405923/2021-0 (MCS), 406062/2023-4 (MCS), 441811/2020-5 (JS), ENV grant FRG 0339638 (OD), PQ 313129/2022-3 (ACB); CONACYT Consejo Nacional de Ciencia y Tecnología México, a master fellowship (MILH); grant CB2016-283134 (JVD & LRCE), CONAFOR-CONACYT grant C01-234547 (JCP), CONAFOR-CONACYT grant CONAFOR-2014 (LRCE); CONCYTEC Peru & World Bank grant 043-2019-FONDECYT-BM-INC.INV (JGI), and FONDECYT grant BM-INC.INV 039-2019 (MEF); CONICET (FAR) and grant PIP-11220200102929CO (MEF); Copel Geração e Transmissão S.A. grant PD-06491-0405-2015 (AKM); CUOMO Foundation (MM); German Academic Exchange Service (DAAD) (MR & MI); Deutsche Forschungsgemeinschaft (DFG) (AA), grants BR 1895/15-1, BR 1895/23-1, and BR 1895/29-1 (AB); Dirección de Investigación de la Universidad Nacional de Loja (DPC); Dirección General de Asuntos del Personal Académico of the UNAM (Mexico) (RB); Estudios dendrocronológicos en las Sierras Madre Occidental, Oriental y del Sur de México, grant 38111-4251030012346 (ECO); FACEPE grant IBPG-1418-5.00/21 (DBCC); FAPEAM

grant 01.02.016301.02630/2022-76 (JS); FAPEMAT (FAL); FAPEMIG grant APQ-01544-22 (ACB), grant APQ-02541-14 (GAP); FAPESC grant 2019TR65 (TABF); Fundação de Amparo à Pesquisa do Estado de São Paulo - FAPESP grants 2012/50457-4 (GC), 2009/53951-7 (MTF), 2018/01847-0 (PG), 2018/07632-6 (MGV), 2018/22914-8 (DROR & NB), 2019/08783-0 (GML), 2019/09813-0 (MGV), 2020/04608-7 (DROR), 2019/26350-4 (NB), FAPESP-NERC grant 2018/50080-4 (GC); FAPITEC (CSL & ASR); FAPITEC/SE/FUNTEC grant 01/2011 (MAP); FCT - Portuguese Foundation for Science and Technology grant UIDB/04033/2020 (JLPCL); Fulbright Fellowship (BJE); German Research Council (MM); HELVETAS Swiss Intercooperation (MEF); IAI-SGP-CRA grant 2047 (JVD); IFS grant D/5466-1 (JHS & JN); Inter-American Institute for Global Change Research IAI (FAR); ITTO Fellowship Award grant 046/12S (EJRR); Lamont Climate Center (BMB); Mahidol University grant FRB660042/0185 (NP); María Zambrano postdoctoral research program MZ2021 (RaAS); Minciencias grant 1118-714-51372 (IRNM); Ministerio Ciencia grant TED2021-129770B-C22 (RSS); National Geographic Global Exploration Fund grant GEFNE80-13 (IR); National Natural Science Foundation of China grant 31870591 (PF); National Research Council of Thailand (NRCT) grant N42A660392 (PT); National Science Foundation - NSF (BMB), grant AGS-1501321 (DGS & GAP), NSF CREST grant 0833211 (KSF); IBN-9801287 (AJL); AGS-2102888 (JM); AGS-2102938 (GLH); GER-9553623 (BJE); NSF Postdoctoral Fellowship (BJE); NSF-FAPESP PIRE grant 2017/50085-3 (CF, MTF, GC, GML, MGV & DROR); NSF-FAPESP PIRE grant 2019/27110-7 (CF); NUFFIC (JHS & JN); NUFFIC-NICHE (EM); PROCAD-AM grant 88887.625854/2021-00 (BH); Agencia Estatal de Investigación – Spain, Proyectos de generacion de conocimiento grant PID2021-123675OB-C44 (RSS); Schlumberger Foundation (JHS & JN); Short-term fellowship from the Smithsonian Tropical Research Institute (RaAS); Sigma Xi (AJL); Spanish Agency for International Development Cooperation (KPV); Thailand Science Research and Innovation Fund Chulalongkorn University (PT); The Copperbelt University (JHS & JN); Instituto Nacional de Ciência e Tecnologia em Fisiologia de Plantas em Condições de Estresse (INCT Fisiologia do Estresse) grant 406455/2022-8 (FAL); UK NERC grant NE/K01353X/1 (EG); UKRI grant EP/X025098/1 (NJL), and NE/B501504 (NJL); UNAM-PAPIIT grant IN110223 (LVS); Universidad Autónoma Agraria Antonio Narro (ECO); World Wildlife Fund (WWF) (KPV); Xunta de Galicia grant ED431C 2023/19 and ED481D 2023/012 (GPL)

838 **Author contributions:**

839 Conceptualization: PG, PAZ, FB, VT, ZXF, DGS, GML, MM, SP, NP; Data contribution: all co-  
840 authors except FB; Data analyses: PG, PAZ, FB; Writing – original draft: PG and PAZ, with important  
841 contributions from FB, VT, ZXF, DGS, GML, MM, SP, NP; Writing – review & editing: all  
842 coauthors.



**Table 1. Associations of climate sensitivity with mean climate and the quality of climate data.**

The proxy for climate sensitivity used is the maximum correlation between ring-width index (RWI) and monthly precipitation or  $T_{\max}$ . Only prevailing correlations are used, so: positive correlations for precipitation and negative for  $T_{\max}$ . Shown are estimates (Est, unscaled) of coefficients, significance levels and relative importance values (RI) per explanatory variable. N=number of chronologies. \*= $p < 0.05$ , \*\*= $p < 0.01$ , \*\*\*= $p < 0.001$ .

Explanatory variables	Precipitation sensitivity		$T_{\max}$ sensitivity	
	Est	RI	Est	RI
Intercept	0.43***		-0.52***	
MAP (mm)	-0.000017*	0.23	0.000030**	0.24
MAT (°C)	NS	-	0.0034***	0.3
Distance to nearest meteorological station (km)	-0.00029***	0.77	0.00049***	0.46
$R^2$	0.07		0.19	
Degrees of freedom	361		326	

**Table 2. Results of multiple regression models relating chronology characteristics with mean climate and climate sensitivity.** Shown are estimates (Est, unscaled) of coefficients, significance levels and relative importance values (RI) per explanatory variable. Dashes (-) indicate variables not initially included in the model. N=number of chronologies. \*= $p < 0.05$ , \*\*= $p < 0.01$ , \*\*\*= $p < 0.001$ .

Explanatory variables	Growth synchronicity (Rbar)		Chronology length (y)	
	Est	RI	Est	RI
Intercept	0.616***		295***	
MAT(°C)	-0.00427***	0.26	-7.78***	0.43
MAP (mm)	-0.0000621***	0.36	0.0191*	0.02
Highest $T_{\max}$ correlation	-0.0730*	0.22	-	-
Highest P correlation	0.0649***	0.16	-	-
Gymnosperms	-	-	134***	-
$R^2$	0.19		0.25	
Degrees of freedom	479		493	

855    **Appendices**

856

857    Appendix 1 and 2 – Separate PDF

- Anderegg, W.R.L., Trugman, A.T., Badgley, G., Anderson, C.M., Bartuska, A., Ciais, P., Cullenward, D., Field, C.B., Freeman, J., Goetz, S.J., Hicke, J.A., Huntzinger, D., Jackson, R.B., Nickerson, J., Pacala, S., Randerson, J.T., 2020. Climate-driven risks to the climate mitigation potential of forests. *Science* 368. <https://doi.org/10.1126/science.aaz7005>
- Aragão, J.R.V., Zuidema, P.A., Groenendijk, P., 2022. Climate-growth relations of congeneric tree species vary across a tropical vegetation gradient in Brazil. *Dendrochronologia* 71, 125913. <https://doi.org/10.1016/j.dendro.2021.125913>
- Babst, F., Alexander, M.R., Szejnér, P., Bouriaud, O., Klesse, S., Roden, J., Ciais, P., Poulter, B., Frank, D., Moore, D.J.P., Trouet, V., 2014. A tree-ring perspective on the terrestrial carbon cycle. *Oecologia* 176, 307–322. <https://doi.org/10.1007/s00442-014-3031-6>
- Babst, F., Bodesheim, P., Charney, N., Friend, A.D., Girardin, M.P., Klesse, S., Moore, D.J.P., Seftigen, K., Björklund, J., Bouriaud, O., Dawson, A., DeRose, R.J., Dietze, M.C., Eckes, A.H., Enquist, B., Frank, D.C., Mahecha, M.D., Poulter, B., Record, S., Trouet, V., Turton, R.H., Zhang, Z., Evans, M.E.K., 2018. When tree rings go global: Challenges and opportunities for retro- and prospective insight. *Quaternary Science Reviews* 197, 1–20. <https://doi.org/10.1016/j.quascirev.2018.07.009>
- Babst, F., Bouriaud, O., Poulter, B., Trouet, V., Girardin, M.P., Frank, D.C., 2019. Twentieth century redistribution in climatic drivers of global tree growth. *Science Advances* 5, eaat4313. <https://doi.org/10.1126/sciadv.aat4313>
- Babst, F., Friend, A.D., Karamihalaki, M., Wei, J., von Arx, G., Papale, D., Peters, R.L., 2021. Modeling Ambitions Outpace Observations of Forest Carbon Allocation. *Trends in Plant Science* 26, 210–219. <https://doi.org/10.1016/j.tplants.2020.10.002>
- Baker, J.C.A., Hunt, S.F.P., Clerici, S.J., Newton, R.J., Bottrell, S.H., Leng, M.J., Heaton, T.H.E., Helle, G., Argollo, J., Gloor, M., Brien, R.J.W., 2015. Oxygen isotopes in tree rings show good coherence between species and sites in Bolivia. *Global and Planetary Change* 133, 298–308. <https://doi.org/10.1016/j.gloplacha.2015.09.008>
- Baker, J.C.A., Santos, G.M., Gloor, M., Brien, R.J.W., 2017. Does Cedrela always form annual rings? Testing ring periodicity across South America using radiocarbon dating. *Trees*. <https://doi.org/10.1007/s00468-017-1604-9>
- Barichivich, J., Peylin, P., Launois, T., Daux, V., Risi, C., Jeong, J., Luyssaert, S., 2021. A triple tree-ring constraint for tree growth and physiology in a global land surface model. *Biogeosciences* 18, 3781–3803. <https://doi.org/10.5194/bg-18-3781-2021>
- Beck, J., Böller, M., Erhardt, A., Schwanghart, W., 2014. Spatial bias in the GBIF database and its effect on modeling species' geographic distributions. *Ecological Informatics* 19, 10–15. <https://doi.org/10.1016/j.ecoinf.2013.11.002>
- Belmecheri, S., Babst, F., Wahl, E.R., Stahle, D.W., Trouet, V., 2016. Multi-century evaluation of Sierra Nevada snowpack. *Nature Climate Change* 6, 2–3. <https://doi.org/10.1038/nclimate2809>
- Black, B.A., Griffin, D., van der Sleen, P., Wanamaker, A.D., Speer, J.H., Frank, D.C., Stahle, D.W., Pederson, N., Copenheaver, C.A., Trouet, V., Griffin, S., Gillanders, B.M., 2016. The value of crossdating to retain high-frequency variability, climate signals, and extreme events in environmental proxies. *Global Change Biology* 22, 2582–2595. <https://doi.org/10.1111/gcb.13256>
- Boninsegna, J.A., Argollo, J., Aravena, J.C., Barichivich, J., Christie, D., Ferrero, M.E., Lara, A., Le Quesne, C., Luckman, B.H., Masiokas, M., Morales, M., Oliveira, J.M., Roig, F., Srur, A., Villalba, R., 2009. Dendroclimatological reconstructions in South America: A review. *Palaeogeography, Palaeoclimatology, Palaeoecology* 281, 210–228. <https://doi.org/10.1016/j.palaeo.2009.07.020>
- Brien, R.J.W., Schöngart, J., Zuidema, P.A., 2016. Tree Rings in the Tropics: Insights into the Ecology and Climate Sensitivity of Tropical Trees, in: *Tropical Tree Physiology*. Springer International Publishing, pp. 439–461.
- Brien, R.J.W., Zuidema, P.A., 2005. Relating tree growth to rainfall in Bolivian rain forests: a test for six species using tree ring analysis. *Oecologia* 146, 1–12. <https://doi.org/10.1007/s00442-005-0160-y>
- Buckley, B.M., Anchukaitis, K.J., Penny, D., Fletcher, R., Cook, E.R., Sano, M., Nam le, C., Wichienkeo, A., Minh, T.T., Hong, T.M., 2010. Climate as a contributing factor in the demise of Angkor, Cambodia. *PNAS* 107, 6748–52. <https://doi.org/10.1073/pnas.0910827107>
- Bunn, A., Korpela, M., Biondi, F., Campelo, F., Mérian, P., Qeadan, F., Zang, C., 2023. dplR: Dendrochronology Program Library in R (R package version 1.7.6).
- Bunn, A.G., 2008. A dendrochronology program library in R (dplR). *Dendrochronologia* 26, 115–124. <https://doi.org/10.1016/j.dendro.2008.01.002>
- Caetano-Andrade, V.L., Clement, C.R., Weigel, D., Trumbore, S., Boivin, N., Schöngart, J., Roberts, P., 2020. Tropical Trees as Time Capsules of Anthropogenic Activity. *Trends in Plant Science* 25, 369–380. <https://doi.org/10.1016/j.tplants.2019.12.010>
- Chamberlain, S., Barve, V., Mcglinn, D., Oldoni, D., Desmet, P., Geffert, L., Ram, K., 2023. rgbif: Interface to the Global Biodiversity Information Facility API.
- Cook, B.I., Smerdon, J.E., Cook, E.R., Williams, A.P., Anchukaitis, K.J., Mankin, J.S., Allen, K., Andreu-Hayles, L., Ault, T.R., Belmecheri, S., Coats, S., Coulthard, B., Fosu, B., Grierson, P., Griffin, D., Herrera, D.A., Ionita, M., Lehner, F., Leland, C., Marvel, K., Morales, M.S., Mishra, V., Ngoma, J., Nguyen, H.T.T., O'Donnell, A., Palmer, J., Rao, M.P., Rodriguez-Caton, M., Seager, R., Stahle, D.W., Stevenson, S., Thapa, U.K., Varuolo-Clarke, A.M., Wise, E.K., 2022. Megadroughts in the Common Era and the Anthropocene. *Nature Reviews Earth & Environment* 3, 741–757. <https://doi.org/10.1038/s43017-022-00329-1>
- Cook, E.R., Anchukaitis, K.J., Brendan M. Buckley, D'Arrigo, R.D., Jacoby, G.C., Wright, W.E., 2010. Asian Monsoon Failure and Megadrought During the Last Millennium. *Science* 328, 486–489. <https://doi.org/10.1126/science.1185188>
- Corlett, R.T., 2013. Where are the Subtropics? *Biotropica* 45, 273–275. <https://doi.org/10.1111/btp.12028>
- Crowther, T.W., Glick, H.B., Covey, K.R., Bettigole, C., Maynard, D.S., Thomas, S.M., Smith, J.R., Hintler, G., Duguid, M.C., Amatulli, G., Tuanmu, M.N., Jetz, W., Salas, C., Stam, C., Piotto, D., Tavani, R., Green, S., Bruce, G., Williams, S.J., Wiser, S.K., Huber, M.O., Hengeveld, G.M., Nabuurs, G.J., Tikhonova, E., Borchardt, P., Li, C.F., Powrie, L.W., Fischer, M., Hemp, A., Homeier, J., Cho, P., Vibrans, A.C., Umunay, P.M., Piao, S.L., Rowe, C.W., Ashton, M.S., Crane, P.R., Bradford, M.A., 2015. Mapping tree density at a global scale. *Nature* 525, 201–205. <https://doi.org/10.1038/nature14967>
- D'Arrigo, R., Palmer, J., Ummenhofer, C.C., Kyaw, N.N., Krusic, P., 2011. Three centuries of Myanmar monsoon climate variability inferred from teak tree rings. *Geophysical Research Letters* 38. <https://doi.org/10.1029/2011GL049927>

- De Mil, T., Vannoppen, A., Beeckman, H., Van Acker, J., Van Den Bulcke, J., 2016. A field-to-desktop toolchain for X-ray CT densitometry enables tree ring analysis. *Ann Bot* 117, 1187–1196. <https://doi.org/10.1093/aob/mcw063>
- Fan, L., Wigneron, J.-P., Ciais, P., Chave, J., Brandt, M., Fensholt, R., Saatchi, S.S., Bastos, A., Al-Yaari, A., Hufkens, K., Qin, Y., Xiao, X., Chen, C., Myneni, R.B., Fernandez-Moran, R., Mialon, A., Rodriguez-Fernandez, N.J., Kerr, Y., Tian, F., Peñuelas, J., 2019. Satellite-observed pantropical carbon dynamics. *Nature Plants*. <https://doi.org/10.1038/s41477-019-0478-9>
- Feeley, K.J., Malhi, Y., Zelazowski, P., Silman, M.R., 2012. The relative importance of deforestation, precipitation change, and temperature sensitivity in determining the future distributions and diversity of Amazonian plant species. *Glob Change Biol* 18, 2636–2647. <https://doi.org/10.1111/j.1365-2486.2012.02719.x>
- Feeley, K.J., Wright, S.J., Supardi, M.N.N., Kassim, A.R., Davies, S.J., 2007. Decelerating growth in tropical forest trees. *Ecology Letters* 10, 461–469. <https://doi.org/10.1111/j.1461-0248.2007.01033.x>
- Fichtler, E., Worbes, M., 2010. Wood anatomy and tree-ring structure and their importance for tropical dendrochronology, in: Junk, W.J., Piedade, M.T.F., Wittmann, F., Schongart, J., Parolin, P. (Eds.), *Amazonian Floodplain Forests: Ecophysiology, Biodiversity and Sustainable Management*. Springer, Dordrecht, pp. 329–346.
- Fisher, R.A., Koven, C.D., Anderegg, W.R.L., Christoffersen, B.O., Dietze, M.C., Farrior, C.E., Holm, J.A., Hurtt, G.C., Knox, R.G., Lawrence, P.J., Lichstein, J.W., Longo, M., Matheny, A.M., Medvigy, D., Muller-Landau, H.C., Powell, T.L., Serbin, S.P., Sato, H., Shuman, J.K., Smith, B., Trugman, A.T., Viskari, T., Verbeeck, H., Weng, E., Xu, C., Xu, X., Zhang, T., Moorcroft, P.R., 2018. Vegetation demographics in Earth System Models: A review of progress and priorities. *Global Change Biology* 24, 35–54. <https://doi.org/10.1111/gcb.13910>
- Friedlingstein, P., O'Sullivan, M., Jones, M.W., Andrew, R.M., Hauck, J., Olsen, A., Peters, G.P., Peters, W., Pongratz, J., Sitch, S., Le Quéré, C., Canadell, J.G., Ciais, P., Jackson, R.B., Alin, S., Aragão, L.E.O.C., Arneeth, A., Arora, V., Bates, N.R., Becker, M., Benoit-Cattin, A., Bittig, H.C., Bopp, L., Bultan, S., Chandra, N., Chevallier, F., Chini, L.P., Evans, W., Florentie, L., Forster, P.M., Gasser, T., Gehlen, M., Gilfillan, D., Gkritzalis, T., Gregor, L., Gruber, N., Harris, I., Hartung, K., Haverd, V., Houghton, R.A., Ilyina, T., Jain, A.K., Joetzier, E., Kadono, K., Kato, E., Kitidis, V., Korsbakken, J.I., Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Liu, Z., Lombardozzi, D., Marland, G., Metzl, N., Munro, D.R., Nabel, J.E.M.S., Nakaoka, S.I., Niwa, Y., O'Brien, K., Ono, T., Palmer, P.I., Pierrot, D., Poulter, B., Resplandy, L., Robertson, E., Rödenbeck, C., Schwinger, J., Séférian, R., Skjelvan, I., Smith, A.J.P., Sutton, A.J., Tanhua, T., Tans, P.P., Tian, H., Tilbrook, B., van der Werf, G., Vuichard, N., Walker, A.P., Wanninkhof, R., Watson, A.J., Willis, D., Wiltshire, A.J., Yuan, W., Yue, X., Zaehle, S., 2020. Global Carbon Budget 2020. *Earth Syst. Sci. Data* 12, 3269–3340. <https://doi.org/10.5194/essd-12-3269-2020>
- Friend, A.D., Eckes-Shephard, A.H., Tupker, Q., 2022. Wood structure explained by complex spatial source-sink interactions. *Nat Commun* 13, 7824. <https://doi.org/10.1038/s41467-022-35451-7>
- Furnival, G.M., Wilson, R.W., 1974. Regressions by leaps and bounds. *Technometrics* 16, 499–511. <https://doi.org/10.1080/00401706.1974.10489231>
- Garcia, M.N., Hu, J., Domingues, T.F., Groenendijk, P., Oliveira, R.S., Costa, F.R.C., 2022. Local hydrological gradients structure high intraspecific variability in plant hydraulic traits in two dominant central Amazonian tree species. *Journal of Experimental Botany* 73, 939–952. <https://doi.org/10.1093/jxb/erab432>
- Gebrekirstos, A., Bräuning, A., Sass-Klassen, U., Mbow, C., 2014. Opportunities and applications of dendrochronology in Africa. *Current Opinion in Environmental Sustainability* 6, 48–53. <https://doi.org/10.1016/j.cosust.2013.10.011>
- Giraldo, J.A., del Valle, J.I., González-Caro, S., David, D.A., Taylor, T., Tobón, C., Sierra, C.A., 2023. Tree growth periodicity in the ever-wet tropical forest of the Americas. *Journal of Ecology* 111, 889–902. <https://doi.org/10.1111/1365-2745.14069>
- Godoy-Veiga, M., Ceccantini, G., Pitsch, P., Krottenthaler, S., Anhof, D., Locosselli, G.M., 2018. Shadows of the edge effects for tropical emergent trees: the impact of lianas on the growth of *Aspidosperma polyneuron*. *Trees* 32, 1073–1082. <https://doi.org/10.1007/s00468-018-1696-x>
- Granato-Souza, D., Stahle, D.W., Barbosa, A.C., Feng, S., Torbenson, M.C.A., de Assis Pereira, G., Schöngart, J., Barbosa, J.P., Griffin, D., 2019. Tree rings and rainfall in the equatorial Amazon. *Clim Dyn* 52, 1857–1869. <https://doi.org/10.1007/s00382-018-4227-y>
- Groenendijk, P., Bongers, F., Zuidema, P.A., 2017. Using tree-ring data to improve timber-yield projections for African wet tropical forest tree species. *Forest Ecology and Management*. <https://doi.org/10.1016/j.foreco.2017.05.054>
- Groenendijk, P., Sass-Klaassen, U., Bongers, F., Zuidema, P.A., 2014. Potential of tree-ring analysis in a wet tropical forest: A case study on 22 commercial tree species in Central Africa. *Forest Ecology and Management* 323, 65–78. <https://doi.org/10.1016/j.foreco.2014.03.037>
- Heilman, K.A., Dietze, M.C., Arizpe, A.A., Aragon, J., Gray, A., Shaw, J.D., Finley, A.O., Klesse, S., DeRose, R.J., Evans, M.E.K., 2022. Ecological forecasting of tree growth: Regional fusion of tree-ring and forest inventory data to quantify drivers and characterize uncertainty. *Global Change Biology* 28, 2442–2460. <https://doi.org/10.1111/gcb.16038>
- Heinrich, I., Weidner, K., Helle, G., Vos, H., Banks, J.C.G., 2008. Hydroclimatic variation in Far North Queensland since 1860 inferred from tree rings. *Palaeogeography, Palaeoclimatology, Palaeoecology* 270, 116–127. <https://doi.org/10.1016/j.palaeo.2008.09.002>
- Hijmans, R.J., 2021. geosphere: Spherical Trigonometry. R package version 1.5-14.
- Hughes, M.K., Swetnam, T.W., Diaz, H.F., 2010. *Dendroclimatology: progress and prospects*. Springer Science & Business Media.
- Humphrey, V., Zscheischler, J., Ciais, P., Gudmundsson, L., Sitch, S., Seneviratne, S.I., 2018. Sensitivity of atmospheric CO<sub>2</sub> growth rate to observed changes in terrestrial water storage. *Nature* 560, 628–631. <https://doi.org/10.1038/s41586-018-0424-4>
- Jansma, E., Brewer, P.W., Zandhuis, I., 2010. TRiDaS 1.1: The tree-ring data standard. *Dendrochronologia* 28, 99–130. <https://doi.org/10.1016/j.dendro.2009.06.009>
- Jeong, J., Barichivich, J., Peylin, P., Haverd, V., McGrath, M.J., Vuichard, N., Evans, M.N., Babst, F., Luyssaert, S., 2020. Using the International Tree-Ring Data Bank (ITRDB) records as century-long benchmarks for land-surface models. *Geosci. Model Dev. Discuss.* 2020, 1–45. <https://doi.org/10.5194/gmd-2020-29>
- Klesse, S., DeRose, R.J., Babst, F., Black, B.A., Anderegg, L.D.L., Axelson, J., Ettinger, A., Griesbauer, H., Guiterman, C.H., Harley, G., Harvey, J.E., Lo, Y.-H., Lynch, A.M., O'Connor, C., Restaino, C., Sauchyn, D., Shaw, J.D., Smith, D.J., Wood, L., Villanueva-Diaz, J., Evans, M.E.K., 2020. Continental-scale tree-ring-based projection of Douglas-fir growth:

- Testing the limits of space-for-time substitution. *Global Change Biology* 26, 5146–5163.  
<https://doi.org/10.1111/gcb.15170>
- Loader, N.J., Walsh, R.P.D., Robertson, I., Bidin, K., Ong, R.C., Reynolds, G., McCarroll, D., Gagen, M., Young, G.H.F., 2011. Recent trends in the intrinsic water-use efficiency of ringless rainforest trees in Borneo. *Phil. Trans. R. Soc. B* 366, 3330–3339. <https://doi.org/10.1098/rstb.2011.0037>
- Locosselli, G.M., Brien, R.J.W., de Souza Leite, M., Gloor, M., Krotenthaler, S., de Oliveira, A.A., Barichivich, J., Anhu, D., Ceccantini, G., Schöngart, J., Buckeridge, M., 2020. Global tree-ring analysis reveals rapid decrease in tropical tree longevity with temperature. *Proceedings of the National Academy of Sciences of the United States of America* 117, 33358–33364. <https://doi.org/10.1073/PNAS.2003873117>
- Lopez, L., Stahle, D., Villalba, R., Torbenson, M., Feng, S., Cook, E., 2017. Tree ring reconstructed rainfall over the southern Amazon Basin. *Geophysical Research Letters* 44, 7410–7418. <https://doi.org/10.1002/2017GL073363>
- Lumley, T. (based on F. code by A.M., 2020. *leaps: Regression Subset Selection*. R package version 3.1.
- McLeod, A.I., Xu, C., Lai, Y., 2020. Bestglm: Best Subset GLM and Regression Utilities. R Package Version 0.37.3.
- Menne, M.J., Durre, I., Vose, R.S., Gleason, B.E., Houston, T., \, ara G, 2012. An Overview of the Global Historical Climatology Network-Daily Database. *Journal of Atmospheric and Oceanic Technology* 29, 897–910. <https://doi.org/10.1175/jtech-d-11-00103.1>
- Meyer, C., Weigelt, P., Kreft, H., 2016. Multidimensional biases, gaps and uncertainties in global plant occurrence information. *Ecol Lett* 19, 992–1006. <https://doi.org/10.1111/ele.12624>
- Morales, M.S., Cook, E.R., Barichivich, J., Christie, D.A., Villalba, R., LeQuesne, C., Srur, A.M., Ferrero, M.E., González-Reyes, Á., Couvreur, F., Matuskovsky, V., Aravena, J.C., Lara, A., Mundo, I.A., Rojas, F., Prieto, M.R., Smerdon, J.E., Bianchi, L.O., Masiokas, M.H., Urrutia-Jalabert, R., Rodríguez-Catón, M., Muñoz, A.A., Rojas-Badilla, M., Alvarez, C., Lopez, L., Luckman, B.H., Lister, D., Harris, I., Jones, P.D., Williams, A.P., Velazquez, G., Aliste, D., Aguilera-Betti, I., Marcotti, E., Flores, F., Muñoz, T., Cuq, E., Boninsegna, J.A., 2020. Six hundred years of South American tree rings reveal an increase in severe hydroclimatic events since mid-20th century. *Proceedings of the National Academy of Sciences* 202002411. <https://doi.org/10.1073/pnas.2002411117>
- Pan, Y., Birdsey, R.A., Phillips, O.L., Houghton, R.A., Fang, J., Kauppi, P.E., Keith, H., Kurz, W.A., Ito, A., Lewis, S.L., Nabuurs, G.-J., Shvidenko, A., Hashimoto, S., Lerink, B., Schepaschenko, D., Castanho, A., Murdiyars, D., 2024. The enduring world forest carbon sink. *Nature* 631, 563–569. <https://doi.org/10.1038/s41586-024-07602-x>
- Pastorello, G., Trotta, C., Canfora, E., Chu, H., Christianson, D., Cheah, Y.-W., Poindexter, C., Chen, J., Elbashandy, A., Humphrey, M., Isaac, P., Polidori, D., Reichstein, M., Ribeca, A., van Ingen, C., Vuichard, N., Zhang, L., Amiro, B., Ammann, C., Arain, M.A., Ardö, J., Arkebauer, T., Arndt, S.K., Arriga, N., Aubinet, M., Aurela, M., Baldocchi, D., Barr, A., Beamesderfer, E., Marchesini, L.B., Bergeron, O., Beringer, J., Bernhofer, C., Berveiller, D., Billesbach, D., Black, T.A., Blanken, P.D., Bohrer, G., Boike, J., Bolstad, P.V., Bonal, D., Bonnefond, J.-M., Bowling, D.R., Bracho, R., Brodeur, J., Brümmer, C., Buchmann, N., Burban, B., Burns, S.P., Buysse, P., Cale, P., Cavagna, M., Cellier, P., Chen, S., Chini, I., Christensen, T.R., Cleverly, J., Collalti, A., Consalvo, C., Cook, B.D., Cook, D., Coursolle, C., Cremonese, E., Curtis, P.S., D'Andrea, E., da Rocha, H., Dai, X., Davis, K.J., Cinti, B.D., Grandcourt, A. de, Ligne, A.D., De Oliveira, R.C., Delpierre, N., Desai, A.R., Di Bella, C.M., Tommasi, P. di, Dolman, H., Domingo, F., Dong, G., Dore, S., Duce, P., Dufréne, E., Dunn, A., Dušek, J., Eamus, D., Eichmann, U., ElKhidir, H.A.M., Eugster, W., Ewenz, C.M., Ewers, B., Famulari, D., Fares, S., Feigenwinter, I., Feitz, A., Fensholt, R., Filippa, G., Fischer, M., Frank, J., Galvagno, M., Gharun, M., Gianelle, D., Gielen, B., Gioli, B., Gitelson, A., Gode, I., Goeckede, M., Goldstein, A.H., Gough, C.M., Goulden, M.L., Graf, A., Griebel, A., Gruening, C., Grünwald, T., Hammerle, A., Han, S., Han, X., Hansen, B.U., Hanson, C., Hatakka, J., He, Y., Hehn, M., Heinesch, B., Hinko-Najera, N., Hörtnagl, L., Hutley, L., Ibrom, A., Ikawa, H., Jackowicz-Korczynski, M., Janouš, D., Jans, W., Jassal, R., Jiang, S., Kato, T., Khomik, M., Klatt, J., Knohl, A., Knox, S., Kobayashi, H., Koerber, G., Kolbe, O., Kosugi, Y., Kotani, A., Kowalski, A., Kruijt, B., Kurbatova, J., Kutsch, W.L., Kwon, H., Launiainen, S., Laurila, T., Law, B., Leuning, R., Li, Yingnian, Liddell, M., Limousin, J.-M., Lion, M., Liska, A.J., Lohila, A., López-Ballesteros, A., López-Blanco, E., Loubet, B., Loustau, D., Lucas-Moffat, A., Lüers, J., Ma, S., Macfarlane, C., Magliulo, V., Maier, R., Mammarella, I., Manca, G., Marcolla, B., Margolis, H.A., Marras, S., Massman, W., Mastepanov, M., Matamala, R., Matthes, J.H., Mazzenga, F., McCaughey, H., McHugh, I., McMillan, A.M.S., Merbold, L., Meyer, W., Meyers, T., Miller, S.D., Minerbi, S., Moderow, U., Monson, R.K., Montagnani, L., Moore, C.E., Moors, E., Moreaux, V., Moureaux, C., Munger, J.W., Nakai, T., Neirynck, J., Nesic, Z., Nicolini, G., Noormets, A., Northwood, M., Noisetto, M., Nouvellon, Y., Novick, K., Oechel, W., Olesen, J.E., Ourcival, J.-M., Papuga, S.A., Parmentier, F.-J., Paul-Limoges, E., Pavelka, M., Peichl, M., Pendall, E., Phillips, R.P., Pilegaard, K., Pirk, N., Posse, G., Powell, T., Prasse, H., Prober, S.M., Rambal, S., Rannik, Ü., Raz-Yaseef, N., Rebmann, C., Reed, D., Dias, V.R. de, Restrepo-Coupe, N., Reverte, B.R., Roland, M., Sabbatini, S., Sachs, T., Saleska, S.R., Sánchez-Cañete, E.P., Sanchez-Mejia, Z.M., Schmid, H.P., Schmidt, M., Schneider, K., Schrader, F., Schroder, I., Scott, R.L., Sedláč, P., Serrano-Ortiz, P., Shao, C., Shi, P., Shironya, I., Siebicke, L., Šigut, L., Silberstein, R., Sirca, C., Spano, D., Steinbrecher, R., Stevens, R.M., Sturtevant, C., Suyker, A., Tagesson, T., Takanashi, S., Tang, Y., Tapper, N., Thom, J., Tomassucci, M., Tuovinen, J.-P., Urbanski, S., Valentini, R., van der Molen, M., van Gorsel, E., van Huissteden, K., Varlagin, A., Verfaillie, J., Vesala, T., Vincke, C., Vitale, D., Vygodskaya, N., Walker, J.P., Walter-Shea, E., Wang, H., Weber, R., Westermann, S., Wille, C., Wofsy, S., Wohlfahrt, G., Wolf, S., Woodgate, W., Li, Yuelin, Zampedri, R., Zhang, J., Zhou, G., Zona, D., Agarwal, D., Biraud, S., Torn, M., Papale, D., 2020. The FLUXNET2015 dataset and the ONEFlux processing pipeline for eddy covariance data. *Scientific Data* 7, 225. <https://doi.org/10.1038/s41597-020-0534-3>
- Perret, D.L., Evans, M.E.K., Sax, D.F., 2024. A species' response to spatial climatic variation does not predict its response to climate change. *Proc. Natl. Acad. Sci. U.S.A.* 121, e2304404120. <https://doi.org/10.1073/pnas.2304404120>
- Portal-Cahuana, L.A., Fontana, C., Assis-Pereira, G., Groenendijk, P., Roig, F.A., Tomazello-Filho, M., 2023. Thirty-four years of dendrochronological studies in Perú: A review of advances and challenges. *Dendrochronologia* 78, 126058. <https://doi.org/10.1016/j.dendro.2023.126058>
- Poulter, B., Frank, D., Ciais, P., Myneni, R.B., Andela, N., Bi, J., Broquet, G., Canadell, J.G., Chevallier, F., Liu, Y.Y., Running, S.W., Sitch, S., van der Werf, G.R., 2014. Contribution of semi-arid ecosystems to interannual variability of the global carbon cycle. *Nature* 509, 600–603. <https://doi.org/10.1038/nature13376>
- Pumijumngong, N., 2013. Dendrochronology in Southeast Asia. *Trees* 27, 343–358. <https://doi.org/10.1007/s00468-012-0775-7>

- Quesada-Román, A., Ballesteros-Cánovas, J.A., St. George, S., Stoffel, M., 2022. Tropical and subtropical dendrochronology: Approaches, applications, and prospects. *Ecological Indicators* 144, 109506. <https://doi.org/10.1016/j.ecolind.2022.109506>
- Quintilhan, M.T., Santini, L., Ortega Rodriguez, D.R., Guillemot, J., Cesilio, G.H.M., Chambi-Legoas, R., Nouvellon, Y., Tomazello-Filho, M., 2021. Growth-ring boundaries of tropical tree species: Aiding delimitation by long histological sections and wood density profiles. *Dendrochronologia* 69, 125878. <https://doi.org/10.1016/j.dendro.2021.125878>
- R Core Team, 2023. R: A Language and Environment for Statistical Computing.
- Rayback, S.A., Duncan, J.A., Schaberg, P.G., Kosiba, A.M., Hansen, C.F., Murakami, P.F., 2020. The DendroEcological Network: A cyberinfrastructure for the storage, discovery and sharing of tree-ring and associated ecological data. *Dendrochronologia* 60, 125678. <https://doi.org/10.1016/j.dendro.2020.125678>
- Rifai, S.W., Girardin, C.A.J., Berenguer, E., del Aguila-Pasquel, J., Dahlsjö, C.A.L., Doughty, C.E., Jeffery, K.J., Moore, S., Oliveras, I., Riutta, T., Rowland, L.M., Murakami, A.A., Addo-Danso, S.D., Brando, P., Burton, C., Ondo, F.E., Duah-Gyamfi, A., Amézquita, F.F., Freitag, R., Pacha, F.H., Huasco, W.H., Ibrahim, F., Mbou, A.T., Mihindou, V.M., Peixoto, K.S., Rocha, W., Rossi, L.C., Seixas, M., Silva-Espejo, J.E., Abernethy, K.A., Adu-Bredu, S., Barlow, J., da Costa, A.C.L., Marimon, B.S., Marimon-Junior, B.H., Meir, P., Metcalfe, D.B., Phillips, O.L., White, L.J.T., Malhi, Y., 2018. ENSO Drives interannual variation of forest woody growth across the tropics. *Phil. Trans. R. Soc. B* 373, 20170410. <https://doi.org/10.1098/rstb.2017.0410>
- Schöngart, J., Bräuning, A., Barbosa, A.C.M.C., Lisi, C.S., de Oliveira, J.M., 2017. Dendroecological Studies in the Neotropics: History, Status and Future Challenges, in: Amoroso, M.M., Daniels, L.D., Baker, P.J., Camarero, J.J. (Eds.), *Dendroecology: Tree-Ring Analyses Applied to Ecological Studies*. Springer International Publishing, Cham, pp. 35–73.
- Schulman, E., 1956. Dendroclimatic Changes in Semiarid America.
- Staal, A., Tuinenburg, O.A., Bosmans, J.H.C., Holmgren, M., van Nes, E.H., Scheffer, M., Zemp, D.C., Dekker, S.C., 2018. Forest-rainfall cascades buffer against drought across the Amazon. *Nature Climate Change* 8, 539–543. <https://doi.org/10.1038/s41558-018-0177-y>
- Stahle, D.W., Cook, E.R., Burnette, D.J., Villanueva, J., Cerano, J., Burns, J.N., Griffin, D., Cook, B.I., Acuña, R., Torbenson, M.C.A., Szejner, P., Howard, I.M., 2016. The Mexican Drought Atlas: Tree-ring reconstructions of the soil moisture balance during the late pre-Hispanic, colonial, and modern eras. *Quaternary Science Reviews* 149, 34–60. <https://doi.org/10.1016/j.quascirev.2016.06.018>
- Stahle, D.W., Diaz, J.V., Burnette, D.J., Paredes, J.C., Heim Jr, R.R., Fye, F.K., Soto, R.A., Therrell, M.D., Cleaveland, M.K., Stahle, D.K., 2011. Major Mesoamerican droughts of the past millennium. *Geophysical Research Letters* 38. <https://doi.org/10.1029/2010GL046472>
- Therrell, M.D., Stahle, D.W., Ries, L.P., Shugart, H.H., 2006. Tree-ring reconstructed rainfall variability in Zimbabwe. *Climate Dynamics* 26, 677–685. <https://doi.org/10.1007/s00382-005-0108-2>
- Van der Sleen, P., Groenendijk, P., Vlam, M., Anten, N.P.R., Boom, A., Bongers, F., Pons, T.L.T.L., Terburg, G., Zuidema, P.A., 2015a. No growth stimulation of tropical trees by 150 years of CO<sub>2</sub> fertilization but water-use efficiency increased. *Nature Geoscience* 8, 24–28. <https://doi.org/10.1038/ngeo2313>
- Van der Sleen, P., Groenendijk, P., Zuidema, P.A., 2015b. Tree-ring  $\delta^{18}O$  in African mahogany (*Entandrophragma utile*) records regional precipitation and can be used for climate reconstructions. *Global and Planetary Change* 127, 58–66. <https://doi.org/10.1016/j.gloplacha.2015.01.014>
- Villarreal, S., Vargas, R., 2021. Representativeness of FLUXNET Sites Across Latin America. *JGR Biogeosciences* 126. <https://doi.org/10.1029/2020JG006090>
- Wagner, F.H., Hérault, B., Bonal, D., Stahl, C., Anderson, L.O., Baker, T.R., Sebastian Becker, G., Beeckman, H., Boanerges Souza, D., Cesar Botosso, P., Bowman, D.M.J.S., Bräuning, A., Brede, B., Irving Brown, F., Julio Camarero, J., Camargo, P.B., Cardoso, F.C.G., Carvalho, F.A., Castro, W., Koloski Chagas, R., Chave, J., Chidumayo, E.N., Clark, D.A., Regina Capellotto Costa, F., Couralet, C., Henrique Da Silva Mauricio, P., Dalitz, H., Resende De Castro, V., Milani, J.E.D.F., Consuelo De Oliveira, E., De Souza Arruda, L., Devineau, J.L., Drew, D.M., Dünisch, O., Durigan, G., Elifuraha, E., Fedele, M., Ferreira Fedele, L., Figueiredo Filho, A., Finger, C.A.G., César Franco, A., Jnior, L.F., Galvão, F., Gebrekirstos, A., Gliniars, R., Mauricio Lima De Alencastro Graça, P., Griffiths, A.D., Grogan, J., Guan, K., Homeier, J., Raquel Kanieski, M., Khoon Kho, L., Koenig, J., Valerio Kohler, S., Krepkowski, J., Lemos-Filho, J.P., Lieberman, D., Eugene Lieberman, M., Sergio Lisi, C., Longhi Santos, T., Ayala, J.L.L., Eijji Maeda, E., Malhi, Y., Maria, V.R.B., Marques, M.C.M., Marques, R., Maza Chamba, H., Mbawambo, L., Liana Lisboa Melgaço, K., Angela Mendivelso, H., Murphy, B.P., O'Brien, J.J., F. Oberbauer, S., Okada, N., Plissier, R., Prior, L.D., Alejandro Roig, F., Ross, M., Rodrigo Rossatto, D., Rossi, V., Rowland, L., Rutishauser, E., Santana, H., Schulze, M., Selhorst, D., Rodrigues Silva, W., Silveira, M., Spann, S., Swaine, M.D., Toledo, J.J., Miranda Toledo, M., Toledo, M., Toma, T., Tomazello Filho, M., Ignacio Valdez Hernández, J., Verbesselt, J., Aparecida Vieira, S., Vincent, G., Volkmer De Castilho, C., Volland, F., Worbes, M., Lea Bolzan Zanon, M., Aragão, L.E.O.C., 2016. Climate seasonality limits leaf carbon assimilation and wood productivity in tropical forests. *Biogeosciences* 13, 2537–2562. <https://doi.org/10.5194/bg-13-2537-2016>
- Wang, J., Zeng, N., Wang, M., 2016. Interannual variability of the atmospheric CO<sub>2</sub> growth rate: roles of precipitation and temperature. *Biogeosciences* 13, 2339–2352.
- Williams, A.P., Cook, B.I., Smerdon, J.E., 2022. Rapid intensification of the emerging southwestern North American megadrought in 2020–2021. *Nature Climate Change* 12, 232–234. <https://doi.org/10.1038/s41558-022-01290-z>
- Wils, T.H.G., Robertson, I., Eshetu, Z., Koprowski, M., Sass-Klaassen, U.G.W., Touchan, R., Loader, N.J., 2010. Towards a reconstruction of Blue Nile baseflow from Ethiopian tree rings. *The Holocene* 20, 837–848. <https://doi.org/10.1177/0959683610365940>
- Woodborne, S., Hall, G., Robertson, I., Patrut, A., Rouault, M., Loader, N.J., Hofmeyr, M., 2015. A 1000-Year Carbon Isotope Rainfall Proxy Record from South African Baobab Trees (*Adansonia digitata* L.). *PLoS ONE* 10, e0124202. <https://doi.org/10.1371/journal.pone.0124202>
- Worbes, M., 2002. One hundred years of tree-ring research in the tropics – a brief history and an outlook to future challenges. *Dendrochronologia* 20, 217–231. <https://doi.org/10.1078/1125-7865-00018>

- Xu, X., Van Der Sleen, P., Groenendijk, P., Vlam, M., Medvigy, D., Moorcroft, P., Petticord, D., Ma, Y., Zuidema, P.A., 2024. Constraining long-term model predictions for woody growth using tropical tree rings. *Global Change Biology* 30, e17075. <https://doi.org/10.1111/gcb.17075>
- Zhao, S., Pederson, N., D'Orangeville, L., HilleRisLambers, J., Boose, E., Penone, C., Bauer, B., Jiang, Y., Manzanedo, R.D., 2019. The International Tree-Ring Data Bank (ITRDB) revisited: Data availability and global ecological representativity. *Journal of Biogeography* 46, 355–368. <https://doi.org/10.1111/jbi.13488>
- Zuidema, P.A., Babst, F., Groenendijk, P., Trouet, V., Abiyu, A., Acuña-Soto, R., Adeniesky-Filho, E., Alfaro-Sánchez, R., Aragão, J.R.V., Assis-Pereira, G., Bai, X., Barbosa, A.C., Battipaglia, G., Beeckman, H., Botosso, P.C., Bradley, T., Bräuning, A., Brien, R., Buckley, B.M., Camarero, J.J., Carvalho, A., Ceccantini, G., Centeno-Erguera, L.R., Cerano-Paredes, J., Chávez-Durán, Á.A., Cintra, B.B.L., Cleaveland, M.K., Couralet, C., D'Arrigo, R., del Valle, J.I., Dünisch, O., Enquist, B.J., Esemann-Quadros, K., Eshetu, Z., Fan, Z.-X., Ferrero, M.E., Fichtler, E., Fontana, C., Francisco, K.S., Gebrekirstos, A., Gloor, E., Granato-Souza, D., Haneca, K., Harley, G.L., Heinrich, I., Helle, G., Inga, J.G., Islam, M., Jiang, Y., Kaib, M., Khamisi, Z.H., Koprowski, M., Kruijt, B., Layme, E., Leemans, R., Leffler, A.J., Lisi, C.S., Loader, N.J., Locosselli, G.M., Lopez, L., López-Hernández, M.I., Lousada, J.L.P.C., Mendiavelso, H.A., Mokria, M., Montóia, V.R., Moors, E., Nabais, C., Ngoma, J., Nogueira Júnior, F. de C., Oliveira, J.M., Olmedo, G.M., Pagotto, M.A., Panthi, S., Pérez-De-Lis, G., Pucha-Cofrep, D., Pumijumong, N., Rahman, M., Ramirez, J.A., Requena-Rojas, E.J., Ribeiro, A. de S., Robertson, I., Roig, F.A., Rubio-Camacho, E.A., Sass-Klaassen, U., Schöngart, J., Sheppard, P.R., Slotta, F., Speer, J.H., Therrell, M.D., Toirambe, B., Tomazello-Filho, M., Torbenson, M.C.A., Touchan, R., Venegas-González, A., Villalba, R., Villanueva-Diaz, J., Vinya, R., Vlam, M., Wils, T., Zhou, Z.-K., 2022. Tropical tree growth driven by dry-season climate variability. *Nature Geoscience* 15, 269–276. <https://doi.org/10.1038/s41561-022-00911-8>
- Zuidema, P.A., Heinrich, I., Rahman, M., Vlam, M., Zwartsenberg, S.A., van der Sleen, P., 2020. Recent CO2 rise has modified the sensitivity of tropical tree growth to rainfall and temperature. *Global Change Biology* 26, 4028–4041. <https://doi.org/10.1111/gcb.15092>
- Zuidema, P.A., Poulter, B., Frank, D.C., 2018. A Wood Biology Agenda to Support Global Vegetation Modelling. *Trends in Plant Science* 23, 1006–1015. <https://doi.org/10.1016/j.tplants.2018.08.003>
- Zuquim, G., Costa, F.R.C., Tuomisto, H., Moulatlet, G.M., Figueiredo, F.O.G., 2020. The importance of soils in predicting the future of plant habitat suitability in a tropical forest. *Plant Soil* 450, 151–170. <https://doi.org/10.1007/s11104-018-03915-9>