

LABORATORY SOURCEBOOK
Advances in Physiology Education

TITLE: Large scale practical cardiovascular classes with *Danio rerio*: overcoming ethical, financial and logistical challenges associated with mammalian models

Laura F Corns¹, Nicholas S Freestone², James L Rouse³, Aidan Seeley⁴

¹ School of Biosciences, University of Sheffield, Sheffield, UK

² School of Life Sciences, Pharmacy and Chemistry, Kingston University, London, UK

³ School of Biology, University of Leeds, Leeds, UK

⁴ Swansea University Medical School, Swansea University, Swansea, UK

CORRESPONDING AUTHOR

Laura Corns

School of Biosciences

University of Sheffield

Western Bank

Sheffield

South Yorkshire

S10 2TN

0114 222 1092

l.f.corns@sheffield.ac.uk

ABBREVIATED TITLE Large scale practical cardiovascular classes with *D. rerio*

KEYWORDS 3Rs, partial replacement, Zebrafish

ABSTRACT (250 WORD LIMIT):

Traditional laboratory practicals exploring cardiovascular physiology and pharmacology rely on mammalian models, presenting ethical, financial, and logistical challenges. *Danio rerio* (zebrafish) larvae offer a compelling alternative that aligns with the partial replacement principle of the 3Rs, whilst providing an opportunity for students to develop desirable *in-vivo* skills to improve their employability. Here we introduce an engaging set of *in-vivo* laboratory practicals suitable for large undergraduate cohorts, that utilizes larval zebrafish to investigate cardiac ion channels and receptors. The practical involves two 3-hour sessions where students measure heart rate in 72 and 96 hours post-fertilisation larvae in response to various treatments. The first session introduces students to handling larval zebrafish before exploring the effects of a reduced ambient temperature and application of the commonly used zebrafish anaesthetic Tricaine (MS-222) on both heart rate and the zebrafish startle reflex. Finally, students apply the well known adrenergic agonist, adrenaline. The second session empowers students to develop their own testable hypothesis regarding which ion channels or receptors are likely to influence zebrafish heart rate, providing them with the autonomy to select two pharmacologically active drugs from a carefully curated list (e.g. isoproterenol (β -adrenergic receptor agonist), propranolol (β -adrenergic receptor antagonist) and nifedipine (L-type calcium channel blocker)) that will enable them to address

50 their hypothesis. Students' subsequent data for analysis allows them to develop an understanding of the
51 conserved and divergent aspects of cardiac physiology between zebrafish and mammalian systems, and an
52 appreciation of the importance of appropriate model selection in physiological and pharmacological
53 research.

54

55 **NEW AND NOTEWORTHY**

56 The document outlines how large-scale undergraduate practical classes involving *Danio rerio* (zebrafish)
57 can be used to teach cardiovascular physiology. It emphasizes the educational value of using live
58 zebrafish to explore heart rate, drug effects, and homeostasis. The process supports active, inquiry-based
59 learning, fostering engagement, critical thinking, and collaborative skills. It also addresses ethical and
60 logistical considerations. Overall, the approach effectively combines hands-on experimental experience
61 with core physiological concepts in an impactful educational format.

62

63 **INTRODUCTION**

64 Exploring the cardiovascular and autonomic nervous systems is a fundamental part of any physiology and
65 pharmacology education. To gain an understanding of human cardiac tissue, including the ion channels
66 and receptors that regulate its activity, traditional laboratory practicals have often relied on mammalian
67 models as a physiologically relevant alternative. However, mammalian models such as the Langendorff
68 preparation and other isolated heart preparations (1, 2), come with significant ethical, financial, and
69 logistical challenges. As an alternative to mammalian models, educators often turn to computational
70 models which can allow students to develop a good understanding of cardiac physiology (3, 4). However,
71 computational models do not address concerns around a reduction in the number of students graduating
72 without any *in-vivo* experience (5), which has led to an *in-vivo* skills gap in graduates (6-8). This
73 necessitates the exploration of alternative, robust, and educationally valuable non-mammalian *in-vivo*
74 models to study cardiovascular physiology and pharmacology.

75 Zebrafish (*Danio rerio*) offer a compelling alternative to mammalian models, particularly in larval stages,
76 aligning with the partial replacement aspect of the 3Rs (replacement, reduction, refinement). In the UK,
77 zebrafish larvae are not considered protected animals until they reach 5 days post-fertilisation (dpf),
78 aligning with the point at which they are capable of independent feeding (9, 10). Although not a full
79 replacement, this significantly reduces the ethical burden associated with their use, allowing for large-
80 scale, non-invasive experiments. Furthermore, zebrafish husbandry is relatively inexpensive and requires
81 less specialised infrastructure compared to mammalian housing, contributing to substantial reductions in
82 financial and logistical overheads. Here, we describe a laboratory practical using zebrafish which provides
83 students with hands-on experience of an *in-vivo* lower vertebrate model to investigate the ion channels
84 and receptors present in cardiac tissue.

85

86 **Objectives and Overview**

87

88 The modulation of zebrafish heart rate practical classes are suitable for large cohorts of undergraduate
89 students, and use the simple measurement of heart rate in larval zebrafish *in-vivo*. Students apply various
90 drugs, including those with anaesthetic properties and drugs known to influence human heart rate, and
91 observe the effects in the *in-vivo* model, generating their own data for subsequent analysis. This practical

also provides an excellent starting point for discussions around the ethics of animal use and the concepts of the 3Rs. The practical class described can be easily expanded for individual or group work projects, such as a final year capstone project.

The objectives of the described practical classes are (i) to use zebrafish appropriately to measure a key physiological variable, heart rate, *in-vivo*, (ii) to design appropriate experiments to investigate the effect of drugs on zebrafish heart rate, (iii) to analyse data relating to changes in heart rate and use this to evaluate the similarities and differences between mammalian and zebrafish hearts and (iv) to introduce students to the importance of appropriate models to investigate physiological function and evaluate pharmacological compounds.

Background

Zebrafish are a valued model across the biosciences from drug screening to understanding human disease, including cardiovascular disease (11-13). Zebrafish have a single circulatory system, which requires a two chambered heart consisting of a single atrium and a single ventricle; a significant structural difference from the four chambered mammalian heart. Despite this structural difference, the range of heart rate in an adult zebrafish (120-180 bpm; 14) is more similar to humans than that of common mammalian models such as the mouse. Adult zebrafish electrocardiograms show distinct P, QRS, and T waves, and the QT duration suggests a similar cardiac repolarisation time to humans (14). Most of the specialised cardiac cell types, for example, pacemaker cells, are found in both human and zebrafish hearts. The overall shape of a cardiac action potential is also similar to that of humans, with the depolarizing phase of the action potential mediated by voltage-gated Na^+ channels, the plateau phase mediated by voltage-gated L-type Ca^{2+} channels and the repolarizing phase mediated by the rapid delayed rectifier current (I_{Kr} ; 15-17). These similarities can be observed pharmacologically, as the voltage-gated Na^+ channel blockers, tetrodotoxin and tricaïne (also known as MS-222), reduce the action potential upstroke and the L-type Ca^{2+} channel blocker, nifedipine, shortens the plateau phase of zebrafish cardiac action potentials (16, 18). Although the zebrafish cardiac function is not fully developed in larval zebrafish, the ECG profile at 72 hours post-fertilisation (hpf) is very similar to that of adult zebrafish described above (19). Heart rate does vary with development, steadily increasing between 24 hpf to 96 hpf, where it can reach over 200 bpm, before slowing again as they reach adulthood (16, 18). Regardless of this increase, the larval zebrafish heart rate is still lower than that of commonly used mammalian models. Unlike endothermic mammals, zebrafish are ectotherms whose heart rate can be influenced by ambient temperature, with a substantial decrease in heart rate as ambient temperature is reduced from 28°C to 18°C (19). This makes zebrafish amenable to anaesthesia by rapid cooling, particularly in larval zebrafish (20), however, it is a major difference between zebrafish and mammals.

The heart rate of both humans and zebrafish is dynamically regulated by the autonomic nervous system, comprising the sympathetic and parasympathetic branches. Sympathetic stimulation, mediated by the release of norepinephrine from sympathetic neurons and activation of β -adrenergic receptors expressed by pacemaker cells and cardiac myocytes, increases heart rate and contractile strength. Conversely, parasympathetic stimulation, via acetylcholine (ACh) release from the vagus nerve, activates muscarinic receptors expressed by pacemaker cells and results in a decrease in heart rate. Some evidence suggests that this autonomic regulation can be seen early in zebrafish development, with increases in heart rate in response to the application of norepinephrine and the β -adrenergic agonist isoprenaline (also known as isoproterenol), observed from 96 hpf (18, 21, 22). The development of functional sympathetic innervation and thus the endogenous release of norepinephrine can be assessed by the observation of a decrease in heart rate in response to the β -adrenergic antagonist propranolol; in zebrafish this is first observed at 5 dpf

(21, 22). In relation to the parasympathetic system, muscarinic M2 receptors are expressed in cardiac tissue from 30 hpf (23), with the non-selective cholinergic agonist, Carbachol, evoking bradycardia from 72 hpf. In contrast to this, responses to the non-selective cholinergic agonist, ACh, are not seen until 5 dpf (21). The lack of response to the non-selective muscarinic antagonist, atropine, until 11-12 dpf has suggested that parasympathetic innervation does not occur until this time point (21). However, mild electrical stimulation at 0.5 Hz of 5-7 dpf zebrafish has been observed to induce atropine sensitive bradycardia and propranolol sensitive tachycardia (24), evidencing that central autonomic regulation of cardiac function is active in larval zebrafish.

These practicals give students a valuable opportunity to observe the effects of drugs on a simple measurement in an *in-vivo* model, something that simply cannot be achieved for logistical and ethical grounds in mammalian models. The similarities and differences between zebrafish and mammalian systems highlighted above offer excellent discussion points for students when considering zebrafish as a model organism for understanding cardiac physiology and testing cardiac drugs. This allows students to consider how aspects such as developmental age of a model and conditions such as temperature and the use of an anaesthetic can affect their results.

Learning Objectives:

After completing this activity, the student will be able to:

1. Explain what the term 'protected animal' means in relation to performing scientific experiments using *Danio rerio* (zebrafish) within the UK.
2. Design experiments to investigate the effect of drugs on heart rate in the *in-vivo* model, zebrafish.
3. Calculate and perform drug dilutions to administer drug compounds to the *in-vivo* model, zebrafish.
4. Use a dissecting microscope to observe changes in heart rate in the *in-vivo* model, zebrafish.
5. Analyse the data through data visualisation and using descriptive and inferential statistics, to determine the effect of drugs on heart rate in *Danio rerio*.
6. Using the data collected, discuss whether Tricaine is a useful anaesthetic when investigating zebrafish heart rate.
7. Using the data collected and knowledge of the mammalian heart, discuss the similarities and differences between the mammalian heart and the zebrafish heart.

Activity Level

The activity presented in its current form is suitable for all levels of undergraduate students in the general biomedical science disciplines; it acts as an early opportunity for students to engage with *in vivo* techniques and learn about the importance of choosing an appropriate model organism for their given question. Should the range of drugs detailed later feel inappropriate for a particular cohort of students, this can be amended to use a smaller, simpler range of drugs. Alternatively, this could also be easily adapted for students to complete as a final year capstone project or as a project for taught Master's students and Master's by Research students. If adapting for a project, students could have more freedom in choosing the drugs that they wished to investigate.

Prerequisite Student Knowledge or Skills

Before doing this activity, students should have a basic understanding of cardiovascular and autonomic physiology, more specifically the basic complement of ion channels that are expressed in mammalian cardiac tissue and the basic receptor subtypes used by the sympathetic and parasympathetic branches of the autonomic nervous system to regulate heart rate. It is also important that students have been taught how to behave appropriately when using a model organism, to be introduced to the zebrafish as the model organism that they will be using and the concepts of replacement, reduction and refinement (the 3Rs). Students should know how to use a standard laboratory micropipette, calculate drug dilutions and use a standard laboratory dissecting microscope.

Time Required

Zebrafish must be marbled 4 days in advance of the first session if one wishes to use zebrafish at 72 hpf (this is the age when zebrafish typically hatch). Marbling refers to the technique of placing a layer of marbles at the bottom of the fish tank to prevent predation of the zebrafish eggs. Here the eggs are protected by settling in between the marbles. The age at which initial experiments are performed can be adjusted slightly if needed to fit your timetable, however, be aware that if using fish earlier than 72 hpf, you will either have to dissect the zebrafish from their chorion or teach the students how to do this. This can be technically challenging and is only recommended if students have been taught microdissection previously.

The students require at least one 3 hour session to explore the basics of measuring zebrafish heart rate and modulating this by reducing the ambient temperature, applying the anaesthetic tricaine and the adrenergic receptor agonist, adrenaline (epinephrine). However, to incorporate experimental design and allow students to develop further hypotheses, it works better as two 3 hour sessions. The second session can be used to allow students to select two or three drugs from a curated selection to allow them to test their own specific hypothesis. For example, students could choose to focus on agonists and/or antagonists that are selective for α -adrenergic receptors, or for cholinergic receptors.

METHODS

Equipment and Supplies

Modulation of Zebrafish heart rate

The following should be available for a student to set up their workstation (Figure 1):

- 3 small petri dishes containing 5 ml of E3 medium and 72 hpf (for experiment 1) or 96 hpf (for experiment 2) zebrafish (nacre mutant).
- E3 media contains; 5 mM NaCl, 0.17 mM KCl, 0.33 mM CaCl₂, and 0.33 mM MgSO₄, pH 7.2–7.3, dissolved oxygen >6.3 mg/L, total hardness: 65 mg/L (as CaCO₃), temperature: 28 ± 1°C.
- Dissecting microscope
- Large petri dish (to use as an ice bath)
- Dissecting seeker
- Pasteur pipette
- Counter
- Stopwatch
- Thermometer
- Container of ice/ access to an ice machine
- Pipettes and tips (P5000, P1000 and P100)

- Waste bucket
- An Eppendorf with 5 mM tricaine
- An Eppendorf of 1 mM adrenaline

Eppendorfs containing the appropriate stock concentrations of the remaining drugs (Table 1) are available on request from the session leader. These drugs should be kept on ice during the practical to prevent degradation.

Ethical responsibilities of staff and students

A strain of zebrafish, nacre (25), is most useful for these experiments, as it lacks melanophores and thus allows the heart to be visualised more easily. However, most other zebrafish strains, including wild-type, could be used for these experiments as the pigmentation is relatively low and does not obstruct visualisation of the heart at the ages used. The zebrafish used in the experiment are used up to 96 hpf and are therefore exempt from the UK's Animals (Scientific Procedures) Act, 1986. Regardless, students are taught to look out for any signs of distress in the zebrafish and to notify the teaching staff if they suspect that the zebrafish has been damaged. Adopters of this activity are responsible for obtaining permission for human or animal research from their home institution. For a summary of Guiding Principles for Research Involving Animals and Human Beings, please see <https://www.physiology.org/mm/Publications/Ethical-Policies/Animal-and-Human-Research>.

Instructions

Preparation for the practicals

1. Production and care of zebrafish embryos
 - a. Zebrafish should be maintained at 28°C, ideally on a 14 hours light/10 hours dark cycle. Zebrafish can be maintained within a simple tank in tap water which requires regular cleaning (26) or a purpose built aquatic system which contains a circulating system that filters and aerates the water (27).
 - b. For breeding large numbers for an experiment, Westerfield (26) provides excellent detailed advice. Briefly, add the female to the tank with the male whilst adding a layer of marbles at the bottom of the tank. The following morning, the onset of light will initiate breeding and fertilised eggs will lay within the layer of marbles. The eggs can then be collected using a net, any debris removed and placed in a petri dish of E3 embryo media. Methylene blue (0.01%; Sigma-Aldrich) can be added as a fungicide. Embryos should be incubated at 28.5°C. Embryos can be sorted under a microscope to remove any unfertilised eggs; they should be cleaned daily.
 - c. Zebrafish embryos will hatch from their chorion between 48-72 hpf (28), making 72 hours a useful time to start experiments as one does not have to remove the chorion through dissection.
2. Stock solutions of Tricaine (5 mM; Ethyl 3-aminobenzoate methanesulfonate salt; Sigma-Aldrich) and all other experimental drugs (table 1) should be prepared in Eppendorfs in advance and frozen until the day of the practical.

Laboratory Session 1

Note that part of the aims of this session can be for students to decide on exactly how they will perform the experiments. The instructions given below are more extensive than those currently given to students.

Students are each given 3 zebrafish, each in a petri dish with 5 ml E3 medium, and perform all experiments described for session 1 on the same 3 zebrafish. You can choose to allow students to use a different zebrafish for each condition, however, in the interest of reducing the number of animals used for this practical class, this is not how it is currently delivered.

1. Label each petri dish so that control, drug and wash out recordings can be attributed to the correct zebrafish.
2. Test and record the temperature of the E3 medium for your control recordings.
3. Using the dissecting microscope identify the heart (highlighted by the white arrow in Figure 2) and measure the heart rate for each of your 72 hpf zebrafish in E3 medium. Use the counter to keep a record of how many beats are observed over a 30 second period then double this to calculate heart rate in beats per minute. Collect 3 repeats in total for each of the 3 zebrafish.
4. Test the presence of the zebrafish startle reflex by using the dissecting seeker to gently nudge the zebrafish tail; zebrafish that are not anaesthetised should swim away.
5. Assess the effect of reducing temperature to 15°C:
 - Create a cold water bath by adding a small amount of ice and some cold water to the large petri dish, testing the temperature using the thermometer until your water bath is just below 15°C. Transfer one of your petri dishes to this cold water bath, ensuring no cold water enters the petri dish as this will affect the osmolarity of the solution, and leave for a few minutes to reach the required temperature of 15°C. Add more ice to your water bath if required to achieve the correct temperature.
 - Once the correct temperature is achieved, measure heart rate 3 times and test the startle reflex before removing the petri dish containing the zebrafish back to the bench to return to your original control temperature. Repeat for the remaining 2 zebrafish.
 - Once the zebrafish have returned to control temperature, measure heart rate 3 times and test the startle reflex. A warm water bath can be used to speed up this return to room temperature.
6. Using the same 3 zebrafish, assess the effect of 500 µM Tricaine on zebrafish heart rate and immobilisation.
 - Dilute 5 mM Tricaine to 500 µM Tricaine within the petri dish containing the zebrafish. Avoiding the zebrafish, gently remove 500 µl of E3 medium from the first petri dish and add 500 µl of 5 mM Tricaine.
 - After 5 minutes, use the seeker to test the startle reflex. If there is no reflex, measure the heart rate 3 times.
 - Use a pasteur pipette to transfer the zebrafish to a fresh petri dish containing 5 ml of E3 medium. After 5 minutes, use the seeker to test the startle reflex. Once the reflex has returned, measure the heart rate 3 times.
7. Using the same 3 zebrafish, assess the effect of 10 µM adrenaline (epinephrine) on zebrafish heart rate.
 - Avoiding the zebrafish, gently remove 50 µl of E3 medium from the petri dish and add 50 µl of 1 mM adrenaline. After 5 minutes, measure the heart rate 3 times.
 - Use a pasteur pipette to transfer the zebrafish to a fresh petri dish containing 5 ml of E3 medium. After 5 minutes, measure the heart rate. If it has not returned to control values, wait a further 5 minutes and measure heart rate again.
8. Any zebrafish used in experiments should be euthanized by placing them in 1% sodium hypochlorite (bleach) for at least five minutes. Your session lead will dispose of euthanized zebrafish.

Laboratory session 2

- 339 9. Using the same protocols as Laboratory session 1, first measure the heart rate for each of your 96
340 hpf zebrafish in E3 medium. This is important as you are using 3 new zebrafish and they are now
341 24 hours older, so you must establish a new baseline heart rate.
- 342 10. Once you have developed your hypothesis regarding zebrafish cardiac ion channels and/or
343 receptors, choose 2 appropriate drugs from Table 1 to test this hypothesis. Assess the effect of
344 your 2 chosen drugs on zebrafish heart rate using the methods from laboratory session 1. You can
345 also apply two drugs together if this helps to test your hypothesis. The drugs (Table 1) are not an
346 exhaustive list of what can be given to students but are those often used for convenience and
347 relevance to our students.
- 348 11. Any zebrafish used in experiments should be euthanized by placing them in 1% sodium
349 hypochlorite (bleach) for at least five minutes. Your session lead will dispose of euthanized
350 zebrafish.
351
352

353 **Table 1:** Adrenaline (also known as epinephrine) and other optional compounds. Suggested stock and
354 working concentrations are provided, along with the known molecular target and known effect in humans.
355 *see 'Expected results' for a more detailed description of this expected effect.

356
357
358
359
360
361
362
363

359
360
361
362
363

Drug	Known target in mammals	Expected effect	Stock conc	Working conc	Supplier details
Adrenaline (epinephrine)	Non-selective α/β adrenergic agonist, some selectivity for β_2	Increase in heart rate	1 mM	10 μ M	Sigma Aldrich, Cat No E4250
Isoprenaline	Non-selective β adrenergic agonist	Increase in heart rate	10 mM	100 μ M	Sigma Aldrich, Cat No I5627
Salbutamol	β_2 adrenergic agonist	Increase in heart rate	1 mM	10 μ M	Sigma Aldrich, Cat No S8260
Propranolol	Non-selective β adrenergic antagonist	Decrease in heart rate*	1 mM	10 μ M	Sigma Aldrich, Cat No P0689
Atenolol	Selective β_1 adrenergic antagonist	Decrease in heart rate*	10 mM	1 mM	Sigma Aldrich, Cat No A7655
Acetylcholine (ACh)	Non-selective ACh receptor agonist	Decrease in heart rate*	10 mM	1 mM	Sigma Aldrich, Cat No A6625
Bethanechol	Muscarinic ACh receptor agonist	Decrease in heart rate*	10 mM	100 μ M	Sigma Aldrich, Cat No PHR2357
Nifedipine	L-type Ca^{2+} channel blocker	Decrease in heart rate*	1 mM	10 μ M	Sigma Aldrich, Cat No N7634

364
365
366
367
368
369
370

372
373

individual variation in the development, with some zebrafish not exhibiting a startle reflex (30). This is no reason to abandon this zebrafish. If the tricaine does not appear to be inducing a loss of the startle reflex, students should first consider whether they have pipetted the correct amount. Subsequently, they should check whether they have overstimulated the zebrafish with excessive light and thus an increase in ambient temperature or through excessive movement of the fluid and use of the dissecting seeker. These considerations should also be made if the heart rate does not return to baseline levels following removal of the anaesthetic. Variations in responses are expected and this can prove a useful point to discuss the variability of real biological data. If zebrafish responses become more variable over the course of the application of drugs, this can be used a useful discussion point for the balance between collecting good quality data and reducing the number of animals used.

Recommended working concentrations of drugs are given based on their potency and considerations given to the maximum concentrations considered safe for undergraduate students to work with. The working concentrations suggested for drugs all evoke responses where expected based on the expression of receptors and development of the autonomic nervous system (see 'Expected Results'). Higher working concentrations could be used if instructors deem this safe for their students on consultation of the material safety data sheets. It should be noted that the Atenolol and ACh stocks are made up in E3 media, so the 1:10 dilution should not affect the osmolarity of the solution or introduce another factor which could affect heart rate. This could be used as point at which you can introduce the concept that a solvent used to dissolve a drug could influence the experiment, and that where this is the case appropriate solvent controls should be used.

Zebrafish can become more active at 96 hpf, swimming around the petri dish more, which can make it more difficult to visualise the heart for a long enough time for counting. Students can be encouraged to use Tricaine as an anaesthetic or create an ice bath to lower the body temperature of the zebrafish and thus reduce zebrafish movement. If expanding this into a project, students could be taught to mount zebrafish in 3% methylcellulose to prevent excessive movement and increase accuracy of heart rate measurements. To achieve this, prepare the 3% methylcellulose (Sigma, M-0387) in E3 medium and place on a shaker to dissolve overnight. Transfer a small amount of 3% methylcellulose onto a shallow glass depression slide (SLS Select Slides Single Cavity; MIC3450). Transfer the zebrafish on top of the methylcellulose and cover with E3 medium, with or without your test drug, to fill the depression. Gently press the zebrafish into the methylcellulose and adjust it to the correct orientation to view the heart (31). To change the solution simply remove the solution from the top of the methylcellulose, replace with the new solution and leave to equilibrate for 2 minutes.

At 96 hpf the zebrafish heart rate also increases (18) making it more difficult to count. If not using a camera, it can be easier to encourage students to select drugs which are more likely to slow down the heart rate. One could complete both experiments using 72 hpf zebrafish if both sessions are run on the same day, however, one is likely to see responses to fewer of the drugs (see 'Expected Results'). As a single batch of zebrafish embryos can be 200 or more, to reduce the use of animals and prevent excess waste of zebrafish, the sessions could be run on the same or concurrent days.

Safety Considerations

For animal welfare, we advise students to use a low intensity of light from the dissecting microscope.

Due to the potentially toxic nature of some of the drugs, stock solutions are kept in small quantities in Eppendorf tubes and students are required to ask staff for the drugs when ready to perform their experiments. Recommended working concentrations of drugs are given.

Before any students in the UK undertake laboratory work of the type described in this paper they must fill out forms relating to the identification of hazards relating to the planned experiments. These COSHH forms (Control of Substances Hazardous to Health) are a ubiquitous feature of experimental work in UK universities and students are well versed in these safety protocols and acceptable ways of working.

Students are therefore required to provide confirmation that they have read the COSHH form for this experiment before they can take part. Additionally, students are trained how to take gloves off appropriately to ensure the skin is never in contact with the external aspect of the gloves.

RESULTS

Expected Results

Students measuring heart rate of larval zebrafish by eye *in-vivo* found that the heart rate at 72 hpf was 156 ± 25 bpm ($n = 120$) and this decreased significantly (73 ± 27 bpm, $n = 120$; $p < 0.001$) when the ambient temperature was reduced to 15°C (Figure 3A). Heart rate was capable of a full recovery to baseline when zebrafish were returned to the original control temperature. At 72 hpf, 80% of zebrafish displayed a startle reflex ($n = 120$), whereas only 26% displayed a startle reflex at 15°C ($n = 120$); as with heart rate a full recovery was observed when returned to control temperatures.

Tricaine (MS-222) is voltage-gated Na^{+} channel blocker and a commonly used anaesthetic in zebrafish. It was expected to abolish the startle reflex and decrease heart rate (9). Student experiments showed a decrease in the number of zebrafish exhibiting a startle response from 78% in control conditions to 24% in $500 \mu\text{M}$ tricaine ($n = 117$), with the vast majority of zebrafish (71%; $n = 116$) showing a recovered startle reflex on return to E3 media. A small but significant decrease in heart rate was observed by students (control, 160 ± 26 bpm, $n = 117$; tricaine, 155 ± 27 bpm, $n = 117$, return to control, 167 ± 26 bpm, $n = 116$; $p < 0.01$; Figure 3B).

When turning attention to drugs known to regulate heart rate in humans, as expected, students observed a reversible increase in heart rate on application of the non-selective adrenergic agonist adrenaline (control, 168 ± 29 bpm, $n = 78$; $10 \mu\text{M}$ adrenaline, 194 ± 36 bpm, $n = 78$, return to control, 168 ± 27 bpm, $n = 71$; $p < 0.0001$; Figure 3C). Moving to the optional drugs that can be chosen by students based on their specific hypothesis, the isoprenaline, a non-selective β adrenergic agonist (18), whilst the effects of Salbutamol, a β_2 adrenergic agonist, are more varied due to the expression of two different β_2 adrenergic receptor genes in the zebrafish heart which have been shown to have complex effects on heart rate (22). The potential effects of Salbutamol, provides a platform to discuss the possible cardiac side effects of its use in the treatment of asthma. It also allows for a separate discussion about how many animals, including zebrafish, have multiple genes for a single protein compared to humans having a single gene and what this means for studying these species.

The application of the non-selective β adrenergic antagonist, propranolol, and the selective β_1 adrenergic antagonist, atenolol, are not expected to elicit a change in heart rate at 96 hpf. Responses have previously been observed at 5 dpf (21, 22), suggesting sympathetic regulation is not active until this point. However, at 96 hpf the increase in heart rate observed in response to adrenaline can be reduced by the co-application of either propranolol and atenolol. These differences can be used as useful discussion points for choosing the correct developmental timepoints for studying a certain function within a given organism, and for a discussion around the mechanism of action of antagonists in the presence and absence of endogenous and exogenous agonists.

Moving to the drugs modulating the parasympathetic control of cardiac function, the non-selective cholinergic agonist, acetylcholine (ACh) and the muscarinic cholinergic agonist, bethanechol, may be expected to decrease heart rate. However, given the varied developmental time points (3-5 dpf) at which responses in heart rate to cholinergic agonists have been observed (23, 21), it is not surprising that many students do not observe responses to these agonists. Again, this does provide an opportunity to discuss the importance of choosing a model organism at an appropriate time point for your given hypothesis.

Finally, the L-type Ca^{2+} channel blocker, nifedipine, usually produces a decrease in heart rate (32). However, an increase in heart rate is sometimes observed, which could be analogous to the reflex tachycardia seen in humans in response to high doses of nifedipine (33). This response allows for a discussion around the complexities of *in vivo* studies and the importance of considering how drugs may impact on the vasculature as well as the heart.

Misconceptions

As mentioned when troubleshooting, there are misconceptions from students about zebrafish being defective if they do not exhibit a startle reflex under control conditions. Although the tactile startle reflex generally develops by 48 hpf (28), there can be inter-individual variation in the development, with some zebrafish not exhibiting a startle reflex (30).

Evaluation of Student Work

Inquiry Applications

This practical allows students to experience facilitated inquiry, where there are broad guidelines for the research question and methods, but the students need to specify the exact question in choosing their drugs and design the finer details of the method. They are given the question 'Are zebrafish a useful model for investigating cardiac physiology?' and told to focus on the similarities in changes of heart rate in response to drugs which have known actions in humans. The first session is very guided and then the second allows for the development and testing of their own hypotheses. They are given the freedom to choose whether to perform the second day's experiments with Tricaine anaesthesia or at a lower temperature, and whether they wish to apply drugs alone or together.

The activity could be made more student-centred by allowing them to choose which aspect of zebrafish physiology or behaviour they wish to investigate to determine whether they are a useful model. They could be allowed to research the experimental drugs from scratch rather than choosing from a defined list, or consider manipulating ion concentrations in the E3 medium to assess the effect on homeostasis.

Wider Educational Applications

As discussed in the expected results section, this practical series allows one to explore numerous topics including the mechanism of action of antagonists, the localised expression of specific receptor subtypes and mechanisms underlying side effects of clinical drugs, the complications arising from using an *in vivo* model and the importance of using appropriate developmental ages for studying different systems. Any of these topics could be expanded upon to create additional problem-based learning studies.

ACKNOWLEDGMENTS

To Professor Louise Robson for her support and guidance when initially developing this practical. To the Biological Services aquarium team at the University of Sheffield for breeding the zebrafish and the School of Biosciences teaching technical team, for their yearly preparation of these practicals. To all the students who have applied themselves and engaged positively with this practical and the principles that it is teaching.

REFERENCES

1. **Bell, RM, Mocanu, MM, Yellon, DM.** (2011). Retrograde heart perfusion: The Langendorff technique of isolated heart perfusion. *Advances in Physiology Education Sourcebook*, 50(6), 940–950. <https://doi.org/10.1016/j.yjmcc.2011.02.018>.
2. **Liao R, Podesser BK, Lim CC.** The continuing evolution of the Langendorff and ejecting murine heart: new advances in cardiac phenotyping. *Am J Physiol Heart Circ Physiol*, 303(2), H156–67, 2012. <https://doi.org/10.1152/ajpheart.00333.2012>.
3. **Dewhurst DG, Ulyott RT.** Computer Simulated Demonstration of the Actions of Drugs on the Isolated Perfused Mammalian Heart (Langendorff Preparation). *Alternatives to Laboratory Animals*, 19(3), 316–322, 1991. <https://doi.org/10.1177/026119299101900305>.
4. **Samsel, RW, Schmidt, GA, Hall, JB, Wood, LD, Shroff, SG, & Schumacker, PT.** Cardiovascular physiology teaching: computer simulations vs. animal demonstrations. *Advances in Physiology Education*, 266(6), S36, 1994. <https://doi.org/10.1152/advances.1994.266.6.S36>.
5. **British Pharmacological Society.** A survey of integrative physiology/pharmacology teaching undertaken by the BPS and the Physiological Society. *pA2 Online* 3: 10–11, 2004.
6. **The Association of the British Pharmaceutical Industry.** Sustaining the Skills Pipeline in the Pharmaceutical and Biopharmaceutical Industries [Online]. <https://www.abpi.org.uk/media/ou0hnrf2/sustain-skills-pipeline.pdf>.
7. **The Association of the British Pharmaceutical Industry.** Skills Needs for Biomedical Research [Online]. <https://www.abpi.org.uk/media/bufdzm3b/skills-biomedical-research.pdf>.
8. **The Association of the British Pharmaceutical Industry.** Bridging the skills gap in the biopharmaceutical industry [Online]. <https://www.abpi.org.uk/publications/bridging-the-skills-gap-in-the-biopharmaceutical-industry-2022/>
9. Animals (Scientific Procedures) Act 1986, c.1. Available at: <https://www.gov.uk/government/publications/the-operation-of-the-animals-scientific-procedures-act-1986> (Accessed: 24 April 2025).
10. **Strähle, U, Scholz, S, Geisler, R, Greiner, P, Hollert, H, Rastegar, S, Schumacher, A, Selderslaghs, I, Weiss, C, Witters, H, Braunbeck, T.** Zebrafish embryos as an alternative to animal experiments—A commentary on the definition of the onset of protected life stages in animal welfare regulations. *Reproductive Toxicology*, 33(2), 128–132, 2012. <https://doi.org/10.1016/j.reprotox.2011.06.121>.
11. **Bowley, G, Kugler, E, Wilkinson, R, Lawrie, A, van Eeden, F, Chico, TJA, Evans, PC, Noël, ES, Serbanovic-Canic, J.** Zebrafish as a tractable model of human cardiovascular disease. *Br J Pharmacol* 179:900–917, 2022. <https://doi.org/10.1111/bph.15473>.
12. **Choi, T, Choi, TI, Lee, YR, Choe, SK, Kim CH.** Zebrafish as an animal model for biomedical research. *Exp Mol Med* 53:310–317, 2021. <https://doi.org/10.1038/s12276-021-00571-5>
13. **Patton, EE, Zon, LI, Langenau, DM.** Zebrafish disease models in drug discovery: from preclinical modelling to clinical trials. *Nat Rev Drug Discov* 20: 611–628, 2021. <https://doi.org/10.1038/s41573-021-00210-8>
14. **Milan, DJ, Jones, IL, Ellinor, PT, MacRae, CA.** In vivo recording of adult zebrafish electrocardiogram and assessment of drug-induced QT prolongation. ¹ *American Journal of Physiology-Heart and Circulatory Physiology*, 291(1), H469–H477, 2006. <https://doi.org/10.1152/ajpheart.00960.2005>

15. Baker, K, Warren, KS, Yellen, G, & Fishman, MC. Defective “pacemaker” current (I_h) in a zebrafish mutant with a slow heart rate, *Proc. Natl. Acad. Sci. U.S.A.* 94 (9) 4554-4559, 1997. <https://doi.org/10.1073/pnas.94.9.4554>.
16. Nemtsas, P, Wettwer, E, Christ, T, Weidinger, G, Ravens, U. Adult zebrafish heart as a model for human heart? An electrophysiological study. *Advances in Physiology Education*, 48(1), 161–171, 2010. <https://doi.org/10.1016/j.yjmcc.2009.08.034>.
17. Hassel, D, Scholz, EP, Trano, N, Friedrich, O, Just, S, Meder, B, Weiss, DL, Zitron, E, Marquart, S, Vogel, B, Karle, CA, Seemann, G, Fishman, MC, Katus, HA, Rottbauer, W. Deficient Zebrafish Ether-à-Go-Go-Related¹ Gene Channel Gating Causes Short-QT Syndrome in Zebrafish Reggae Mutants. *Circulation*, 117(7), 857–864, 2008. <https://doi.org/10.1161/CIRCULATIONAHA.107.75222>
18. De Luca, E, Zaccaria, GM, Hadhoud, M, Rizzo, G, Ponzini, R, Morbiducci, U, Santoro, MM. ZebraBeat: a flexible platform for the analysis of the cardiac rate in zebrafish embryos. *Sci Rep*, 4:4898. 2014. <https://doi.org/10.1038/srep04898>.
19. Dhillon, SS, Dóró, É, Magyary, I, Egginton, S, Sík, A, Müller, F. Optimisation of Embryonic and Larval ECG Measurement in Zebrafish for Quantifying the Effect of QT Prolonging Drugs. *PLoS ONE*, 8(4), e60552, 2013. <https://doi.org/10.1371/journal.pone.0060552>
20. Chen, K, Wang, C-Q, Fan, Y-Q, Xie, Y-S, Yin, Z-F, Xu, Z-J, Zhang, H-L, Cao, J-T, Han, Z-H, Wang, Y, Sun, D-Q. The Evaluation of Rapid Cooling As an Anesthetic Method for the Zebrafish. *Zebrafish*, 11(1), 91–96, 2014. <https://doi.org/10.1089/zeb.2012.0858>
21. Schwerte, T, Prem, C, Mairösl, A, Pelster, B. Development of the sympatho-vagal balance in the cardiovascular system in zebrafish (*Danio rerio*) characterized by power spectrum and classical signal analysis. *The Journal of experimental biology*, 209(Pt 6), 1093–1100 2006. <https://doi.org/10.1242/jeb.02117>.
22. Steele, SL, Yang, X, Debais-Thibaud, M, Schwerte, T, Pelster, B, Ekker, M, Tiberi, M, Perry, SF In vivo and in vitro assessment of cardiac β -adrenergic receptors in larval zebrafish (*Danio rerio*). *J Exp Biol*, 214(9), 1445–1457, 2011. <https://doi.org/10.1242/jeb.052803>.
23. Hsieh, DJY, Liao, C-F. Zebrafish M2 muscarinic acetylcholine receptor: cloning, pharmacological characterization, expression patterns and roles in embryonic bradycardia. *British Journal of Pharmacology*, 137(6), 782–792, 2002. <https://doi.org/10.1038/sj.bjp.0704930>.
24. Lister, JA, Robertson, CP, Lepage, T, Johnson, SL, and Raible, DW. nacre encodes a zebrafish microphthalmia-related protein that regulates neural-crest-derived pigment cell fate. *Development*, 126(17), 3757–3767 1999, <https://doi.org/10.1242/dev.126.17.3757>
25. Mann, KD, Hoyt, C, Feldman, S, Blunt, L, Raymond, A, Page-McCaw, PS. Cardiac response to startle stimuli in larval zebrafish: sympathetic and parasympathetic components. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 298(5), R1288-R1297, 2010. <https://doi.org/10.1152/ajpregu.00302.2009>
26. Westerfield, M. The Zebrafish Book, 5th Edition; A guide for the laboratory use of zebrafish (*Danio rerio*), 2007 Eugene, University of Oregon Press. Paperback.
27. Avdesh, A, Chen, M, Martin-Iverson, MT, Mondal, A, Ong, D, Rainey-Smith, S, Taddei, K, Lardelli, M, Groth, DM, Verdile, G, Martins, RN. Regular Care and Maintenance of a Zebrafish (*Danio rerio*) Laboratory: An Introduction. *J. Vis. Exp.* (69), e4196, 2012. <https://doi.org/10.3791/4196>.
28. Kimmel, CB, Ballard, WW, Kimmel, SR, Ullmann, B, Schilling, TF. Stages of embryonic development of the zebrafish. *Dev Dyn.* 203(3), p253-310, 1995. <https://doi.org/10.1002/aja.1002030302>.
29. Saint-Amant, L, Drapeau, P. Time course of the development of motor behaviors in the zebrafish embryo. *Journal of neurobiology*, 37(4), 622–632, 1998, [https://doi.org/10.1002/\(sici\)1097-4695\(199812\)37:4](https://doi.org/10.1002/(sici)1097-4695(199812)37:4)

30. Fitzgerald, JA, Kirla, KT, Zinner, CP, & Vom Berg, CM. Emergence of consistent intra-individual locomotor patterns during zebrafish development. *Scientific reports*, 9(1), 13647. 2019 <https://doi.org/10.1038/s41598-019-49614-y>.
31. Liang, JO, Ahmed, S, Akusoba, C, Amacher, S, Bell, A, Brick, J, Butto, A, Byrne, R, Carson, H, Chau, E, Chen, J, Connor, E, Deininger, K, Erickson, J, Freeman, E, Friedenson, B, Grinblat, Y, Halloran, M, Holt, V, Ilagan, K, Ionescu, R, Karanovic, D, Kong, N, Kramek, H, Lee, J-H, Manieri, A, Manohar, P, Michelson, A, Morris, J, Neyman, A, Noon, K, Nothelle, S, Ohtola, J, Pierce, L, Pierce, S, Robinson, J, Romero, M, Schiller, B, Schreiner, A, Shuda, J, Stubblefield-Park, S, Xia, M, Yeganeh, A. *Zebrafish in the Classroom*. 2007. Direct Website Submission (<http://www.zfic.org>).
32. Salgado-Almario, J, Vicente, M, Molina, Y, Martinez-Sielva, A, Vincent, P, Domingo, B, Llopis, J. Simultaneous imaging of calcium and contraction in the beating heart of zebrafish larvae. *Theranostics*, 12(3), 1012–1029, 2022, <https://doi.org/10.7150/thno.64734>
33. Dustan, HP. Calcium channel blockers. Potential medical benefits and side effects. *Hypertension*, 13(5 Suppl), I137–I140. 1989 https://doi.org/10.1161/01.hyp.13.5_suppl.i137

Figure Legends

Figure 1: Equipment needed for the zebrafish experiments. Students require a standard dissecting microscope, 3 small petri dishes containing 5 ml of E3 medium and 72 hpf (for experiment 1) or 96 hpf (for experiment 2) zebrafish (nacre mutant), a counter and a stopwatch to perform basic measurements, a dissecting seeker (highlighted by the white arrow) to enable the startle reflex to be tested, a large petri dish to use as an ice bath and a thermometer to achieve the required temperature, a Pasteur pipette to transfer zebrafish between petri dishes, pipettes and tips (P5000, P1000 and P100) and stock concentrations of tricaine and adrenaline.

Figure 2. A 72 hpf (3 dpf) nacre zebrafish. The arrow indicates the zebrafish heart. Note the ability to see the blood in this region.

Figure 3. Modulation of embryonic zebrafish heart rate. **A)** Decreasing temperature from 22°C to 15°C reduces heart rate in 3 dpf zebrafish (n=119). **B)** Tricaine (500µM) elicits a small decrease in heart rate in 3dpf zebrafish (n = 113). **C)** Adrenaline (10µM) increases heart rate in 4 dpf zebrafish (n=68). Data shown as mean +/- SD.. *p<0.05, **p<0.001, Friedman test and Dunn's multiple comparisons.





